



Nonsteroidal Anti-Inflammatory Drugs (NSAIDs) Therapeutic Class Review (TCR)

June 17, 2020

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, digital scanning, or via any information storage or retrieval system without the express written consent of Magellan Rx Management.

All requests for permission should be mailed to:

Magellan Rx Management
Attention: Legal Department
6950 Columbia Gateway Drive
Columbia, Maryland 21046

The materials contained herein represent the opinions of the collective authors and editors and should not be construed to be the official representation of any professional organization or group, any state Pharmacy and Therapeutics committee, any state Medicaid Agency, or any other clinical committee. This material is not intended to be relied upon as medical advice for specific medical cases and nothing contained herein should be relied upon by any patient, medical professional or layperson seeking information about a specific course of treatment for a specific medical condition. All readers of this material are responsible for independently obtaining medical advice and guidance from their own physician and/or other medical professional in regard to the best course of treatment for their specific medical condition. This publication, inclusive of all forms contained herein, is intended to be educational in nature and is intended to be used for informational purposes only. Send comments and suggestions to PSTCREditor@magellanhealth.com.

June 2020

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.
© 2004-2020 Magellan Rx Management. All Rights Reserved.

MagellanRx
MANAGEMENTSM

FDA-APPROVED INDICATIONS

Oral NSAIDs

Drug	Mfg	OA	RA	JIA	AS	Pain	PD	Other
Single Ingredient Agents								
celecoxib (Celebrex®) ¹	generic, Pfizer	X	X	X	X	X	X	
diclofenac potassium ²	generic	X	X			X	X	
diclofenac potassium (Zipsor®) ³	Assertio					X		
diclofenac sodium (Voltaren®/XR®) ^{4,5}	generic	X	X		X (IR)			
diclofenac submicronized (Zorvolex®) ⁶	generic, Iroko/Egalet/Zyla	X				X		
diflunisal ⁷	generic	X	X			X		
etodolac (Lodine®) ⁸	generic	X	X	X		X		
fenoprofen (Nalfon®) ⁹	generic, Xspire	X	X			X		
flurbiprofen ¹⁰	generic	X	X					
ibuprofen (Motrin®) ¹¹	generic	X	X			X	X	
indomethacin (Indocin®) ¹²	generic, Iroko/Egalet	X	X		X			Treatment of painful shoulder (tendonitis, bursitis) and acute gout
Indomethacin submicronized (Tivorbex®) ¹³	generic, Iroko/Egalet					X		
ketoprofen IR ¹⁴	generic	X	X			X	X	
ketoprofen ER ¹⁵	generic	X	X					

OA = Osteoarthritis, RA = Rheumatoid Arthritis, JIA = Juvenile Idiopathic Arthritis (a.k.a. Juvenile Rheumatoid Arthritis [JRA]), AS = Ankylosing Spondylitis, PD = Primary Dysmenorrhea

FDA-Approved Indications: Oral NSAIDs (continued)

Drug	Mfg	OA	RA	JIA	AS	Pain	PD	Other
Single Ingredient Agents (continued)								
ketorolac tromethamine ¹⁶	generic					X		Short-term (≤ 5 days) management of moderately severe acute pain that requires analgesia at the opioid level, usually in a postoperative setting. Therapy should always be initiated with ketorolac tromethamine injectable formulation (IM/IV) and ketorolac tromethamine tablets are to be used only as continuation treatment, if necessary
meclofenamate ¹⁷	generic	X	X	X	X	X	X	Treatment of primary dysmenorrhea/excessive blood loss Treatment of acute painful shoulder and acute gouty arthritis
mefenamic acid ¹⁸	generic					X < 1 week	X	
meloxicam (Mobic®) ¹⁹	generic, Boehringer- Ingelheim	X	X	X				
meloxicam (Qmiiiz™ ODT) ²⁰	TerSera	X	X	X				
meloxicam submicronized (Vivlodex™) ²¹	Iroko/Egalet, Egalet/Zyla	X						
nabumetone § (Relafen DS™) ^{22,23}	generic, Carwin	X	X					
naproxen (Anaprox® / DS, Naprelan®, EC- / Naprosyn®) ^{24,25}	generic	X	X	X	X	X	X	Treatment of tendonitis, bursitis, and acute gout
oxaprozin (Daypro®) ²⁶	generic, Pfizer	X	X	X				
piroxicam (Feldene®) ²⁷	generic, Pfizer	X	X					
sulindac ²⁸	generic	X	X		X			Treatment of acute painful shoulder and acute gouty arthritis
tolmetin ²⁹	generic	X	X	X				

OA = Osteoarthritis, RA = Rheumatoid Arthritis, JIA = Juvenile Idiopathic Arthritis (a.k.a. Juvenile Rheumatoid Arthritis [JRA]), AS = Ankylosing Spondylitis, PD = Primary Dysmenorrhea

§ Relafen DS is approved under an abbreviated new drug application (ANDA) as a generic³⁰ and provides a new strength (1,000 mg) of nabumetone tablets.

FDA-Approved Indications: Oral NSAIDs (continued)

Drug	Mfg	OA	RA	JIA	AS	Pain	PD	Other
Combination Agents								
celecoxib/ methyl salicylate/ menthol/ capsaicin [¶] (NuDroxiPak [™]) ³¹	NuCare	X	X	X	X	X	X	
diclofenac sodium DR/ capsaicin (Flexipak, Inavix, Inflammacin [™] , Nudiclo [™] TabPak, Xenaflamm [™]) ^{32,33,34,35,36}	Sterling-Knight, Primary, PureTek, Nucare, Shoreline	X	X		X			
diclofenac sodium DR/ capsaicin/ methyl salicylate/menthol [¶] (NuDroxiPak DSDR-50, NuDroxiPak DSDR-75) ³⁷	NuCare	X	X		X			
diclofenac sodium DR/ menthol / camphor ^{**} (Inflatherm [™]) ³⁸	V2	X	X		X			
diclofenac sodium/ misoprostol (Arthrotec [®]) ³⁹	generic, Pfizer	X	X					Indicated for patients who are at high risk for NSAID-induced GI ulcers
etodolac/ capsaicin/ methyl salicylate/ menthol [§] (NuDroxiPak E-400) ⁴⁰	NuCare	X	X			X		

OA = Osteoarthritis, RA = Rheumatoid Arthritis, JIA = Juvenile Idiopathic Arthritis (a.k.a. Juvenile Rheumatoid Arthritis [JRA]), AS = Ankylosing Spondylitis, PD = Primary Dysmenorrhea

¶ NuDroxiPak products contain an oral NSAID co-packaged with topical NuDroxin[®] Pain Relief Roll-on solution that contains capsaicin, menthol, and methyl salicylate for the temporary relief of minor aches and muscle pain associated with arthritis, simple backache, strains, muscle soreness, and stiffness

|| Diclofenac sodium DR/capsaicin kits (Flexipak, Inflammacin, Nudiclo TabPak, Xenaflamm) contain an oral NSAID co-packaged with topical capsaicin cream for the temporary relief of temporary relief of minor aches and muscle pain associated with arthritis, simple backache, strains, and sprains.

** Inflatherm contains an oral NSAID co-packaged with Sore No More roll-on gel (diclofenac/menthol/camphor) that contains menthol and camphor or Asper-Flex cream (diclofenac/trolamine salicylate) that contains trolamine salicylate for the temporary relief of minor aches and pains of muscles and joints associated with simple backaches and arthritis.

FDA-Approved Indications: Oral NSAIDs (continued)

Drug	Mfg	OA	RA	JIA	AS	Pain	PD	Other
Combination Agents (continued)								
esomeprazole/ naproxen (Vimovo®) ⁴¹	generic*, Horizon	X	X		X			Indicated for the relief of signs and symptoms of osteoarthritis, rheumatoid arthritis, and ankylosing spondylitis and the reduction of risk of stomach (gastric) ulcers in patients at risk of developing stomach ulcers due to treatment with NSAIDs Indicated for juvenile idiopathic arthritis (JIA) in adolescent patients ≥ 12 years of age weighing at least 38 kg requiring naproxen for symptomatic relief or arthritis and reduction of risk developing stomach ulcers due to treatment with NSAIDs
ibuprofen/ famotidine (Duexis®) ⁴²	Horizon	X	X					Indicated for the relief of signs and symptoms of rheumatoid arthritis and osteoarthritis and to decrease the risk of developing upper gastrointestinal ulcers
ibuprofen/ capsaicin/ menthol/ methyl salicylate (Comfort Pac™ with Ibuprofen ^{¶¶} ; NuDroxiPak I-800 ^{¶¶}) ^{43,44}	PD-Rx, NuCare	X	X			X	X	
meloxicam/ capsaicin/ menthol/ methyl salicylate (Comfort Pac with Meloxicam ^{¶¶}) ⁴⁵	PD-Rx	X	X	X				
naproxen/ capsaicin/ menthol/ methyl salicylate (Comfort Pac with Naproxen ^{¶¶}) ⁴⁶	PD-Rx	X	X	X	X	X	X	Treatment of tendonitis, bursitis, and acute gout
nabumetone/ capsaicin/ menthol/ methyl salicylate [¶] (NuDroxiPak N-500) ⁴⁷	NuCare	X	X					

OA = Osteoarthritis, RA = Rheumatoid Arthritis, JIA = Juvenile Idiopathic Arthritis (a.k.a. Juvenile Rheumatoid Arthritis [JRA]), AS = Ankylosing Spondylitis, PD = Primary Dysmenorrhea

* The generic version of Vimovo® is indicated for use only in adults, as the brand-name carries marketing exclusivity rights for the pediatric use.

¶ NuDroxiPak products contain an oral NSAID co-packaged with topical NuDroxicin® Pain Relief Roll-on solution that contains capsaicin, menthol, and methyl salicylate for the temporary relief of minor aches and muscle pain associated with arthritis, simple backache, strains, muscle soreness, and stiffness

¶¶ The Comfort Pacs contain an oral NSAID product co-packaged with topical DuraFlex Comfort Gel containing capsaicin, menthol, and methyl salicylate for topical pain relief.

FDA-Approved Indications: Nasal NSAIDs

Drug	Mfg	OA	RA	JIA	AS	Pain	PD	Other
ketorolac tromethamine (Sprix®) ⁴⁸	generic*, Egalet					X		Short-term (up to 5 days) management of moderate to moderately severe pain that requires analgesia at the opioid level

OA = Osteoarthritis, RA = Rheumatoid Arthritis, JIA = Juvenile Idiopathic Arthritis (a.k.a. Juvenile Rheumatoid Arthritis [JRA]), AS = Ankylosing Spondylitis, PD = Primary Dysmenorrhea

*Authorized generic (AG) available

FDA-Approved Indications: Topical NSAIDs

Drug	Mfg	OA	RA	JIA	AS	Pain	PD	Other
diclofenac epolamine (Flector®; Licart™) ^{49,50}	generic*, Pfizer, IBSA					X		Topical treatment of acute pain due to minor strains, sprains, and contusions in patients ≥ 6 years old (Licart labeled for adult use only)
diclofenac sodium (Lexixryl™, Pennsaid® 2% pump, Xrylix™) ^{51,52,53,54}	generic, Shoreline, Horizon, PureTek	X						Treatment of signs and symptoms of osteoarthritis of the knee(s)
diclofenac sodium (Voltaren® Gel, Diclozor) ^{55,56}	generic, Endo, Sterling Knight	X						Relief of pain of osteoarthritis of joints amenable to topical treatment, such as of the knees and hands
diclofenac sodium/ camphor/ lidocaine/ methyl salicylate (Diclovix) ⁵⁷	Primary	X						Treatment of signs and symptoms of osteoarthritis of the knee(s)
diclofenac sodium/ camphor/ menthol (Diclotrex™) ⁵⁸	Puretek	X						Treatment of signs and symptoms of osteoarthritis of the knee(s)
diclofenac sodium/ camphor/ menthol/ methyl salicylate (Inflamma-K Kit) ⁵⁹	Solutech	X						Treatment of signs and symptoms of osteoarthritis of the knee(s)

OA = Osteoarthritis, RA = Rheumatoid Arthritis, JIA = Juvenile Idiopathic Arthritis (a.k.a. Juvenile Rheumatoid Arthritis [JRA]), AS = Ankylosing Spondylitis, PD = Primary Dysmenorrhea

*Authorized generic (AG) available

FDA-Approved Indications: Topical NSAIDs (continued)

Drug	Mfg	OA	RA	JIA	AS	Pain	PD	Other
diclofenac sodium/ capsaicin (CapsFenac Pak, Capsinac, DermacinRx® Lexitral™ PharmaPak, Diclofex DC, Diclopak, Diclosiacin, Diclotral Pak, Nudiclo™ Solupak, Sure Result DSS Premium Pak, Xelitral™) ^{60,61,62,63,64,65,66,67,68,69}	SA3, Bodyshpere, PureTek, Sircle, Sterling-Knight, Chadwick, Patchwerx, Nucare, International Brand Management, Shoreline	X						Treatment of signs and symptoms of osteoarthritis of the knee(s)
diclofenac sodium/ lidocaine/ tape (Trixylitral™) ⁷⁰	Shoreline	X						Treatment of signs and symptoms of osteoarthritis of the knee(s)
diclofenac sodium/ menthol (Dithol Kit, Dimentho) ^{71,72}	Fortus, Levins	X						Treatment of signs and symptoms of osteoarthritis of the knee(s)
diclofenac sodium/ menthol/ tape (Diclovix M) ⁷³	Primary	X						Treatment of signs and symptoms of osteoarthritis of the knee(s)
diclofenac sodium/ menthol/ methyl salicylate (DicloPR) ⁷⁴	19&Pacific	X						Relief of pain of osteoarthritis of joints amenable to topical treatment, such as of the knees and hands
diclofenac sodium/ menthol/ methyl salicylate (Varophen) ⁷⁵	V2	X						Treatment of signs and symptoms of osteoarthritis of the knee(s)

OA = Osteoarthritis, RA = Rheumatoid Arthritis, JIA = Juvenile Idiopathic Arthritis (a.k.a. Juvenile Rheumatoid Arthritis [JRA]), AS = Ankylosing Spondylitis, PD = Primary Dysmenorrhea

Products packaged as kits are discussed in the FDA Indications and Dosages sections, and components are addressed in the Pharmacology section. For other information regarding these products, refer to the primary component throughout the text (e.g., celecoxib, diclofenac, naproxen).

Cambia® (diclofenac potassium powder for oral solution) is only indicated for the acute treatment of migraine attacks in adult patients. It will not be discussed in this therapeutic class review.

OVERVIEW

Nonsteroidal anti-inflammatory drugs (NSAIDs) are commonly used to treat rheumatoid arthritis (RA), osteoarthritis (OA), and pain from various etiologies. NSAIDs are among the most widely used drugs in the United States (US), with approximately 70 million prescriptions being filled yearly; when over-the-counter (OTC) utilization is included, over 30 billion doses of NSAIDs are taken each year in the US.⁷⁶ It is estimated that OTC NSAIDs are used 5 to 7 times more often than prescription NSAIDs.^{77,78} Most oral NSAIDs are now available as generics and are generally considered to be safe and effective.

NSAIDs are associated with adverse effects including gastrointestinal (GI) bleeding, peptic ulcer disease, hypertension, edema, and renal disease. In addition, NSAIDs have been linked to an increased risk of myocardial infarction (MI) which is reflected in the boxed warning for all NSAIDs. In July 2015, the Food and Drug Administration (FDA) issued a Safety Alert strengthening the existing warning on the increased risk of heart attack and stroke risk associated with NSAIDs.⁷⁹

GI adverse effects induced by NSAIDs lead to significant morbidity and mortality. Ulcers are found by endoscopy in 15% to 30% of patients who are using NSAIDs regularly, and the incidence of upper GI clinical events due to NSAIDs is 2.5% to 4.5%. In the US, GI side effects due to NSAIDs in patients with arthritis account for approximately 107,000 hospitalizations and result in 16,500 deaths each year.⁸⁰ Products designed to lessen NSAID GI adverse reactions (Arthrotec, Vimovo, and Duexis) are available.

Celecoxib (Celebrex), which selectively inhibits the cyclooxygenase-2 (COX-2) enzyme, has equal efficacy to many of the other NSAIDs, but the issue of a better safety profile is unclear. Rofecoxib (September 2004) and valdecoxib (April 2005) have been removed from the market due to safety concerns. Celecoxib and all nonselective NSAIDs have come under greater scrutiny due to concerns over their cardiovascular (CV) safety.

Drug delivery technology may overcome the disadvantages of oral drug administration. First pass metabolism may impact oral administration and has the potential for systemic adverse effects.⁸¹ A route of administration that bypasses the systemic exposure would provide an alternative that might improve patient adherence, minimize adverse effects, allow for a longer treatment interval, and serve as a substitute to conventional therapy.

NSAIDs reduce swelling and ease inflammation that can cause pain. NSAIDs are commonly used to treat osteoarthritis and pain from different etiologies. When pharmacological treatment is required, the 2019 American College of Rheumatology (ACR) clinical practice guidelines strongly recommend oral NSAIDs for hand, knee, and hip OA; topical NSAIDs are strongly recommended for knee OA and conditionally recommended for hand OA.⁸² The 2013 treatment guidelines from the American Academy of Orthopedic Surgeons for the treatment of osteoarthritis of the knee do not specify a specific NSAID or route of administration for osteoarthritis symptoms.⁸³ If the risk of GI adverse events is increased, the topical route is preferred among other treatment strategies. Similarly, the American Academy of Orthopedic Surgeons clinical practice guidelines (2017) on the management of osteoarthritis of the hip does not specify a specific NSAID; however, they do note strong evidence to support NSAIDs to improve short-term pain, function, or both in patients with symptomatic osteoarthritis of the hip.⁸⁴ The 2017 American College of Physicians (ACP) clinical practice guidelines for noninvasive treatments for acute, subacute, and chronic low back pain recommend NSAIDs for patients with chronic low back pain as pharmacological therapy for patients who have not had

sufficient relief from nonpharmacological treatments; tramadol or duloxetine are considered 2nd-line therapy in this setting. Opioids for chronic low back pain should only be considered if prior therapy fails and the potential benefits outweigh risks for the patient.⁸⁵

In addition, ACP published gout guidelines in 2016 recommending corticosteroids, NSAIDs, and colchicine to treat patients with gout flare.⁸⁶ Use of NSAIDs is based on high-quality evidence demonstrating their ability to reduce pain in this patient population. Similar efficacy of among NSAIDs (including selective agents) and between NSAIDs and corticosteroids has been demonstrated. Additionally, NSAIDs for prophylactic therapy can also reduce the risk for gout flare in patients starting urate-lowering therapy. Similarly, the 2020 ACR guideline on the management of gout strongly recommends oral colchicine, NSAIDs, or glucocorticoids (oral, intraarticular, or intramuscular) as 1st-line treatment for gout flares over other agents (e.g., interleukin-1 inhibitors, adrenocorticotrophic hormone). Patient factors should determine which agent (colchicine, NSAIDs, glucocorticoids) is used.⁸⁷

PHARMACOLOGY

Both oral and topical NSAIDs inhibit the cyclooxygenase-1 (COX-1) and cyclooxygenase-2 (COX-2) enzymes that catalyze the synthesis of prostaglandins from arachidonic acid. These prostaglandins are partially responsible for the development of pain and inflammation associated with various medical conditions. COX-1 plays a role in maintaining normal gastric mucosa and influences kidney function. COX-2 activity is rapidly upregulated during inflammatory pain conditions and may be involved in the pathogenesis of some malignancies.^{88,89,90,91,92} Selective COX-2 inhibitors provide anti-inflammatory effects and analgesia while theoretically resulting in fewer adverse effects than the nonselective NSAIDs. However, other prostaglandins may be affected that alter platelet aggregation, affecting the CV risk with some of the NSAIDs.

Zorvolex capsules contain diclofenac free acid, whereas other diclofenac products contain a salt of diclofenac (e.g., diclofenac potassium or sodium). The reduction in diclofenac particle size with Zorvolex increases surface area, leading to faster dissolution and absorption of the drug.

Vivlodex capsules contain submicronized meloxicam particles which allow a faster dissolution, resulting in an earlier time to maximum peak concentration compared to the tablet (approximately 2 hours versus 4 hours, respectively).⁹³

The inhibition of platelet aggregation seen with NSAIDs is due to the inhibition of COX-1 in platelets, causing decreased levels of platelet thromboxane A₂ and increased bleeding time. The inhibition of platelet aggregation is reversible.

Esomeprazole, a component of Vimovo, is a proton pump inhibitor (PPI) that works by blocking H⁺/K⁺-ATPase in gastric parietal cells, resulting in suppression of gastric acid secretions. This activity is dose-dependent.⁹⁴

Misoprostol, a component of Arthrotec, is a synthetic prostaglandin E₁ analog. This agent counteracts the inhibition of prostaglandin synthesis noted with NSAIDs, increasing bicarbonate and mucus production. At doses of 200 mcg or greater, misoprostol is also noted to have significant anti-secretory properties, making the exact nature of its gastroprotective properties unclear.⁹⁵

Famotidine, a component of Duexis, is a competitive inhibitor of histamine-2 receptors, which thereby suppresses both the acid concentration and volume of gastric secretion.⁹⁶ Changes in pepsin secretion are proportional to volume output.

In peripheral sensory neurons, topical capsaicin depletes and limits reaccumulation of substance P, which is thought to be a primary mediator of pain impulses to the central nervous system (CNS). Thus, it limits local pain sensations as these sensations cannot be transmitted to the brain. Topical menthol also has a local effect, including mild analgesia, a cooling sensation, and an irritant/counter-irritant effect. Menthol also causes vasodilation. There are multiple mechanisms of menthol theorized on a cellular level; however, the exact mechanism of its effects is not well defined.⁹⁷ Methyl salicylate acts similarly to menthol and functions as a counterirritant.⁹⁸

The following chart indicates the location of COX-1 and COX-2 enzymes in the body:^{99,100,101}

Location	Brain	Breast Cancer	Colorectal Adenomas, Carcinomas	Endothelial Cells	GI Tract	Head and Neck Cancer	Liver	Lung	Platelets	Renal Cortex, Medullary Interstitial Cells	Renal Medullary Collecting Ducts, Interstitium	Site of Inflammation	Spleen	Synovial Tissue
COX-1				X	X		X	X	X		X		X	
COX-2	X	X	X			X				X		X		X

PHARMACOKINETICS ^{102,103,104,105,106,107,108,109,110,111,112,113,114,115,116,117,118,119,120,121,122,123,124,125,126,127,128,129,130,131,132,133,134,135,136,137,138,139,140,141,142,143,144,145,146,147}

Oral NSAIDs: Drug	Bioavailability (%)	Time to Peak (hr)	Half-Life (hr)	Excretion (%)	
Single Ingredient Agents					
celecoxib (Celebrex)	nr	3	11	Renal: 27 Feces: 57	
diclofenac submicronized (Zorvolex)	50	1	1–2	Renal: 65 Bile: 35	
diclofenac potassium	50	1	1–2	Renal: 65 Bile: 35	
diclofenac potassium (Zipsor)		0.47			
diclofenac sodium (Voltaren)		2.3			2
diclofenac sodium XR/DR (Voltaren XR)		5.3			2
diflunisal	100	2–3	8–12	90-urine	
etodolac (Lodine)	≥ 80	1.4–6.7	6.4–8.4	Renal: 72 Feces: 16	
fenoprofen (Nalfon)	nr	2	3	Renal: 90	
flurbiprofen	96	1.9	7.5	Renal: 70	

Pharmacokinetics (continued)

Oral NSAIDs: Drug	Bioavailability (%)	Time to Peak (hr)	Half-Life (hr)	Excretion (%)
Single Ingredient Agents (continued)				
ibuprofen (Motrin)	< 80	1-2	1.8–2	Renal: 45–79
indomethacin (Indocin)	98	2	4.5	Renal: 60 Feces: 33
indomethacin (Tivorbex)	~ 100	1.67	7.6	Renal: 60 Feces: 33
ketoprofen	90	0.5–7	2–5.4	Renal: 80
ketorolac	100	0.75	2.5–5	Renal: 92 Feces: 6
meclofenamate	~ 100	0.5–2	0.8-5.3	Renal: 70 Feces: 30
mefenamic acid	nr	2–4	2	Renal: 52 Feces: 20
meloxicam (Mobic, Qmiiz ODT)	89	4-5	15–20	Renal: 50 Feces: 50
meloxicam submicronized (Vivlodex)	nr	2	22	Renal: 50 Feces: 50
nabumetone (Relafen DS)	> 80	2.5–4	24	Renal: 80 Feces: 9
naproxen (Anaprox / DS, Naprelan, EC/ Naprosyn)	95	1-6	12–17	Renal: 95
oxaprozin (Daypro)	95	2.5–3	41–55	Renal: 65 Feces: 35
piroxicam (Feldene)	nr	3–5	50	Renal: 95 Feces: 5
sulindac	90	3–4	7.8	Renal: 50 Feces: 25
tolmetin	99	0.5–1	5	Renal: 99
Combination Agents				
diclofenac sodium/ misoprostol (Arthrotec)	50	2	2	Renal: 65 Feces: 35
	nr	0.33	0.5	Renal: 70
esomeprazole/ naproxen (Vimovo)	nr	0.43–1.2	1	Renal: 80
	95	3	15	Renal: 95 Feces: 3
ibuprofen/ famotidine (Duexis)	< 80	1.9	2	Renal: 45–79
	nr	2	4	Renal: 65–70 Metabolic: 30–35

nr = not reported

Nasal NSAIDs: Drug	Bioavailability (%)	Time to Peak (hr)	Half-Life (hr)	Excretion (%)
ketorolac nasal spray (Sprix)	60	0.75	2.5–6	Renal: 92 Feces: 6

Topical NSAIDs

Following a single application of diclofenac epolamine (Flector, Licart) to the upper inner arm, the peak plasma concentrations were noted within 10 to 20 hours. Diclofenac epolamine is 99% protein bound. Diclofenac sodium (Voltaren Gel, diclofenac gel component of co-packaged products) has 17 times less systemic exposure than the orally administered diclofenac. The amount of diclofenac sodium that is absorbed is on average 6% of that from oral diclofenac. Diclofenac sodium (Pennsaid) results in about one-third of the systemic exposure compared to a topical diclofenac gel. The elimination half-life for topical diclofenac is approximately 12 hours. Diclofenac is metabolized through glucuronidation and eliminated through subsequent urinary and biliary excretion.

CONTRAINDICATIONS/WARNINGS ^{148,149,150,151,152,153,154,155,156,157,158,159,160,161,162,163,164,165,166,167,168,169,170,171,172,173,174,175,176,177,178,179,180,181,182,183,184,185,186,187,188,189,190,191,192}

Oral NSAIDs

NSAIDs (non-selective and selective) should not be given to patients who have experienced asthma, urticaria, or allergic-type reactions after taking aspirin, other NSAIDs, or sulfonamides. Severe, rarely fatal, anaphylactic-like reactions to NSAIDs have been reported in such patients.

Ibuprofen/famotidine (Duexis) is contraindicated in patients in late stages of pregnancy. This agent should not be used in patients with a known hypersensitivity to a histamine 2 receptor antagonist (H₂RA).

Diclofenac sodium/misoprostol (Arthrotec) is contraindicated in patients who are pregnant because misoprostol can cause abortion, premature birth, or birth defects. This agent should also not be used in women of childbearing potential unless the benefits clearly outweigh the risks of therapy.

Borderline elevations (< 3 times the upper limit of the normal [ULN] range) or greater elevations of transaminases occurred in about 15% of diclofenac-treated patients in clinical trials of indications other than acute pain. Alanine transaminase (ALT) should be monitored to detect liver injury.

Long-term PPI therapy is associated with an increased risk for osteoporosis-related fractures of the hip, wrist, or spine.¹⁹³ The risk of fracture was increased in patients who received multiple daily doses for a year or longer. Esomeprazole/naproxen (Vimovo) is approved for use twice a day and does not allow for administration of a lower daily dose of the PPI.

Hypomagnesemia has been reported rarely in patients treated with PPIs for at least 3 months, in most cases after a year of therapy. Serious adverse events include tetany, arrhythmias, and seizures.¹⁹⁴

In addition, PPI use may be associated with an increased risk of *Clostridium difficile*-associated diarrhea (CDAD).¹⁹⁵ It is unknown if patients using a H₂RA, such as famotidine, are at increased risk of CDAD.

Acute interstitial nephritis may occur at any time during PPI therapy and has been observed in patients taking PPIs. Esomeprazole/naproxen should be discontinued if interstitial nephritis occurs.

Malabsorption of cyanocobalamin (vitamin B-12) due to hypo-achlorhydria may occur for patients taking medications that suppress acid (PPIs) for longer than 3 months. Although rare, a diagnosis of cyanocobalamin deficiency should be considered if clinical symptoms present.

Cardiovascular (CV) Concerns

NSAIDs may cause an increased risk of serious CV thrombotic events, MI, and stroke, which can be fatal. This risk may increase with duration of use. Patients with CV disease or risk factors for CV disease may be at greater risk. In April 2005, the FDA asked the manufacturers of all marketed prescription NSAIDs (non-selective and COX-2 selective), to revise the labeling for the products to include a boxed warning stating that NSAIDs may cause an increased risk of potentially fatal CV thrombotic events, MI, and stroke. All NSAIDs may have a similar risk, which increases with longer duration of use.¹⁹⁶ Patients with CV disease or CV risk factors may be at greater risk. All NSAIDs are contraindicated for the treatment of peri-operative pain in the setting of coronary artery bypass graft (CABG) surgery. NSAIDs cause an increased risk of potentially fatal bleeding, ulceration, and perforation of the stomach or intestines, occurring at any time during use and without warning. Elderly patients are at greater risk for serious GI events. In July 2015, the FDA issued a Safety Alert strengthening the existing warning on the increased risk of heart attack and stroke risk associated with NSAIDs.¹⁹⁷

In an August 2001 review of the COX-2 inhibitors and risk of CV events, the authors concluded that a prospective trial may be necessary to evaluate the potential risk of CV events with these agents. CLASS and VIGOR studies were designed to examine the GI effects of these medications, not the CV safety.^{198,199,200} However, the VIGOR data were particularly concerning, since it showed a higher incidence of MI in rofecoxib (Vioxx[®]) patients compared to naproxen patients. The implications of these data were unclear because it was unknown if naproxen had a cardioprotective effect or if rofecoxib had adverse effects on the CV system. Data became available that brought the COX-2 inhibitors under more scrutiny; several retrospective studies and meta-analyses questioned the safety of these products.^{201,202,203,204} Newer studies evaluating these agents for use for other indications had more stringent monitoring in place for CV problems, which proved to be a critical step in evaluating their true effects.²⁰⁵

Rofecoxib was withdrawn from the market after the discovery of higher CV risk with the agent in the APPROVe study. Not long after the withdrawal of rofecoxib, the CV effects of the other COX-2 inhibitors were called into question. Valdecoxib (Bextra[®]) was withdrawn from the market in February 2005 following extensive study of available clinical trials by the FDA. Celecoxib was allowed to remain on the market, but the advisement was given to use celecoxib at the lowest effective dose.^{206,207}

The American Heart Association recommended soon afterward that any COX-2 inhibitors be reserved in patients with a history or risk of GI bleeding unless potential benefits of treatment are felt to outweigh the potential CV risks or nonselective NSAID therapy is insufficient.²⁰⁸ The 2007 update to the American Heart Association Scientific Statement on the use of NSAIDs reiterates the reservation of COX-2 inhibitor use in patients with history of or at risk of CV disease and strengthens the point by

suggesting COX-2 selective agents be used as a last resort with the prior steps being non-pharmacologic treatments (physical therapy, weight loss, exercise, etc.) followed by a stepped pharmacologic approach.²⁰⁹ The first-line agents recommended are acetaminophen, aspirin, or a short-term narcotic analgesic. If therapeutic alternatives are needed, physicians should consider the nonselective NSAIDs prior to the selective COX-2 inhibitor. This stepped approach focuses on the reported risk of CV events with the need for assessment of risk/benefit ratio at each step.

NSAIDs can lead to the onset of hypertension or worsening of existing hypertension. In addition, fluid retention and edema have been observed in some patients taking NSAIDs. Also, patients taking NSAIDs may have a decreased response to thiazide or loop diuretics. Therefore, monitoring patients with hypertension and patients at risk for the development of edema is recommended with all NSAIDs.

GI Toxicity

NSAIDs can cause an increased risk of serious GI adverse effects including bleeding, ulceration, and perforation of the stomach or intestine, which can be fatal. Serious GI toxicity can occur with or without warning with all NSAIDs. Patients at higher risk for the development of GI toxicity include patients on corticosteroids, anticoagulants, or long duration of NSAID therapy, as well as those with the following risk factors: smoking, alcoholism, poor health status, and older age. NSAIDs can exacerbate inflammatory bowel disease and should therefore be given with caution to patients with a history of this condition.

Renal Toxicity

Long-term use of NSAIDs can lead to renal papillary necrosis, renal insufficiency, acute renal failure, and other renal injuries. Patients with impaired renal function, heart failure, liver dysfunction, the elderly, and those taking diuretics, angiotensin-converting enzyme (ACE) inhibitors, and angiotensin receptor blockers are at greatest risk for this reaction. The FDA notes that discontinuation of NSAID therapy usually is followed by recovery to the pretreatment state.

CNS adverse effects, including seizures, delirium, and coma, have been reported with famotidine in patients with moderate (creatinine clearance [CrCl] < 50 mL/min) and severe (CrCl < 10 mL/min) renal impairment. Since the dosage of the famotidine component in Duexis is fixed, this product is not recommended in patients with moderate to severe renal insufficiency.

Naproxen/esomeprazole (Vimovo) is not recommended in patients with moderate or severe renal impairment.

Safety and efficacy of submicronized meloxicam (Vivlodex) has not been established in patients with severe renal impairment; use is not recommended. The maximum dose in patients on hemodialysis is 5 mg.

Hepatic Impairment

Elevations of 1 or more liver tests may occur in up to 15% of patients taking NSAIDs. Elevations of up to 3 X ULN of alanine aminotransferase (ALT) or aspartate aminotransferase (AST) have been reported in about 1% of patients in clinical trials with NSAIDs. In addition, rare cases of severe hepatic reactions, including jaundice and fatal fulminant hepatitis, liver necrosis, and hepatic failure, some of them with fatal outcomes, have been reported.

Naproxen/esomeprazole (Vimovo) is not recommended in patients with severe hepatic impairment because esomeprazole doses should not exceed 20 mg daily in these patients.

Skin Reactions

NSAIDs can cause serious skin adverse events, such as exfoliative dermatitis, Stevens-Johnson syndrome, and toxic epidermal necrosis, which can be fatal. These conditions can occur without warning. NSAID therapy should be stopped at the first appearance of skin rash or other sign of hypersensitivity.

Hematological

NSAID therapy should be stopped if active and clinically significant bleeding from any source occurs. Anemia is sometimes seen in patients taking NSAIDs. This may be due to fluid retention, occult or gross GI blood loss, or an effect upon erythropoiesis. Patients with initial hemoglobin values of 10 g/dL or less who are to receive long-term therapy with NSAIDs/NSAID combinations should have hemoglobin values assessed periodically.

Pre-existing Asthma

The use of aspirin in patients with aspirin-sensitive asthma has been associated with severe and possibly fatal bronchospasm. Since cross-reactivity between aspirin and the various NSAIDs has been reported in such aspirin-sensitive patients, oral and topical NSAIDs should not be administered to patients with this form of aspirin sensitivity and should be used with caution in patients with pre-existing asthma.

Visual Disturbances

Blurred and/or diminished vision, scotomata, and/or changes in color vision have been reported with ibuprofen/famotidine (Duexis). If a patient develops such complaints while on therapy, the drug should be discontinued, and the patient should have an ophthalmologic examination which includes central visual fields and color vision testing.

Nasal NSAIDs

Ketorolac tromethamine nasal spray (Sprix) is contraindicated in patients with a known hypersensitivity to ethylenediaminetetraacetic acid. Ketorolac nasal spray is contraindicated for use as a prophylactic analgesic before any major surgery. Probenecid decreases the clearance of ketorolac nasal spray; concomitant use is contraindicated. Similarly, the combination of pentoxifylline with ketorolac nasal spray is contraindicated due to increased bleeding risk.

Topical NSAIDs

Diclofenac formulations (Flector, Licart, Pennsaid, Voltaren Gel, diclofenac components of co-packaged products) carry a boxed warning for CV and gastrointestinal risk. These agents may cause an increased risk of serious CV thrombotic events, MI, and stroke, which can be fatal. Formulations are also contraindicated for the treatment of perioperative pain in the setting of coronary artery bypass graft (CABG) surgery. These should not be applied to damaged skin or skin that is not intact.

Patients should be informed of the potential for adverse CV effects associated with all NSAIDs (e.g., risk of CV thrombotic events, new onset or worsening of hypertension, congestive heart failure, and edema). Diclofenac formulations should be used cautiously in patients with these conditions.

NSAIDs, including diclofenac formulations, can cause serious GI adverse events, including inflammation, ulceration, and bleeding and perforation of the stomach, small intestine, or large intestine, which can be fatal. These events can occur at any time during use and without warning symptoms. Elderly patients are at greater risk for serious GI events.

Diclofenac formulations should not be given to patients who have experienced asthma, urticaria, or allergic-type reactions after taking aspirin or other NSAIDs. They should also be avoided in patients with the aspirin triad (a nasal symptom complex typically occurring in asthmatic patients who experience rhinitis with or without nasal polyps or who have severe, potentially fatal bronchospasm after taking aspirin or other NSAIDs).

Capsaicin, copackaged with multiple kit products, is an irritant and should be used with caution to avoid irritation to mucous membranes or the eyes. Menthol/capsaicin patch should not be used on wounds or damaged skin. It should also not be used in combination with any bandage, wrap, or other similar garment. Menthol/methyl salicylate, with or without camphor, should not be used on wounds or damaged skin; contact with eyes, mucous membranes, or rashes should also be avoided.

Medication Guide/Risk Evaluation and Mitigation Strategy (REMS)

A Medication Guide must accompany every prescription NSAID, except for diclofenac potassium 25 mg capsules (Zipsor), at the time of dispensing to better inform patients of possible adverse effects. In June 2005, the FDA requested that the manufacturers of OTC NSAIDs revise their labeling to include more specific information about the potential GI and CV risks. In July 2015, the FDA issued a Safety Alert strengthening the existing warning on the increased risk of heart attack and stroke associated with NSAIDs.²¹⁰ As a result, the FDA is requiring updates to all prescription labels and requesting updates to over-the-counter (OTC) Drug Facts labels.

DRUG INTERACTIONS^{211,212,213,214,215,216,217,218,219,220,221,222,223,224,225,226,227,228,229,230,231,232,233,234,235,236,237,238,239,240,241,242,243,244,245,246,247,248,249,250,251,252,253,254}

Oral NSAIDs

ACE inhibitors/ARBs, or beta-blockers

NSAIDs may diminish the antihypertensive effect of angiotensin converting enzyme (ACE) inhibitors, angiotensin receptor blockers (ARBs), or beta blockers (including propranolol). Deterioration of renal function, including possible acute renal failure may occur with the coadministration of an NSAID with an ACE inhibitor or ARB in elderly patients, those volume depleted or in those with renal impairment. These effects are usually reversible.

aspirin

Concomitant use of aspirin and NSAIDs is not generally recommended because of the potential for GI ulceration. NSAID therapy is not a substitute for aspirin for CV prophylaxis. Additionally, NSAIDs can interfere with the antiplatelet activity of aspirin.

bisphosphonates

The risk of GI ulceration is increased with concurrent use of NSAIDs and bisphosphonates.

cyclosporine

NSAIDs may affect renal prostaglandins and increase the nephrotoxic effect of cyclosporine.

diuretics

Due to the inhibition of renal prostaglandin synthesis, NSAIDs may reduce the natriuretic effect of furosemide and thiazide diuretics.

fluconazole

Concomitant use of fluconazole and celecoxib has been noted to increase the celecoxib plasma concentration as much as 2-fold.

voriconazole

Concomitant use of voriconazole increases the systemic exposure to diclofenac. When concomitant voriconazole use is necessary, the total daily dose of diclofenac should not exceed the lowest recommended dose of diclofenac sodium/misoprostol (Arthrotec) 50 twice daily.

lithium

NSAIDs may produce an elevation of plasma lithium levels and a reduction in renal lithium clearance. These effects have been attributed to inhibition of renal prostaglandin synthesis by the NSAID. Patients should be monitored for signs of lithium toxicity when lithium and NSAIDs are given concurrently.

selective serotonin receptor inhibitors (SSRIs)

There is an increased risk of GI bleeding when SSRIs and NSAIDs are given concurrently.

warfarin

The effects of warfarin and NSAIDs on GI bleeding are synergistic, thereby increasing the risk of serious GI bleeding when used together.

methotrexate

Concomitant use of NSAIDs with methotrexate may increase the toxicity of methotrexate. Concomitant use of esomeprazole, a proton pump inhibitor, with methotrexate may increase and prolong serum levels of methotrexate and its metabolite, possibly leading to methotrexate toxicities. In high-dose methotrexate administration, a temporary withdrawal of the esomeprazole/naproxen (Vimovo) may be considered in some patients.

St John's wort

Avoid concomitant use of esomeprazole/naproxen (Vimovo) with St John's wort due to the potential reduction in esomeprazole levels.

rifampin

Avoid concomitant use of esomeprazole/naproxen (Vimovo) with rifampin due to the potential reduction in esomeprazole levels.

tacrolimus

Concomitant administration of esomeprazole/naproxen and tacrolimus may increase the serum levels of tacrolimus.

gastric pH

Esomeprazole inhibits gastric acid secretion and may interfere with the absorption of drugs where gastric pH is an important determinant of bioavailability (e.g., ketoconazole, iron salts, and digoxin).

sodium polystyrene

Intestinal necrosis, possibly fatal has been reported with the concomitant use of sorbitol with sodium polystyrene sulfonate (Kayexalate®). Due to the presence of sorbitol in meloxicam tablets (Mobic) oral suspension, use with Kayexalate is not recommended.

Ketorolac nasal spray is contraindicated in combination with probenecid and pentoxifylline.

Nasal NSAIDs

ACE inhibitors and angiotensin receptor antagonists

NSAIDs may diminish the antihypertensive effect of these agents.

antiepileptic drugs (phenytoin, carbamazepine)

Cases of seizures have been reported with patients taking concomitant ketorolac.

aspirin

Concomitant use of aspirin and NSAIDs is not generally recommended because of the potential for GI ulceration. NSAID therapy is not a substitute for aspirin for CV prophylaxis.

diuretics

Due to the inhibition of renal prostaglandin synthesis, NSAIDs may reduce the natriuretic effect of furosemide and thiazide diuretics.

lithium

NSAIDs may produce an elevation of plasma lithium levels and a reduction in renal lithium clearance. These effects have been attributed to inhibition of renal prostaglandin synthesis by the NSAID. Patients should be monitored for signs of lithium toxicity when lithium and NSAIDs are given concurrently.

methotrexate

Concomitant use of NSAIDs with methotrexate may increase the toxicity of methotrexate.

psychoactive drugs

Hallucinations have been reported in patients taking psychoactive drugs and ketorolac.

selective serotonin receptor inhibitors (SSRIs)

There is an increased risk of GI bleeding when SSRIs and NSAIDs are given concurrently.

warfarin

The effects of warfarin and NSAIDs on GI bleeding are synergistic, thereby increasing the risk of serious GI bleeding when used together.

Topical NSAIDs

All topical diclofenac formulations, including products co-packaged as a kit, have a similar profile to other NSAIDs and may interact with ACE inhibitors, aspirin, diuretics, lithium, methotrexate, and warfarin.

ADVERSE EFFECTS ^{255,256,257,258,259,260,261,262,263,264,265,266,267,268,269,270,271,272,273,274,275,276,277,278,279,280,281,282,283,284,285,286,287,288,289,290,291,292,293,294,295,296,297,298,299,300}

Oral NSAIDs

Drug	Abdominal Pain (%)	Diarrhea (%)	Dyspepsia (%)	Nausea (%)	Headache (%)	Rash (%)	Edema (%)
Single Ingredient Agents							
celecoxib (Celebrex)	4.1 (2.8)	5.6 (3.8)	8.8 (6.2)	3.5 (4.2)	15.8 (20.2)	2.2 (2.1)	2.1 (1.1)
diclofenac potassium	1–10	1–10	1–10	1–10	1–10	1–10	1–10
diclofenac potassium (Zipsor)	7 (3.4)	2.3 (2.8)	1.2 (2.4)	16.5 (20.2)	12.5 (17.1)	nr	nr
diclofenac sodium (Voltaren, Voltaren XR)	1–10	1–10	1–10	1–10	1–10	1–10	1–10
diclofenac submicronized (Zorvolex)	1–10	1–10	2 (1)	27 (37)	13 (15)	1–10	33 (32)
diflunisal	3–9	3–9	3–9	3–9	3–9	3–9	< 1
etodolac (Lodine)	1–10	1–10	1–10	1–10	1–10	1–10	1–10
fenoprofen (Nalfon)	2 (1.1)	1.8 (4.1)	10.3 (2.3)	7.7 (7.1)	8.7 (7.5)	3.7 (0.4)	5 (0.4)
flurbiprofen	≥ 1	≥ 1	≥ 1	≥ 1	≥ 1	≥ 1	≥ 1
ibuprofen (Motrin)	3–9	3–9	reported	3–9	1–3	3–9	1–3
indomethacin (Indocin)	> 1	> 1	> 1	> 1	11.7	< 1	< 1
indomethacin (Tivorbex)	1–2 (1)	2–3 (1)	1–3 (1)	33–34 (36)	11–16 (11)	1–2 (0)	24–26 (32)
ketoprofen (Orudis)	3–9	3–9	11	3–9	3–9	> 1	2
ketorolac	> 10	1–10	> 10	> 10	> 10	1–10	1–10
meclofenamate	nr	10–33	nr	11	3–9	3–9	1–3
mefenamic acid	1–10	1–10	1–10	1–10	1–10	1–10	1–10

Adverse Effects (continued)

Drug	Abdominal Pain (%)	Diarrhea (%)	Dyspepsia (%)	Nausea (%)	Headache (%)	Rash (%)	Edema (%)
Single Ingredient Agents (continued)							
meloxicam (Mobic, Qmiiz ODT)	1.9-4.7 (0.6–2.5)	1.9-7.8 (3.8)	3.8-9.5 (4.5)	2.4–7.2 (2.6–3.2)	2.4–8.3 (6.4–10.2)	0.3–3 (1.7–2.5)	0.6–4.5 (2.5)
meloxicam submicronized (Vivlodex)	≥ 2	≥ 2 (1)	nr	≥ 2	nr	nr	nr
nabumetone (Relafen DS)	12	14	13	3–9	3–9	3–9	3–9
naproxen (Anaprox / DS, Naprelan, EC/ Naprosyn)	3–9	< 3	< 3	< 3	3–9	1–10	3–9
oxaprozin (Daypro)	>1	>1	>1	>1	> 1	>1	>1
piroxicam (Feldene)	1–10	1–10	1–10	1–10	1–10	1–10	1–10
sulindac	10	3–9	3–9	3–9	3–9	3–9	1–3
tolmetin	3–9	3–9	3–9	11	3–9	reported	3–9
Combination Agents							
diclofenac sodium/ misoprostol (Arthrotec)	21	19	14	11	reported	reported	nr
esomeprazole/ naproxen (Vimovo)	4 (3)	6 (4)	8 (12)	4 (4)	3 (5)	reported	3 (1)
ibuprofen/famotidine (Duexis)	2	4–5	5–8	5–6	3	nr	2

Adverse effects data are obtained from prescribing information and therefore, should not be considered comparative or all inclusive. Incidences for the placebo group indicated in parentheses. nr = not reported

*Adverse effects reported are based on the NSAID component.

Nasal NSAIDs

The most commonly reported adverse effects of ketorolac nasal spray (Sprix) (incidence > 2%) and occurring at a rate at least twice that of placebo are: nasal discomfort, rhinalgia, increased lacrimation, throat irritation, oliguria, rash, bradycardia, decreased urine output, increased ALT and/or AST, hypertension, and rhinitis.

Topical NSAIDs

Drug	Pruritus (%)	Dermatitis (%)	Burning (%)	Nausea (%)	Dysgeusia (%)	Headache (%)
diclofenac epolamine (Flector, Licart)	5	2	< 1	3	2	1
diclofenac sodium (Lexixryl, Pennsaid 2% pump, Vopac MDS, Xrylix)	4 (2)	9 (2)	nr	4 (1)	nr	reported
diclofenac sodium (Voltaren Gel, Diclozor, DS Prep Pak)	< 1	4	nr	nr	nr	nr

Adverse effects data are obtained from package inserts and are not meant to be comparative or all inclusive. Incidences for the placebo group are indicated in parentheses. nr = not reported.

Safety information for Pennsaid 2% pump was based on the safety studies done for the Pennsaid 1.5% solution.

SPECIAL POPULATIONS [301,302,303,304,305,306,307,308,309,310,311,312,313,314,315,316,317,318,319,320,321,322,323,324,325,326,327,328,329,330,331,332,333,334,335,336,337,338,339,340,341,342,343,344,345](#)

Oral NSAIDs

Pediatrics

NSAIDs should be used with caution in patients with systemic onset juvenile idiopathic arthritis (JIA), also known as juvenile rheumatoid arthritis (JRA), due to the risk of disseminated intravascular coagulation. Patients with systemic onset JIA should be monitored for the development of abnormal coagulation tests.

Celecoxib (Celebrex) is indicated for the relief of the signs and symptoms of JIA in patients 2 years and older. The use of celecoxib in patients 2 years to 17 years of age with pauciarticular, polyarticular course JIA or in patients with systemic onset JIA was studied in a 12-week, double-blind, active-controlled, pharmacokinetic, safety, and efficacy study, with a 12-week open-label extension. Safety and efficacy have not been studied beyond 6 months in children. Celecoxib has not been studied in patients < 2 years of age, in patients with body weight < 10 kg (22 lbs), and in patients with active systemic features. In some patients with systemic onset JIA, both celecoxib and naproxen were associated with mild prolongation of activated partial thromboplastin time (aPTT) but not prothrombin time (PT). NSAIDs, including celecoxib, should be used only with caution in patients with systemic onset JIA due to the risk of disseminated intravascular coagulation. Patients with systemic onset JIA should be monitored for the development of abnormal coagulation tests. Data on the menthol/capsaicin and lidocaine/menthol patch are limited; a physician should be consulted prior to use in patients < 12 years old.

Safety and efficacy of etodolac extended-release tablets for the relief of signs and symptoms of JIA in patients 6 to 16 years of age are supported by extrapolating data from adequate and well-controlled studies in adult rheumatoid arthritis patients and also by safety, efficacy, and pharmacokinetic data

from an open-label clinical trial in JIA patients 6 to 16 years of age. However, safety and effectiveness of etodolac immediate-release in pediatric patients < 18 years have not been established.

Ibuprofen (Motrin) has been tested in children ≥ 6 months of age. It has not been shown to cause different adverse effects or problems than it does in adults. Over-the-counter strengths of ibuprofen are available for use in children.

Ketorolac tromethamine nasal spray (Sprix) is not for use in patients < 2 years of age. The safety and effectiveness of ketorolac in patients ≤ 17 years of age have not been established.

Meloxicam (Mobic, Qmiiz ODT) is indicated for JIA in patients > 2 years and whose weight ≥ 60 kg. Safety and efficacy of indomethacin (Indocin) and mefenamic acid in children ≤ 14 years of age have not been established.

Safety and efficacy of diflunisal in children < 12 years of age have not been established.

Safety and efficacy of oxaprozin for the relief of signs and symptoms of JIA in patients 6 to 16 years of age are supported by extrapolating from adequate and well-controlled studies in adult rheumatoid arthritis patients. Safety and efficacy of oxaprozin in pediatric patients < 6 years of age have not been established.

Safety and efficacy of tolmetin sodium have not been established in children < 2 years of age.

Safety and efficacy of naproxen (EC-Naprosyn, Naprosyn, Anaprox, and Anaprox DS) in patients < 2 years of age have not been established. The use of naproxen suspension (Naprosyn Suspension) is recommended for JIA in children ≥ 2 years of age because it allows for more flexible dose titration based on the child's weight. Single doses of 2.5 to 5 mg/kg, with total daily dose not exceeding 15 mg/kg/day, are well tolerated in pediatric patients > 2 years of age. Pharmacokinetic studies of naproxen were not performed in pediatric patients < 5 years of age. Per the pharmacokinetic section of the manufacturer prescribing information, naproxen delayed-release (EC-Naprosyn) has not been studied in patients < 18 years of age. Naproxen controlled-release (Naprelan) has not been studied in patients < 18 years of age.

Vimovo is indicated for juvenile idiopathic arthritis (JIA) in adolescent patients ≥ 12 years of age who weigh ≥ 38 kg, who require symptomatic relief of arthritis, and are at risk of developing NSAID-related gastric ulcers.

Safety and efficacy have not been established in patients < 18 years of age for the following agents: diclofenac (Zorvolex), diclofenac potassium (Zipsor), diclofenac sodium (Voltaren, Voltaren XR), diclofenac sodium/misoprostol (Arthrotec), flurbiprofen, prescription strength ibuprofen (Motrin), ibuprofen/famotidine (Duexis), indomethacin (Tivorbex), ketoprofen, meclofenamate, meloxicam submicronized (Vivlodex), nabumetone (Relafen DS), piroxicam (Feldene), and sulindac.

NSAIDs combined with proton pump inhibitors have no data supporting pediatric use; however, esomeprazole (Nexium®) is indicated for use in patients > 1 year of age, and naproxen has been proven safe and effective in patients ≥ 2 years.

Pregnancy

Use of NSAIDs during the third trimester of pregnancy increases the risk of premature closure of the fetal ductus arteriosus. Avoid use of NSAIDs, including ibuprofen/famotidine and naproxen/esomeprazole, in pregnant women starting at 30 weeks gestation. Previously, oral NSAIDs

were considered Pregnancy Category C prior to 30 weeks gestation and Category D after 30 weeks. Diclofenac sodium/misoprostol (Arthrotec) is Category X and has a boxed warning because misoprostol may cause abortions in pregnant women. Also, meclufenamate may be associated with miscarriage and minor skeletal malformations.

NSAIDs, including meloxicam (Mobic, Qmiiz ODT), may be associated with a reversible delay in ovulation. Therefore, in women who have difficulties conceiving, or who are undergoing investigation of infertility, use of meloxicam is not recommended.

Renal Insufficiency

Please see Warnings section of this review.

The maximum meloxicam submicronized (Vivlodex) dose for patients on hemodialysis is 5 mg per day.

Hepatic Insufficiency

The daily dose of celecoxib in patients with moderate hepatic impairment should be decreased by 50%; celecoxib use in patients with severe hepatic impairment is not recommended.

Patients who are known or suspected to be poor CYP2C9 metabolizers based on previous history and/or experience with other CYP2C9 substrates (such as warfarin, phenytoin) should be administered celecoxib with caution. Consider starting treatment at half the lowest recommended dose in poor metabolizers. Consider using alternative management in JIA patients who are poor metabolizers.

Elderly

NSAIDs should be used with caution in elderly patients (65 years of age and older) since advancing age appears to increase the possibility of adverse effects. Elderly patients may be less tolerant of GI ulceration or bleeding than other individuals, and fatal GI reactions have been reported in this population. Indomethacin may cause confusion or, on rare occasions, psychosis.

NSAIDs and famotidine are known to be substantially excreted by the kidney, and the risk of toxic effects to NSAIDs may be greater in patients with renal function impairment. Because elderly patients are more likely to have decreased renal function, take care in dose selection, and it may be useful to monitor renal function.

For patients > 65 years, the dosage of ketorolac nasal spray (Sprix) is reduced to 15.75 mg (1 spray in only 1 nostril) every 6 to 8 hours for a daily maximum dose of 63 mg.

Nasal NSAIDs

Pediatrics

Safety and efficacy of oral ketorolac in patients < 17 years of age has not been established. Ketorolac nasal spray (Sprix) is not for use in pediatric patients less than 2 years of age.

Pregnancy

Ketorolac nasal spray (Sprix) is Pregnancy Category C prior to 30 weeks gestation; thereafter, it is Category D in late pregnancy since it can cause premature closure of the ductus arteriosus and should therefore be avoided.

Renal Impairment

Long-term administration of NSAIDs has resulted in renal papillary necrosis and other renal injury.

Hepatic Impairment

Elevations of 1 or more liver tests may occur in up to 15% of patients taking NSAIDs including diclofenac formulations. Notable elevations of alanine aminotransferase (ALT) or aspartate aminotransferase (AST) (approximately 3 x ULN) have been reported in about 1% of patients in clinical trials with NSAIDs. In addition, rare cases of severe hepatic reactions, including jaundice and fatal fulminant hepatitis, liver necrosis, and hepatic failure, some with fatal outcomes, have been reported.

Patients with severe hepatic disease are at greater risk of developing toxic blood concentrations of lidocaine because of their inability to metabolize lidocaine normally.

Geriatrics

Exercise caution when using ketorolac nasal spray (Sprix) in elderly patients.

Topical NSAIDs

Pediatrics

Some diclofenac epolamine topical systems (e.g., Flector and generics) are FDA-approved in patients as young as 6 years old; others (e.g., Licart) are labeled for adult use only. Safety and effectiveness in pediatric patients for the other topical NSAID products in this review have not been established.

Pregnancy

Diclofenac containing formulations are Pregnancy Category C prior to 30 weeks gestation; avoid use (Category D) starting at 30 weeks gestation due to the risk of premature closure of the fetal ductus arteriosus.

Renal Impairment

Long-term administration of NSAIDs has resulted in renal papillary necrosis and other renal injury. Diclofenac formulations are not recommended for use in patients with advanced renal disease.

Hepatic Impairment

Elevations of 1 or more liver tests may occur in up to 15% of patients taking NSAIDs including diclofenac formulations. Notable elevations of alanine aminotransferase (ALT) or aspartate aminotransferase (AST) (approximately 3 x ULN) have been reported in about 1% of patients in clinical trials with NSAIDs. In addition, rare cases of severe hepatic reactions, including jaundice and fatal fulminant hepatitis, liver necrosis, and hepatic failure, some with fatal outcomes, have been reported.

Patients with severe hepatic disease are at greater risk of developing toxic blood concentrations of lidocaine because of their inability to metabolize lidocaine normally.

Geriatrics

Diclofenac, as with any NSAID, is known to be substantially excreted by the kidney, and the risk of toxic reactions to diclofenac formulations may be greater in patients with impaired renal function. Because

elderly patients are more likely to have decreased renal function, care should be taken when using diclofenac formulations in the elderly, and it may be useful to monitor renal function.

DOSAGES ^{346,347,348,349,350,351,352,353,354,355,356,357,358,359,360,361,362,363,364,365,366,367,368,369,370,371,372,373,374,375,376,377,378,379,380,381,382,383,384,385,386,387,388,389,390,391,392,393,394,395,396,397,398,399,400,401,402,403,404,405,406,407,408,409,410,411,412,413,414,415,416,417,418,419,420}

Oral NSAIDs

Drug	Recommended Dosages	Maximum Daily Dose (MDD)	Availability
Single Ingredient Agents			
celecoxib (Celebrex)	OA: 200 mg daily or 100 mg twice daily RA: 100 to 200 mg twice daily JIA: 50 mg twice daily (patients ≥ 10 kg to ≤ 25 kg); 100 mg twice daily (patients > 25 kg) AS: 200 mg daily or 100 mg twice daily; may increase to 400 mg/day after 6 weeks Dysmenorrhea or acute pain: 400 mg once with an additional 200 mg on day 1, then 200 mg twice daily, as needed	OA: 200 mg RA: 400 mg JIA: 100 mg in patients ≥ 10 kg; 200 mg in patients > 25 kg AS: 400 mg Dysmenorrhea/Acute Pain: 400 mg	Capsules: 50 mg, 100 mg, 200 mg, 400 mg
diclofenac potassium	OA: 50 mg 2 to 3 times daily RA: 50 mg 3 to 4 times daily Pain & Primary Dysmenorrhea (PD): 50 mg 3 times daily OR initial dose of 100 mg, followed by 50 mg 3 times daily	OA: 200 mg RA: 225 mg Pain & PD: 200 mg on initial day, followed by 150 mg	Tablets: 50 mg
diclofenac potassium (Zipsor)	Pain: 25 mg 4 times daily	Pain (mild to moderate acute): 100 mg	Capsules: 25 mg
diclofenac sodium (Voltaren)	OA: 50 mg 2 to 3 times daily, or 75 mg twice daily RA: 50 mg 3 to 4 times daily, or 75 mg twice daily AS: 25 mg 4 times daily. May repeat 25 mg dose at bedtime, if necessary PD: Initial dose of 50 mg to 100 mg daily; may titrate up to a maximum of 200 mg/day	OA: 150 mg RA: 200 mg AS: 125 mg PD: 200 mg	Tablets, delayed-release: 25 mg, 50 mg, 75 mg
diclofenac sodium XR/DR (Voltaren XR)	OA: 100 mg once daily RA: 100 mg once daily; may be increased to 100 mg twice daily	OA: 100 mg RA: 200 mg	Tablets, extended-release: 100 mg

MDD = Maximum Daily Dose.

Dosages: Oral NSAIDs (continued)

Drug	Recommended Dosages	Maximum Daily Dose (MDD)	Availability
Single Ingredient Agents (continued)			
diclofenac submicronized (Zorvolex)	Pain: 18 to 35 mg 3 times daily OA: 35 mg 3 times daily (can be taken with or without food, but food can decrease its effectiveness)	Pain (mild to moderate acute) & OA: 105 mg	Capsules: 18 mg, 35 mg Take on an empty stomach
diflunisal	OA & RA: 250 mg once daily to 500 mg twice daily Pain: An initial dose of 1,000 mg followed by 500 mg every 12 hours; Following the initial dose, some patients may require 500 mg every 8 hours; A lower dosage may be appropriate depending on factors such as pain severity, patient response, weight, or advanced age (e.g., 500 mg initially, followed by 250 mg every 8 to 12 hours)	OA & RA: 1,500 mg Pain maintenance: 1,500 mg	Tablets: 500 mg Tablets should be swallowed whole, not crushed or chewed
etodolac (Lodine)	OA & RA: 300 mg 2 to 3 times daily, 400 mg twice daily, or 500 mg twice daily for immediate-release; 400 to 1,000 mg daily for extended-release Pain: 200 to 400 mg every 6 to 8 hours, up to 1000 mg/day for immediate-release JIA (extended-release only): Daily, based on body weight: <ul style="list-style-type: none"> ▪ 20 to 30 kg: 400 mg ▪ 31 to 45 kg: 600 mg ▪ 46 to 60 kg: 800 mg ▪ > 60 kg: 1,000 mg 	OA & RA: 1,000 mg Pain: 1,000 mg JIA: 1,000 mg	Capsules: 200 mg, 300 mg Tablets: 400 mg, 500 mg Tablets, extended-release: 400 mg, 500 mg, 600 mg
fenoprofen (Nalfon)	OA & RA: 300 to 600 mg 3 to 4 times daily Pain: 200 mg every 4 to 6 hours, as needed	OA & RA: 3,200 mg Pain: 1,200 mg	Capsules: 200 mg, 400 mg Tablets: 600 mg
flurbiprofen	OA & RA: 200 to 300 mg per day administered in divided doses 2 to 4 times a day; the largest recommended single dose in a multiple-dose daily regimen is 100 mg	OA & RA: 300 mg	Tablets: 50 mg, 100 mg
ibuprofen (Motrin)	OA & RA: 300 mg 4 times daily; 400, 600, or 800 mg 3 or 4 times daily Pain: 400 mg every 4 to 6 hours, as needed PD: 400 mg every 4 hours, as needed	OA & RA: 3,200 mg Pain: 2,400 mg PD: 2,400 mg	Tablets: 400 mg, 600 mg, 800 mg Suspension: 100 mg/5 mL

MDD = Maximum Daily Dose.

Dosages: Oral NSAIDs (continued)

Drug	Recommended Dosages	MDD	Availability
Single Ingredient Agents (continued)			
indomethacin (Indocin)	<p>OA, RA & AS: 25 mg 2 to 3 times daily for immediate release; 75 mg once daily for extended release (a large portion of total daily dose may be administered in the evening rectally with suppository retained for at least 60 minutes)</p> <p>Acute painful shoulder: 75 to 150 mg daily in 3 or 4 divided doses immediate release; 75 mg twice daily for ER capsules; when 150 mg is prescribed, give as 1 capsule twice daily; Continue therapy until the signs and symptoms of inflammation have been controlled for several days; the usual course of therapy is 7 to 14 days</p> <p>Acute gouty arthritis (immediate-release): 50 mg 3 times daily until pain is tolerable; rapidly reduce dose to complete cessation of the drug</p>	<p>OA, RA & AS: 200 mg</p> <p>ER capsules: 75 mg twice daily</p> <p>Acute painful shoulder: 150 mg</p> <p>Acute gouty arthritis: 150 mg</p>	<p>Capsules, oral: 25 mg, 50 mg</p> <p>Capsules, sustained-release: 75 mg</p> <p>Suppository: 50 mg</p> <p>Suspension: 25 mg/5 mL</p> <p>ER capsules can be administered once a day and can be substituted for indomethacin 25 mg capsules 3 times a day</p>
indomethacin (Tivorbex)	20 mg 3 times daily or 40 mg 2 to 3 times daily	120 mg	Capsules, oral: 20 mg, 40 mg
ketoprofen [†]	<p>OA & RA: 75 mg 3 times daily or 50 mg 4 times daily for immediate-release; 200 mg once daily for extended-release</p> <p>Pain & PD: 25 to 50 mg every 6 to 8 hours, as necessary for immediate-release</p>	<p>OA & RA: 200 to 300 mg</p> <p>Pain & PD: 300 mg</p>	<p>Capsules: 50 mg, 75 mg</p> <p>Capsules, extended-release: 200 mg</p>
ketorolac tromethamine	<p>Short-term (≤ 5 days) for acute pain: Adult patients younger than 65 years of age: Following conversion from injectable formulation, first dose may be 1 or 2 tablets followed by 1 tablet every 4 to 6 hours, not to exceed 40 mg in 24 hours Patients 65 years of age and older, renally impaired, or < 50 kg (110 lbs) of body weight: Following conversion from injectable formulation, 1 tablet every 4 to 6 hours, not to exceed 40 mg in 24 hours</p>	<p>Short-term (≤ 5 days) for acute pain: Adult patients younger than 65 years of age: 40 mg Patients 65 years of age and older, renally impaired, or < 50 kg (110 lbs) of body weight: 40 mg</p>	<p>Tablet: 10 mg</p> <p>Oral formulation should not be given as initial dose</p>
meclofenamate sodium	<p>OA & RA: 200 to 400 mg per day, administered in 3 to 4 equal doses</p> <p>Excessive menstrual blood loss & PD: 100 mg 3 times a day, for up to 6 days, starting at the onset of menstrual flow</p> <p>Pain: 50 mg to 100 mg every 4 to 6 hours</p>	<p>OA & RA: 400 mg</p> <p>Excessive menstrual blood loss & PD: 300 mg</p> <p>Pain: 400 mg</p>	Capsules: 50 mg, 100 mg

MDD = Maximum Daily Dose.

Dosages: Oral NSAIDs (continued)

Drug	Recommended Dosages	MDD	Availability
Single Ingredient Agents (continued)			
mefenamic acid	Pain in patients ≥ 14 years of age : 500 mg as an initial dose followed by 250 mg every 6 hours, as needed, usually not to exceed 1 week PD: 500 mg as an initial dose followed by 250 mg every 6 hours, starting with the onset of bleeding and associated symptoms; treatment should not be necessary for more than 2 to 3 days	Pain: 1,000 mg PD: 1,000 mg	Capsules: 250 mg
meloxicam (Mobic, Qmiiz ODT)	OA & RA: 7.5 mg once daily JIA: 7.5mg once daily for children who weigh ≥ 60 kg	OA & RA: 15 mg once daily JIA: 7.5 mg once daily	Tablets: 7.5 mg, 15 mg Orally Disintegrating Tablets (Qmiiz ODT): 7.5 mg, 15 mg
meloxicam submicronized (Vivlodex)	OA: 5 mg once daily, initially; may increase dose to 10 mg once daily for those requiring additional analgesia	OA: 10 mg once daily	Capsules: 5 mg, 10 mg
nabumetone (Relafen DS)	OA & RA: 1,000 mg once or twice daily; some patients may obtain more relief from 1,500 to 2,000 mg/day which can be given as a single or twice daily dose	OA & RA: 2,000 mg	Tablets: 500 mg, 750 mg, 1,000 mg (Relafen DS)
naproxen (Naprosyn)	OA, RA & AS: 750 to 1,000 mg once daily or 250 to 500 mg twice daily JIA: 5 mg/kg given twice daily; suspension is recommended for patients 2 years of age and older to allow for more accurate titration of the dose Pain, PD & acute tendonitis/bursitis: 1,000 mg to 1,500 mg once daily for a limited period; thereafter, the total daily dose should not exceed 1,000 mg Alternatively, a 500 to 550 mg first dose, followed by 500 to 550 mg every 12 hours or 250 to 275 mg every 6 to 8 hours Acute gout: The starting dose is 1,000 to 1,500 mg once daily, then 1,000 mg once daily Alternatively, 750 to 825 mg to start, followed by 250 to 275 mg every 8 hours until the attack subsides	OA, RA & AS: 550 to 1,500 mg JIA: 10 mg/kg Pain, PD & acute tendonitis/bursitis: The initial total daily dose should not exceed 1,250 to 1,500 mg; thereafter, the total daily dose should not exceed 750 to 1,100 mg Acute gout: The initial total daily dose should not exceed 1,250 to 1,500 mg; thereafter, the total daily dose should not exceed 750 to 1,000 mg	Tablets: 250 mg, 275 mg, 375 mg, 500 mg, 550 mg Tablet, delayed-release: 375 mg, 500 mg Tablets, controlled-release: 375 mg, 500 mg, 750 mg Suspension: 125 mg/5 mL
oxaprozin (Daypro)	OA & RA: 1,200 mg once daily JIA: For patients 22 to 31 kg, give 600 mg; for 32 to 54 kg, give 900 mg; for ≥ 55 kg, give 1,200 mg	OA & RA: 1,800 mg JIA: 26 mg/kg	Tablet: 600 mg
piroxicam (Feldene)	OA & RA: 10 mg twice daily or 20 mg once daily	OA & RA: 20 mg	Capsules: 10 mg, 20 mg

MDD = Maximum Daily Dose.

Dosages: Oral NSAIDs (continued)

Drug	Recommended Dosages	MDD	Availability
Single Ingredient Agents (continued)			
sulindac	OA, RA & AS: 150 mg twice daily with food Acute shoulder pain: 200 mg twice daily with food Acute gouty arthritis: 200 mg twice daily with food	OA & RA: 400 mg Acute shoulder pain: 400 mg Acute gouty arthritis: 400 mg	Tablets: 150 mg, 200 mg
tolmetin	OA & RA: 400 mg 3 times daily JIA: Starting dosage is 20 mg/kg/day in 3 to 4 divided doses; the usual dose is 15 to 30 mg/kg/day once control is achieved	OA & RA: 1,800 mg JIA: 30 mg/kg/day	Tablets: 200 mg, 600 mg Capsules: 400 mg
Combination Agents			
celecoxib/ methyl salicylate/ menthol/ capsaicin (NuDroxiPak)	Celecoxib: dosing based on celecoxib component (described above); the lowest effective dose and treatment duration should be used to achieve the treatment goal for each patient Methyl salicylate/menthol/capsaicin solution: apply to affected area as directed	Celecoxib: 200 mg/day Methyl salicylate/menthol/capsaicin solution: as directed	Kit containing 100 oral capsules of celecoxib 200 mg co-packaged with methyl salicylate 25%/menthol 6%/capsaicin 0.025% solution (NuDroxicin Pain Relief Roll-on)
diclofenac sodium DR/capsaicin (Flexipak, Inavix, Inflammacin, Nudiclo TabPak, Xenaflamm)	Diclofenac sodium: dosing for AS, OA, and RA based on diclofenac sodium component (described above); the lowest effective dose and treatment duration should be used to achieve the treatment goal for each patient Capsaicin: apply thin film to affected area 3 to 4 times daily; wash and dry hands after use	Diclofenac sodium: 200 mg/day Capsaicin: 4 times daily	Flexipak, Inavix, Nudiclo: kit containing sixty 75 mg tablets co-packaged with capsaicin cream 0.025% (60 g tube) Inflammacin, Xenaflamm: kit containing sixty 75 mg tablets co-packaged with capsaicin cream 0.025% (237 mL tube)
diclofenac sodium DR/ capsaicin/ methyl salicylate/ menthol (NuDroxiPak DSDR-50, NuDroxiPak DSDR-75)	Diclofenac sodium: dosing for AS, OA, and RA based on diclofenac sodium component (described above); the lowest effective dose and treatment duration should be used to achieve the treatment goal for each patient Methyl salicylate/menthol/capsaicin solution: apply topically to affected area as directed	Diclofenac sodium: 200 mg/day Methyl salicylate/menthol/capsaicin solution: as directed	Kit containing sixty 50 mg (DSDR-50) or 75 mg (DSDR-75) diclofenac sodium delayed-release tablets co-packaged with methyl salicylate 25%/menthol 6%/capsaicin 0.025% solution (NuDroxicin Pain Relief Roll-on)
diclofenac sodium DR / menthol / camphor (Inflatherm™)	Diclofenac sodium: dosing for AS, OA, and RA based on diclofenac sodium component (described above); the lowest effective dose and treatment duration should be used to achieve the treatment goal for each patient Menthol/camphor gel: apply topically to affected area 3 to 4 times daily	Diclofenac sodium: 200 mg/day Menthol/camphor: 4 times per day	Kit containing sixty 75 mg diclofenac sodium delayed-release tablets co-packaged with menthol 3%/camphor 3% (Sore No More Natural Pain Relieving Gel)

MDD = Maximum Daily Dose.

Dosages: Oral NSAIDs (continued)

Drug	Recommended Dosages	MDD	Availability
Combination Agents (continued)			
diclofenac sodium/ misoprostol (Arthrotec)	OA: 50 mg/200 mcg 2 to 3 times daily; or 75 mg/200 mcg twice daily RA: 50 mg/200 mcg 2 to 4 times daily; or 75 mg/200 mcg twice daily	OA: 150 mg of diclofenac RA: 225 mg of diclofenac Note: Limit misoprostol to 200 mcg at any 1 time.	Tablets: 50 mg/200 mcg, 75 mg/200 mcg
etodolac/ capsaicin/ methyl salicylate/ menthol (NuDroxiPak E-400)	Etodolac: dosing for OA, RA, and acute pain is based on etodolac component (described above); the lowest effective dose and treatment duration should be used to achieve the treatment goal for each patient Methyl salicylate/menthol/capsaicin solution: apply topically to affected area as directed	Etodolac: 1,000 mg/day Methyl salicylate/ menthol/capsaicin solution: as directed	Kit containing 100 oral tablets of etodolac 400 mg co-packaged with methyl salicylate 25%/menthol 6%/ capsaicin 0.025% solution (NuDroxacin Pain Relief Roll- on)
esomeprazole/ naproxen (Vimovo)	OA, RA, AS: 375 mg or 500 mg of naproxen with 20 mg of esomeprazole twice daily JIA: Adolescents ≥ 12 years weighing 38 kg to < 50 kg 375 mg/20 mg tab twice daily	OA, RA, AS: 1,000 mg of naproxen If total dose of < 40mg esomeprazole is more appropriate, consider an alternate treatment Controlled studies do not extend beyond 6 months	Tablets: 375 mg/20 mg, 500 mg/20 mg Do not split, crush or dissolve; take at least 30 minutes before meals
ibuprofen/ famotidine (Duexis)	OA, RA: 1 tablet 3 times daily (not considered interchangeable with single-component products)	3 tablets	Tablets: 800 mg/26.6 mg Do not chew, divide, or crush tablets
ibuprofen/ capsaicin/ menthol/ methyl salicylate (Comfort Pac with Ibuprofen; NuDroxiPak I-800)	Ibuprofen: dosing based on ibuprofen component (as described above); the lowest effective dose and treatment duration should be used to achieve the treatment goal for each patient Methyl salicylate/menthol/capsaicin: as directed	Ibuprofen: see above for individually packaged naproxen Methyl salicylate/ menthol/capsaicin: As directed	Tablets: 800 mg Co-packaged with: Duraflex Comfort Gel: 60 mL (capsaicin/ menthol/methyl salicylate) – For Comfort Pac OR NuDroxacin Pain Relief Roll- on: capsaicin 0.025%/ menthol 6%/ methyl salicylate 25% solution – For NuDroxiPak

MDD = Maximum Daily Dose.

Dosages: Oral NSAIDs (continued)

Drug	Recommended Dosages	MDD	Availability
Combination Agents (continued)			
meloxicam/ capsaicin/ menthol/ methyl salicylate (Comfort Pac with Meloxicam)	Meloxicam: dosing based on meloxicam component (as described above); the lowest effective dose and treatment duration should be used to achieve the treatment goal for each patient Methyl salicylate/menthol/capsaicin: As directed	Meloxicam: see above for individually packaged naproxen Methyl salicylate/menthol/capsaicin: As directed	Tablets: 15 mg Co-packaged with: Duraflex Comfort Gel: 60 mL (capsaicin/ menthol/ methyl salicylate) – For Comfort Pac OR NuDroxicin Pain Relief Roll-on: capsaicin 0.025%/ menthol 6%/ methyl salicylate 25% solution – For NuDroxiPak
nabumetone/ capsaicin/ menthol/ methyl salicylate (NuDroxiPak N-500)	Nabumetone: dosing based on nabumetone component (as described above); the lowest effective dose and treatment duration should be used to achieve the treatment goal for each patient Methyl salicylate/menthol/capsaicin: As directed	Naproxen: see above for individually packaged nabumetone Methyl salicylate/menthol/capsaicin: As directed	Tablets: 500 mg Co-packaged with: NuDroxicin Pain Relief Roll-on: capsaicin 0.025%/ menthol 6%/ methyl salicylate 25% solution
naproxen/ capsaicin/ menthol/ methyl salicylate (Comfort Pac with Naproxen)	Naproxen: dosing based on naproxen component (as described above); the lowest effective dose and treatment duration should be used to achieve the treatment goal for each patient Methyl salicylate/menthol/capsaicin: As directed	Naproxen: see above for individually packaged naproxen Methyl salicylate/menthol/capsaicin: As directed	Tablets: 500 mg Co-packaged with: Duraflex Comfort Gel: 60 mL (capsaicin/ menthol/ methyl salicylate) – For Comfort Pac

MDD = Maximum Daily Dose.

† A micronized ketoprofen 10% cream (Frotek) by Nubrotori is also available in 30 mL bottles.⁴²¹

Different formulations of oral diclofenac (e.g., diclofenac sodium enteric-coated tablets, diclofenac sodium extended-release tablets, or diclofenac potassium immediate-release tablets) may not be bioequivalent even if the milligram strength is the same. Diclofenac capsules are not interchangeable with other oral diclofenac formulations.

Meloxicam submicronized (Vivlodex) capsules are not interchangeable with other formulations of oral meloxicam, even if the mg strength is the same.

Nasal NSAIDs

Drug	Recommended Dosages	Maximum Daily Dose (MDD)	Availability
ketorolac nasal spray (Sprix)	<p>Short-term (≤ 5 days) for acute pain: Adult patients younger than 65 years of age: 31.5 mg (1 spray [15.75mg] in each nostril) every 6 to 8 hours</p> <p>Patients 65 years of age and older, renally impaired, or < 50 kg (110 lbs) of body weight: 15.75 mg (1 spray [15.75 mg] in only 1 nostril) every 6 to 8 hours; maximum daily dose is 63 mg (4 doses)</p>	<p>Short-term (≤ 5 days) for acute pain: Adult patients younger than 65 years of age: 126 mg (4 doses per 24 hours)</p> <p>Patients 65 years of age and older, renally impaired, or < 50 kg (110 lbs) of body weight: 63 mg (4 doses per 24 hours)</p>	Nasal spray: 8 sprays per bottle. Bottle must be discarded 24 hours after priming

Topical NSAIDs

Drug	Adult Dosage	Special Handling and Disposal	Availability
diclofenac epolamine (Flector, Licart)	Apply 1 topical system to the most painful area twice daily	<p>Hand washing is recommended after applying, handling, or removing this patch</p> <p>Do not wear patch during bathing/showering</p> <p>Place only on intact skin</p> <p>If patch begins to “peel back” it may be taped down or use a non-occlusive mesh netting sleeve</p> <p>Storage envelope should remain sealed at all times when not in use</p>	1.3% topical system
diclofenac sodium (Lexixryl, Xrylix)	40 drops per knee (applying 10 drops at a time), 4 times daily and spread evenly around knee	Wash and dry hands after use	1.5% topical solution Lexixryl/Xrylix: kit containing 1.5% topical solution co-packaged with 30 Xrylix sheets (kinesiology tape/cross tape)
diclofenac sodium (Vopac MDS kit)	6 sprays per knee (equivalent to 40 drops), every 6 to 8 hours	Clean area with alcohol prep pads (in kit), allow area to dry prior to application; wash and dry hands after use	1.5% topical solution (150 mL) co-packaged with 100 alcohol prep pads, 1 pair sterile gloves, and a metered dose spray bottle
diclofenac sodium (Pennsaid 2% pump)	2 pump actuations per knee 2 times per day and spread evenly around knee	Wash and dry hands after use	2% topical solution

Dosages: Topical NSAIDs (continued)

Drug	Adult Dosage	Special Handling and Disposal	Availability
diclofenac sodium (Voltaren Gel, Diclozor, DS Prep Pak)	<p>Lower extremities: Apply 4 g to the affected area 4 times daily</p> <p>Upper extremities: Apply 2 g to the affected area 4 times daily</p>	<p>Do not apply more than 16 g daily to any 1 of the affected joints of the lower extremities</p> <p>Do not apply more than 8 g daily to any 1 of the affected joints of the upper extremities</p> <p>Total dose should not exceed 32 g per day over all affected joints</p> <p>Patient should wash hands after use, unless the hands are the treated joint; in which case wait at least 1 hour after the application before washing</p> <p>Showering/bathing should be avoided for at least 1 hour after the application</p> <p>The gel should be measured using the dosing card supplied and can be used to apply the gel; card should be rinsed and dried after use</p>	<p>1% gel (Voltaren, generic)</p> <p>Diclozor: kit containing 1% gel co-packaged with a 10-count of occlusive dressings</p> <p>DS Prep Pak: kit containing 1% gel co-packaged with 20 gloves and 20 antiseptic wipes</p>

Dosages: Topical NSAIDs (continued)

Drug	Adult Dosage	Special Handling and Disposal	Availability
<p>diclofenac sodium/capsaicin (CapsFenac Pak, Capsinac, DermacinRx Lexitral PharmaPak, Diclofex DC, Diclopak, Diclosiacin, Diclotral Pak, Nudiclo Solupak, Sure Result DSS Premium Pak, Xelitral)</p>	<p>40 drops diclofenac sodium per knee (applying 10 drops at a time), 4 times daily and spread evenly around knee; Co-packaged capsaicin cream may be applied up to 4 times daily</p>	<p>Wash and dry hands after use</p>	<p>CapsFenac Pak, Diclopak, Nudiclo Solupak: package contains (1) 1.5% diclofenac sodium topical solution and (2) capsaicin 0.025% cream Capsinac: package contains (1) 1.5% diclofenac sodium topical solution and (2) capsaicin 0.025% cream DermacinRx Lexitral PharmaPak: package contains (1) 1.5% diclofenac sodium topical solution and (2) DermacinRx Penetral™ cream, containing a time released 0.025% capsaicin with acai berry oil and omega 3, 6, and 9 oils Diclofex DC: package contains (1) 1.5% diclofenac sodium topical solution and (2) Bio-Sorb™ Cream, containing 0.025% capsaicin Diclosiacin: package contains (1) 1.5% diclofenac sodium topical solution and (2) capsaicin 0.025% cream Diclotral Pak: package contains (1) 1.5% diclofenac sodium topical solution and (2) Dermafrice™ Cream, containing 0.025% capsaicin Sure Result DSS Premium Pak: package contains (1) 1.5% diclofenac sodium topical solution and (2) Sure Result SR Relief Cream, containing 0.025% capsaicin Xelitral: package contains (1) 1.5% diclofenac sodium topical solution and (2) Penetral™ Cream, containing 0.025% capsaicin</p>
<p>diclofenac sodium/camphor/lidocaine/methyl salicylate (Diclovix)</p>	<p>40 drops diclofenac sodium per knee (applying 10 drops at a time), 4 times daily and spread evenly around knee; Co-packaged patch should be applied to affected area 1 or 2 times daily and leave in place for 8-12 hours</p>	<p>Wash and dry hands after use Wash and dry hands after applying or removing patch</p>	<p>Diclovix kit: package contains (1) 1.5% diclofenac sodium topical solution and (2) Viva Patch, containing camphor 2%, lidocaine 2.5%, and methyl salicylate 4%</p>

Dosages: Topical NSAIDs (continued)

Drug	Adult Dosage	Special Handling and Disposal	Availability
diclofenac sodium/ camphor/ menthol (Diclotrex)	40 drops diclofenac sodium per knee (applying 10 drops at a time), 4 times daily and spread evenly around knee; Co-packaged gel is applied as a liberal amount to the affected area using the roll-on applicator not more than 3 to 4 times per day	Wash and dry hands after use	Package contains (1) 1.5% diclofenac sodium topical solution and (2) Camphotrex™ gel roll-on applicator containing 4% camphor and 10% menthol
diclofenac sodium/ camphor/ menthol/ methyl salicylate (Inflamma-K Kit)	40 drops diclofenac sodium per knee (applying 10 drops at a time), 4 times daily and spread evenly around knee; Co-packaged patch should be applied over affected area following diclofenac application and may be used up to 4 times daily; remove patch after 8 hours	Wash and dry hands after use	Package contains (1) 1.5% diclofenac sodium topical solution and (2) Salonpas patch, containing camphor 3.1%, menthol 6%, and methyl salicylate 10%
diclofenac sodium/ lidocaine/tape (Trixylytral)	40 drops diclofenac sodium per knee (applying 10 drops at a time), 4 times daily and spread evenly around knee; Apply thin lidocaine film to affected area 2 to 3 times per day	Wash and dry hands after use Avoid skin-to-skin contact, covering with clothing, or application of other topical substances until area is completely dry	Package contains (1) 1.5 % diclofenac sodium topical solution and (2) 3.88% lidocaine cream (Lidotral) and (3) kinesiology/cross tape (Xrylix)
diclofenac sodium/ menthol (Dithol Kit, Dimentho)	40 drops diclofenac sodium per knee (applying 10 drops at a time), 4 times daily and spread evenly around knee; Apply menthol to affected area not more than 4 times daily in one direction (not circular)	Wash and dry hands after use Avoid skin-to-skin contact, covering with clothing, or application of other topical substances until area is completely dry	Dithol Kit: Package contains (1) 1.5% diclofenac sodium topical solution and (2) menthol 10% topical liquid as a roll-on Dimentho: Package contains (1) 1.5% diclofenac sodium topical solution, (2) menthol 10% topical liquid, and (3) kinesiology tape (20 strips)
diclofenac sodium/ menthol/ tape (Diclovix M)	40 drops diclofenac sodium per knee (applying 10 drops at a time), 4 times daily and spread evenly around knee; Co-packaged gel should be applied directly onto affected area without the need to bandage; repeat if necessary, but do not apply more than 4 times daily.	Wash and dry hands after use	Diclovix M kit: Diclovix kit: package contains (1) 1.5% diclofenac sodium topical solution and (2) ActivICE gel containing 8% menthol and (3) kinesiology tape

Dosages: Topical NSAIDs (continued)

Drug	Adult Dosage	Special Handling and Disposal	Availability
diclofenac sodium/menthol/methyl salicylate (DicloPR)	<p>Lower extremities: Apply 4 g to the affected area 4 times daily</p> <p>Upper extremities: Apply 2 g to the affected area 4 times daily</p> <p>Apply menthol/methyl salicylate to affected area(s) no more than 3 to 4 times daily</p>	<p>Do not apply more than 16 g daily to any 1 affected joint of the lower extremities</p> <p>Do not apply more than 8 g daily to any 1 affected joint of the upper extremities</p> <p>Do not exceed 32 g per day over all affected joints</p> <p>Patient should wash hands after use, unless the hands are the treated joint; in which case wait at least 1 hour after the application before washing</p> <p>Showering/bathing should be avoided for at least 1 hour after the application</p> <p>The gel should be measured using the dosing card supplied and can be used to apply the gel; card should be rinsed and dried after use</p> <p>Menthol/methyl salicylate: Do not use on wounds or damaged skin; do not use with a heating pad; avoid contact with eyes or mucous membranes</p>	Package contains (1) diclofenac sodium 1% gel and (2) Amplify Relief MM cream, containing menthol 10% and methyl salicylate 30%
diclofenac sodium/menthol/methyl salicylate (Varophen)	<p>Diclofenac: apply 40 drops to a knee 4 times a day</p> <p>Apply menthol/methyl salicylate to affected area(s) no more than 3 to 4 times daily</p>	<p>Wash and dry hands after use</p> <p>Avoid contact with eyes or mucous membranes</p>	Package contains (1) diclofenac sodium 1.5% topical solution and (2) Muscle Rub cream, containing menthol 10% and methyl salicylate 15%

CLINICAL TRIALS

Studies were identified through searches performed on PubMed and review of information sent by manufacturers. Search strategy included the FDA-approved use of all drugs in this review. Randomized, comparative, controlled trials comparing agents within this class for the approved indications are considered the most relevant in this category. Studies included for analysis in the review were published in English, performed with human participants, and randomly allocated participants to comparison groups. In addition, studies must contain clearly stated, predetermined outcome measure(s) of known or probable clinical importance, use data analysis techniques consistent with the study question, and include follow-up (endpoint assessment) of at least 80% of participants entering the investigation. Despite some inherent bias found in all studies including those sponsored and/or funded by pharmaceutical manufacturers, the studies in this therapeutic class review were determined to have results or conclusions that do not suggest systematic error in their experimental study design. While the potential influence of manufacturer sponsorship and/or funding must be considered, the studies in this review have also been evaluated for validity and importance.

Oral NSAIDs

Comparative Efficacy of Non-selective NSAIDs

A number of studies have attempted to define relative efficacy of non-selective NSAIDs.^{422,423,424,425,426} These efforts have consistently found that there is generally no significant difference in the efficacy among the non-selective NSAIDs. It was found that there was no statistically significant difference in efficacy, either between non-selective NSAIDs or between a non-selective NSAID and celecoxib. Additionally, no particular non-selective NSAID was associated with increased GI risk when compared to any other non-selective NSAID.

celecoxib (Celebrex) versus diclofenac/omeprazole

Patients who used NSAIDs for arthritis and who presented with ulcer bleeding were screened for study inclusion in a randomized, double-blind trial. Once ulcers healed as determined by endoscopy, patients were randomized to receive either 200 mg of celecoxib twice daily plus daily placebo or 75 mg of diclofenac twice daily plus 20 mg of omeprazole daily for 6 months.⁴²⁷ Patients were negative for *Helicobacter pylori*. Approximately 85% of each group had osteoarthritis. In the intention-to-treat analysis, which included 287 patients, the probability of recurrent bleeding during the 6-month period was 4.9% for celecoxib patients and 6.4% for diclofenac/omeprazole patients (difference, -1.5 percentage points; 95% confidence interval (CI) for the difference, -6.8 to 3.8). The difference between the groups was not significant (p=0.60). A separate analysis of this group performed by the same investigators showed that the probability of recurrent ulcers in 6 months was 18.7% in the celecoxib group and 25.6% in the diclofenac/omeprazole group (p=0.21).⁴²⁸

celecoxib (Celebrex) versus ibuprofen versus diclofenac

A total of 8,059 patients with OA and RA were enrolled in the double-blind, randomized, controlled study of Celecoxib Long-Term Arthritis Safety Study (CLASS).⁴²⁹ A total of 4,573 patients received treatment for 6 months. Patients were randomly assigned to receive celecoxib 400 mg twice daily, ibuprofen 800 mg 3 times daily, or diclofenac 75 mg twice daily. Aspirin use (≤ 325 mg daily) for CV prophylaxis was permitted and was used by 20% of patients. Patients with active GI disease or renal, hepatic, or coagulation disorders were excluded. GI toxicity was defined as upper GI ulcers and ulcer complications including bleeding, perforation, and obstruction. For the entire patient population, the yearly incidence of upper GI complications was 0.76% and 1.45% for celecoxib and NSAIDs, respectively. The overall incidence of upper GI ulcer complications was not statistically different among the groups. When the upper GI complications data were combined with symptomatic gastroduodenal ulcers, celecoxib was found to have a lower annual incidence compared to the NSAIDs (2.08 versus 3.54%, respectively; p=0.02). For patients not taking aspirin, the yearly incidence of upper GI ulcer complications was significantly lower in the celecoxib group (0.44%) versus NSAIDs group (1.27%; p=0.04). Combining the multiple endpoints of the annualized incidence of upper GI ulcer complications and symptomatic ulcers for the patients not receiving aspirin, celecoxib group (1.4%, p=0.02) had significantly fewer events than the NSAIDs group (2.91%). In the patients taking aspirin, the annualized rate of upper GI ulcer complications for the 2 groups was similar (celecoxib, 2.01%; diclofenac, 2.12%; p=0.92). The yearly incidence of upper GI complications for patients taking aspirin was higher in both treatment groups than patients not taking aspirin. Chronic GI blood loss, GI intolerance, and renal or hepatic toxicity occurred less frequently in the celecoxib group. No difference in CV events was noted between celecoxib and NSAIDs, despite aspirin use.

celecoxib (Celebrex) versus ketoprofen

In a 6-week, randomized, double-blind, placebo-controlled trial, celecoxib 100 mg twice daily and ketoprofen 100 mg twice daily were compared in 246 patients who had active ankylosing spondylitis without peripheral synovitis.⁴³⁰ Decrease in pain and functional impairment was greater in the active treatment groups than in the placebo group, with a trend in favor of celecoxib when the 2 active treatments were compared. During treatment, epigastric pain was reported in 8, 14, and 13% of patients in the placebo, ketoprofen, and celecoxib groups, respectively.

celecoxib (Celebrex) versus naproxen

The objective of the multicenter, randomized, double-blind, placebo-controlled study was to compare the efficacy and safety of celecoxib and naproxen for the treatment of OA of the hip.⁴³¹ In the trial, 1,061 patients were randomized to receive celecoxib 100, 200, or 400 mg/day, naproxen 1,000 mg/day, or placebo for 12 weeks. Patients were evaluated at baseline, 2 to 4 days after discontinuing previous NSAID or analgesic therapy, and after 2, 6, and 12 weeks of treatment. All doses of celecoxib and naproxen significantly improved the symptoms of OA at all time points compared with placebo. In terms of pain relief and improvement in functional capacity, celecoxib 200 mg/day and 400 mg/day were similarly efficacious and were as efficacious as naproxen. Both drugs were generally well tolerated.

In a similarly designed trial, 1,003 patients with OA of the knee received celecoxib 50 mg, 100 mg, or 200 mg twice daily, naproxen 500 mg twice daily, or placebo for 12 weeks.⁴³² All celecoxib doses were efficacious compared with placebo, although celecoxib 50 mg twice daily dosage regimen was minimally effective. Improvement observed with the higher dosing regimens of celecoxib was comparable to that seen with naproxen. All doses of celecoxib and naproxen were well tolerated.

In another double-blind, parallel-group, multicenter study, 537 patients with OA or RA were randomized to treatment with celecoxib 200 mg or naproxen 500 mg twice daily for 12 weeks.⁴³³ The 2 agents produced similar improvements in Patient's and Physician's Global Assessments of arthritis efficacy. Incidence of adverse events and withdrawal rates did not differ significantly between treatments. Celecoxib produced a significantly lower incidence rate of both gastric ($p < 0.001$) and duodenal ($p < 0.030$) ulcers.

celecoxib (Celebrex) versus naproxen (Naprosyn) versus diclofenac (Voltaren)

A total of 13,274 OA patients were randomly assigned to treatment with celecoxib 100 mg, celecoxib 200 mg, or nonselective NSAID therapy (diclofenac 50 mg or naproxen 500 mg) twice daily for 12 weeks.⁴³⁴ In the double-blind trial, results from all primary efficacy assessments showed that both dosages of celecoxib were as effective as NSAIDs in treating OA. Significantly more ulcer complications (adjudication based on lesion) occurred within the nonselective NSAID group (0.8/100 patient-years) compared with the celecoxib group (0.1/100 patient-years; OR 7.02; 95% CI, 1.46 to 33.80, $p = 0.008$). The number of CV thromboembolic events was low and not statistically different between the groups.

celecoxib (Celebrex) versus diclofenac (Zorvolex) in acute pain

A randomized, double-blind, placebo-controlled, parallel-arm, single center study in 428 patients with moderate-to-severe pain following bunionectomy evaluated patients randomized to diclofenac 18 mg or 35 mg 3 times daily, celecoxib 200 mg twice daily after a 400-mg loading dose, or placebo.⁴³⁵ At 12 hours, pain intensity with diclofenac 18 and 35 mg was reduced from baseline by 48% and 51%,

respectively, compared to 24% in the placebo group. At 24 hours, pain intensity was reduced by 69%, 73%, versus 52%, in diclofenac 18 and 35 mg versus placebo. All of these differences from placebo were statistically significant. Compared with celecoxib, overall reductions in pain intensity were greater with diclofenac 35 mg and similar with diclofenac 18 mg.

celecoxib (Celebrex) versus other NSAIDs in acute pain

Celecoxib has been studied in numerous head-to-head trials with other NSAIDs, such as ibuprofen, ketoprofen, and naproxen, in the treatment of various acute injuries, such as shoulder tendonitis/bursitis, ankle sprain, and tonsillectomy.^{436,437,438,439,440} Efficacy between celecoxib and the NSAIDs was generally found to be comparable, with no clinical difference in the incidence of adverse effects.

celecoxib (Celebrex) versus ibuprofen (Motrin) versus naproxen (Naprosyn) – CV Outcomes

A total of 24,081 patients with OA or RA were randomized to celecoxib, naproxen, or ibuprofen for a mean of 20 months (mean follow-up of 34 months) to assess for differences in CV risk between the agents.⁴⁴¹ In the intent-to-treat (ITT) group, the primary composite outcome of CV death, nonfatal MI, or nonfatal stroke occurred at similar rates in each group (2.3% for celecoxib, 2.5% for naproxen, and 2.7% for ibuprofen; hazard ratio [HR] for celecoxib versus naproxen, 0.93 [95% CI, 0.76 to 1.13]; HR for celecoxib versus ibuprofen, 0.85 [95% CI, 0.7 to 1.04]; $p < 0.001$ for noninferiority in both comparisons). However, the risk of GI event was significantly lower with celecoxib versus naproxen ($p = 0.01$) or ibuprofen ($p = 0.002$). Likewise, the risk of renal events was significantly lower with celecoxib than with ibuprofen ($p = 0.004$) but was not lower with celecoxib than with naproxen ($p = 0.19$).

diclofenac sodium/misoprostol (Arthrotec) versus diclofenac

Diclofenac was compared to the combination of diclofenac and misoprostol for efficacy, safety, and incidence of endoscopic upper GI ulcers in a 6 week, double-blind trial enrolling 572 patients with OA and a history of ulcers or erosions.⁴⁴² Patients were randomized to diclofenac 75 mg twice daily, diclofenac 50 mg/misoprostol 200 mcg 3 times daily, diclofenac 75 mg/misoprostol 200 mg twice daily, or placebo. All active treatment groups were more effective than placebo in relieving arthritis symptoms. Following the 6 week course of treatment, endoscopic ulcer rates (both gastric and duodenal ulcers) were as follows: diclofenac monotherapy (17%), diclofenac 50 mg with misoprostol (8%), diclofenac 75 mg with misoprostol (7%), and placebo (4%). A higher incidence of flatulence was observed in the diclofenac 75 mg with misoprostol group, whereas diclofenac 50 mg with misoprostol had a higher incidence of diarrhea.

A double-blind, randomized, parallel-group study was conducted to compare the safety and efficacy of a fixed combination of diclofenac 50 mg and misoprostol 200 mcg with a combination of diclofenac 50 mg and placebo in 361 patients with osteoarthritis.⁴⁴³ Patients with no significant gastroduodenal lesions were enrolled and received study medication 2 or 3 times daily for 4 weeks. Post-treatment endoscopic examination of the gastroduodenal mucosa revealed ulcers in 4% of patients in the diclofenac/placebo group compared with none in the diclofenac/misoprostol group ($p = 0.015$). There were no clinically or statistically significant differences between the 2 treatment groups in formal assessments of OA after either 2 or 4 weeks. Discontinuation of study drug due to adverse events was similar in each group (diclofenac/misoprostol group $n = 11$, diclofenac/placebo group $n = 10$). Eight patients in each group discontinued due to GI adverse events.

Similarly, another double-blind, randomized, parallel-group study compared the efficacy of diclofenac 50 mg/misoprostol 200 mcg or diclofenac 50 mg/placebo in treating the signs and symptoms of RA.⁴⁴⁴ A total of 346 patients with RA who had been stabilized on diclofenac for at least 30 days were randomly assigned to receive either combination for 12 weeks. Diclofenac 50 mg/misoprostol 200 mcg demonstrated no statistically significant difference in efficacy in the treatment of the signs and symptoms of RA compared with diclofenac 50 mg/placebo.

diclofenac sodium/misoprostol (Arthrotec) versus nabumetone

In a 6-week trial, diclofenac sodium 75 mg with misoprostol 200 mcg twice daily was compared to nabumetone 1,500 mg once daily or placebo for ulcer rates in 1,203 patients with symptomatic OA of the hip or knee.⁴⁴⁵ All patients enrolled had a history of endoscopically proven ulcers or erosions. Patients were evaluated by endoscopy at baseline, at withdrawal, or at the end of the 6 week time period. The incidence of duodenal and gastric ulcers confirmed with endoscopy was significantly lower in diclofenac/misoprostol group (4%) compared to nabumetone (11%). Duodenal ulcers were similar between the 2 active treatments. Gastric ulcers were significantly less with diclofenac/misoprostol (1%) compared to nabumetone (9%). Types of adverse events were similar for all treatment groups, with GI adverse events predominating. Diclofenac sodium 75 mg with misoprostol 200 mcg was well tolerated by the majority of patients. Withdrawals due to adverse effects were reported as 13% of patients in the diclofenac/misoprostol group, 10% in the nabumetone group, and 9% in the placebo group.

diclofenac (Zorvolex) versus placebo in OA

A randomized, double-blind, parallel-group, placebo-controlled, 12-week, multicenter trial evaluated 305 patients with osteoarthritis of the hip or knee.⁴⁴⁶ Patients were randomized to diclofenac 35 mg 3 times daily, 35 mg twice daily, or placebo. Efficacy parameters included mean change from baseline in Western Ontario and McMaster Universities Arthritis Index (WOMAC) pain subscale score at week 12 (primary efficacy parameter) and in average total WOMAC score over 12 weeks. Submicron diclofenac 35 mg 3 times daily for 12 weeks significantly improved (WOMAC) pain subscale scores from baseline at 12 weeks (-44.1; p=0.0024) compared with placebo (-32.5). The twice-daily regimen was not significantly better (-39; p=0.0795) than placebo. Submicron diclofenac 35 mg 3 times daily (-35.9; p=0.0002) and 35 mg twice daily (-30.3; p=0.0363) improved the average total WOMAC score in patients over 12 weeks compared with placebo (-23.2). Diarrhea, headache, nausea, and constipation were the most common adverse events in the submicron diclofenac groups.

ibuprofen/famotidine (Duexis) versus ibuprofen

Ibuprofen/famotidine was compared to ibuprofen in 2 multicenter, double-blind, active-controlled, randomized studies in patients who were expected to require daily administration of an NSAID for at least 6 months for conditions such as the following: OA, RA, chronic low back pain, chronic regional pain syndrome, and chronic soft tissue pain.⁴⁴⁷ Duexis is FDA approved only for the treatment of OA and RA. The studies compared the incidence of upper GI (gastric and/or duodenal) ulcer formation in a total 1,533 patients, either as a primary or secondary endpoint. Patients were assigned in a 2:1 ratio to ibuprofen/famotidine (800 mg/26.6 mg) or ibuprofen (800 mg) 3 times a day for 24 consecutive weeks. Patient age ranged from 39 to 80 years (median age 55 years). Approximately 15% of the patients in both studies were taking concurrent low-dose aspirin (< or equal to 325 mg daily) and 6% had a history of previous upper gastrointestinal ulcer. *Helicobacter pylori* status was negative at baseline; however,

H. pylori status was not reassessed during the trials. In both trials ibuprofen/famotidine was associated with a statistically significant reduction in the risk of developing upper GI ulcers compared with ibuprofen alone. Each endpoint was analyzed in 2 fashions. In 1 analysis patients who terminated early, without an endoscopic evaluation within 14 days of their last dose of study drug, were classified as not having an ulcer. This analysis reported GI ulcer in 17.4 to 18.6% of patients in the ibuprofen/famotidine group, compared to 31 to 34.3% of patients in the ibuprofen group ($p < 0.0001$). In the second analysis, those patients were classified as having an ulcer, which reported GI ulcer in 8.7 to 10.1% of patients in the ibuprofen/famotidine group, compared to 17.6 to 21.3% of patients in the ibuprofen group ($p \leq 0.0004$). The results of the patients that used low-dose aspirin were consistent with the overall findings of the study. In these clinical studies, 23% of patients 65 years of age and older who were treated with ibuprofen/famotidine developed an upper GI ulcer compared to 27% of those patients who received only ibuprofen. In addition, 25% of patients with a prior history of GI ulcer who were treated with ibuprofen/famotidine developed an upper GI ulcer compared to 24% of those patients who received ibuprofen only.

indomethacin (Tivorbex) versus placebo

Efficacy of indomethacin (Tivorbex) was demonstrated for treatment of acute pain based on 2 phase 3 randomized, double-blind, placebo-controlled, parallel-arm, multicenter, studies that compared 20 mg of the drug 3 times daily, 40 mg 2 times daily, 40 mg 3 times daily, and placebo taken by patients with pain following bunionectomy.⁴⁴⁸ There were a total of 835 patients, randomized equally across the treatment groups, enrolled in the 2 studies. The patients were a mean age of 40 years, ranging from 18 to 68 years, with at least a pain intensity rating of 40 mm on a 100 mm visual analog scale (VAS) during the 9 hour period after anesthetic block was discontinued following the surgery. Mean pain intensity in both studies ranged from 71 to 74 mm. A tablet of 10 mg/325 mg hydrocodone/acetaminophen was allowed every 4 to 6 hours as rescue medication. The use of rescue medication was greater in placebo-treated patients than in Tivorbex-treated patients. In both studies, all 3 Tivorbex dosages demonstrated efficacy in reduction of pain intensity compared to placebo, as measured by the sum of pain intensity difference over 0 to 48 hours after the first dose.

meloxicam submicronized (Vivlodex) versus placebo

A multicenter, randomized, double-blind, parallel-arm, placebo-controlled trial evaluated the safety and efficacy of meloxicam (Vivlodex) for the treatment of osteoarthritis pain of the hip or knee in adult patients ($n=402$).^{449,450} Adult patients were assigned meloxicam 5 mg, meloxicam 10 mg, or placebo orally once daily. The primary efficacy endpoint was the change in the Western Ontario and McMaster University Osteoarthritis Index (WOMAC) Pain Subscale score from baseline to 12 weeks. Both meloxicam doses (5 mg and 10 mg) significantly reduced pain compared to placebo at 12 weeks.

naproxen/esomeprazole (Vimovo) versus naproxen enteric coated (EC)

The manufacturer performed 2 randomized, multicenter, double-blind studies comparing the incidence of gastric ulcer formation in patients with medical conditions expected to require daily NSAID therapy for at least 6 months.^{451,452} If patients ($n=854$) were < 50 years old, they required documented history of gastric or duodenal ulcer within the past 5 years. Patients received naproxen/esomeprazole 500 mg/20 mg twice daily or enteric-coated naproxen 500 mg twice daily. About one quarter of the patients were taking low-dose aspirin also. Naproxen/esomeprazole patients showed statistically

significant reductions in the 6 month cumulative incidence of gastric ulcers compared to naproxen EC (4.1–7.1% of patients with gastric ulcer versus 23.1–24.3%, respectively; $p < 0.001$).

The manufacturer performed two, 12-week, randomized, double-blind, placebo-controlled studies to determine effectiveness of naproxen/esomeprazole in treating the signs and symptoms of OA of the knee.⁴⁵³ Patients receiving naproxen/esomeprazole 500 mg/20 mg twice daily had significantly better results compared to placebo as measured by Western Ontario and McMaster Universities Arthritis Index (WOMAC) pain subscale and WOMAC physical function subscale, as well as a Patient Global Assessment score.

lansoprazole (Prevacid) versus misoprostol (Cytotec)

A prospective, double-blind, multicenter, active- and placebo-controlled study evaluated 537 patients without *H. pylori* who were long-term users of NSAIDs and who had a history of gastric ulcer documented by endoscopy.⁴⁵⁴ Patients were randomized to receive placebo, misoprostol 200 mcg 4 times a day, or lansoprazole 15 or 30 mg once daily for 12 weeks. Patients receiving lansoprazole (15 or 30 mg) remained free from gastric ulcer longer than those who received placebo ($p < 0.001$), but for a shorter time than those who received misoprostol. By week 12, the percentages of gastric ulcer-free patients were as follows: placebo, 51%; misoprostol, 93%; lansoprazole 15 mg, 80%; and lansoprazole 30 mg, 82%. A significantly higher proportion of patients in the misoprostol group reported treatment-related adverse events and early withdrawal from the study. Therapy was successful for 69% of each active treatment group and 35% for the placebo group. Lansoprazole was superior to placebo for the prevention of NSAID-induced gastric ulcers but was not superior to misoprostol 800 mcg per day. When the poor compliance and potential adverse effects associated with misoprostol are considered, however, proton pump inhibitors (PPIs) and full-dose misoprostol are clinically equivalent.^{455,456}

Nasal NSAIDs

ketorolac nasal spray (Sprix) versus placebo

Ketorolac nasal spray was studied in a phase 3, randomized, multicenter, double-blind, placebo-controlled trial of 300 adults who had elective abdominal or orthopedic surgery.⁴⁵⁷ Post-operatively, patients were treated with morphine dosed via patient controlled analgesia (PCA) on an as-needed basis. They were then randomized to the addition of ketorolac nasal spray or placebo, administered every 8 hours for 48 hours. Patients in the ketorolac nasal spray arm had a significantly reduced summed pain intensity difference over 48 hours compared to those in the placebo group. Patients in the ketorolac nasal spray arm required 36% less morphine over 48 hours than those treated with placebo.

A second phase 3, multicenter, double-blind, placebo-controlled study randomized 321 patients who had elective abdominal surgery to treatment with ketorolac nasal spray or placebo.⁴⁵⁸ Post-operatively, patients were treated using morphine PCA on an as-needed basis. In addition, ketorolac nasal spray or placebo was administered every 6 hours for 48 hours. Patients in the ketorolac nasal spray group had a significantly greater reduction in summed pain intensity difference over 48 hours compared to those in the placebo group. Patients treated with ketorolac nasal spray required 26% less morphine over 48 hours compared to those in the placebo group.

Topical NSAIDs

diclofenac patch (Flector) versus placebo

A randomized, double-blind, multicenter, placebo-controlled trial was conducted in 120 patients with traumatic soft tissue injury within 3 hours post-injury.⁴⁵⁹ Patients were randomized to twice daily treatment with either diclofenac patch or placebo over a period of 7 days. The primary efficacy endpoint was the area under the curve (AUC) for tenderness over the first 3 days. The diclofenac patch was significantly more effective than placebo ($p < 0.0001$). The diclofenac patch produced rapid pain relief as reflected by the time to reach resolution of pain at the injured site, which was significantly shorter compared to placebo ($p < 0.0001$). The most frequently observed adverse events with the use of diclofenac patch were mild, local cutaneous adverse events, occurring at the same frequency as placebo.

A multicenter, randomized, placebo-controlled, parallel-design study was conducted to assess the efficacy and safety of diclofenac patch applied directly to the injury site for the treatment of acute minor sports injury pain in 222 adult patients within 72 hours of the injury.⁴⁶⁰ Either a diclofenac or placebo topical patch was applied directly to the skin overlying the injured site twice daily for 2 weeks. Measures of pain intensity were performed in a daily diary and at clinic visits on days 3, 7, and 14. Diclofenac patch was superior to placebo patch in relieving pain. Statistical significance was seen on clinic days 3 ($p = 0.036$) and 14 ($p = 0.048$), as well as the daily diary pain ratings at days 3, 7, and 14 ($p \leq 0.044$). No statistically significant differences were seen in any safety or adverse effect measures with the diclofenac patch as compared to the placebo patch.

diclofenac solution (Pennsaid) versus placebo

Patients ($n = 248$) with osteoarthritis of the knee and at least moderate pain were randomly assigned to apply 1 solution to their painful knee for 4 weeks: diclofenac solution 1.5%, vehicle solution, or placebo solution.⁴⁶¹ The primary efficacy endpoint was pain relief, measured by the Western Ontario and McMaster Universities (WOMAC) LK3.0 Osteoarthritis Index pain subscale. In the intent-to-treat group, the mean change in pain score from baseline to final assessment was significantly greater for the patients who applied the diclofenac solution (-3.9 ; 95% confidence interval [CI], -4.8 to -2.9) than for those who applied the vehicle solution (-2.5 ; 95% CI, -3.3 to -1.7 ; $p = 0.023$) or the placebo solution (-2.5 ; 95% CI, -3.3 to -1.7 ; $p = 0.016$). The diclofenac solution also showed superiority to the vehicle and placebo solutions in physical function, stiffness, and in pain on walking. The Patient Global Assessment scores were significantly better for the patients who applied the diclofenac solution than for those who applied the other solutions ($p = 0.039$ and 0.025 , respectively). The diclofenac solution caused some skin irritation in 36% of patients. In a similarly designed 6-week study, diclofenac solution was again found to be superior to vehicle in 216 patients with osteoarthritis of the knee.⁴⁶² A 12-week trial in 216 patients with osteoarthritis of the knee came to the same conclusions.⁴⁶³

A 12-week, double-blind, double-dummy, randomized controlled trial was performed in 775 subjects with symptomatic primary osteoarthritis of the knee.⁴⁶⁴ This study compared diclofenac solution with a placebo solution, the vehicle solution, oral diclofenac, and the combination of oral diclofenac and diclofenac solution. Subjects applied study solutions 40 drops 4 times daily and took 1 study tablet daily for 12 weeks. Co-primary efficacy variables were WOMAC pain and physical function and a patient overall health assessment. Diclofenac solution was superior to placebo for pain (-6 versus -4.7 ; $p = 0.015$), physical function (-15.8 versus -12.3 ; $p = 0.034$), overall health (-0.95 versus -0.37 ; $p < 0.0001$),

and Patient Global Assessment (-1.36 versus -1.01; $p=0.016$), and was superior to vehicle for all efficacy variables. The most common adverse event associated with diclofenac solution was dry skin. Fewer digestive system and laboratory abnormalities were observed with diclofenac solution than with oral diclofenac.

diclofenac gel (Voltaren Gel) versus placebo

In a randomized, double-blind, placebo-controlled trial, 385 patients with primary osteoarthritis in the dominant hand were assigned to diclofenac 1% gel or vehicle to both hands 4 times daily for 8 weeks.⁴⁶⁵ Primary outcome measures included osteoarthritis pain intensity (100 mm visual analog scale), total Australian/Canadian Osteoarthritis Hand Index (AUSCAN) score, and global rating of disease activity at 4 and 6 weeks. Diclofenac gel decreased pain intensity scores by 42 to 45%, total AUSCAN scores by 35 to 40%, and global rating of disease by 36 to 40%. Significant differences favoring diclofenac gel over vehicle were observed at week 4 for pain intensity and AUSCAN. At week 6, diclofenac gel significantly improved each primary outcome measure compared with vehicle. Secondary outcomes generally supported the primary outcomes. The most common adverse event was application site paresthesia.

In a randomized, double-blind, vehicle-controlled trial, 492 adults with symptomatic knee osteoarthritis were randomized to diclofenac gel 1% or vehicle 4 times daily for 12 weeks.⁴⁶⁶ Primary efficacy outcomes at week 12 were the WOMAC pain subscale, WOMAC physical function subscale, and global rating of disease. At week 12, the diclofenac gel group had significant decreases versus the vehicle group in mean WOMAC pain ($p=0.01$), mean WOMAC physical function ($p=0.001$), and mean global rating of disease ($p<0.001$). Efficacy outcomes significantly favored diclofenac gel versus vehicle beginning at week 1. Application site reactions occurred in 5.1 and 2.5% of patients in the diclofenac gel and vehicle groups, respectively.

SUMMARY

The available clinical data do not suggest that any one nonsteroidal anti-inflammatory drug (NSAID) offers a clear advantage compared to the others in terms of safety or efficacy, given the complex tradeoffs between the many benefits (e.g., pain relief, improved function, and improved tolerability) and harms (e.g., cardiovascular [CV], renal, and gastrointestinal [GI]) involved. Adequate pain relief at the expense of an increase in CV risk could be an acceptable trade-off for some patients. Others may consider even a marginal increase in CV risk unacceptable. When weighing the potential effects of any of these agents, the following patient factors should be considered prior to initiation of therapy: age, comorbid conditions, and concomitant medications. NSAIDs should be used in the lowest effective dose.

The addition of misoprostol to diclofenac sodium (Arthrotec) in an attempt to reduce GI ulcers is efficacious, but many patients have difficulty tolerating the GI adverse effects, including diarrhea, associated with misoprostol. Esomeprazole/naproxen (Vimovo) has been approved to relieve the signs and symptoms of osteoarthritis (OA), rheumatoid arthritis (RA), and ankylosing spondylitis (AS) and to decrease the risk of stomach (gastric) ulcers in patients at risk of developing stomach ulcers from treatment with NSAIDs. Data are available that support use of any proton pump inhibitor with concurrent NSAID administration. Ibuprofen/famotidine (Duexis) is the newest combination agent in this class and is indicated for the relief of signs and symptoms of OA and RA and to decrease the risk of developing upper GI ulcers.

Ketorolac nasal spray (Sprix) offers an alternative method of drug delivery.

The topical NSAIDs are indicated for treatment of acute pain conditions, including strains and sprains, as well as chronic pain conditions like osteoarthritis. For patients at risk for GI or CV events, topical administration of diclofenac (Flector/Licart, Pennsaid, Voltaren gel, and diclofenac-containing kits) provides an alternative method of drug delivery.

Multiple products in this class are marketed as kits, which most commonly contain an oral or topical NSAID that is already available as a single agent co-packaged with another non-prescription topical agent (e.g., capsaicin or menthol).

REFERENCES

- 1 Celebrex [package insert]. New York, NY; Pfizer; May 2019.
- 2 Diclofenac potassium [package insert]. Morgantown, WV; Mylan; August 2017.
- 3 Zipsor [package insert]. Newark, CA; DepoMed; May 2016.
- 4 Diclofenac sodium tablet, delayed release [package insert]. Parsippany, NJ; Actavis; October 2017.
- 5 Diclofenac sodium tablet, film coated, extended release [package insert]. Parsippany, NJ; Actavis; June 2017.
- 6 Zorvolex [package insert]. Philadelphia, PA; Iroko; May 2016.
- 7 Diflunisal [package insert]. Sellersville, PA; Teva; July 2018.
- 8 Etodolac [package insert]. Hawthorne, NY; Taro; December 2015.
- 9 Nalfon [package insert]. Ridgeland, MS; Xspire; May 2016.
- 10 Flurbiprofen [package insert]. Morgantown, WV; Mylan; May 2016.
- 11 Ibuprofen tablet [package insert]. Bridgewater, NJ; Amneal; May 2019.
- 12 Indocin [package insert]. Philadelphia, PA; Iroko; March 2019.
- 13 Tivorbex [package insert]. Philadelphia, PA; Iroko; March 2019.
- 14 Ketoprofen IR [package insert]. Morgantown, WV; Teva; September 2018.
- 15 Ketoprofen ER [package insert]. North Wales, PA; Teva; May 2017.
- 16 Ketorolac tromethamine tablet [package insert]. Morgantown, WV; Mylan; July 2015.
- 17 Meclofenamate [package insert]. Morgantown, WV; Mylan; July 2015.
- 18 Mefenamic acid [package insert]. Basking Ridge, NJ; Micro; January 2020.
- 19 Mobic [package insert]. Ridgefield, CT; Boehringer-Ingelheim; June 2016.
- 20 Qmiiz ODT [package insert]. Lake Forest, IL; TerSera; October 2018.
- 21 Vivlodex [package insert]. Philadelphia, PA; Iroko; October 2015.
- 22 Nabumetone [package insert]. Parsippany, NJ; Actavis; May 2016.
- 23 Relafen DS [package insert]. Hazlet, NJ; Carwin; January 2020.
- 24 Naprelan [package insert]. San Diego, CA; Victory; July 2019.
- 25 Anaprox/Anaprox DS [package insert]. Alpharetta, GA; Canton; July 2019.
- 26 Daypro [package insert]. New York, NY; Pfizer; May 2019.
- 27 Feldene [package insert]. New York, NY; Pfizer; May 2019.
- 28 Sulindac [package insert]. Parsippany, NJ; Actavis; May 2016.
- 29 Tolmetin [package insert]. Morgantown, WV; Mylan; July 2015.
- 30 ANDA Approval Letter. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/applletter/2019/203166Orig1s000ltr.pdf. Accessed June 22, 2020.
- 31 NuDroxiPak Pain Relief Pak [package insert]. NuCare Pharmaceuticals. September 2019. Available at: <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=64bc6acc-58bf-0881-e053-2a91aa0a35e4&audience=consumer>. Accessed June 2, 2020.
- 32 Flexipak [package insert]. Ripley, MS; Sterling-Knight; January 2018.
- 33 Inavix [package insert]. Ocean Springs, MS; Primary; July 2020.
- 34 Inflammacin [package insert]. San Fernando, CA; PureTek; April 2019. Available at: <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=91d25301-fac9-41f0-8a4e-e22d44926347&audience=consumer>. Accessed June 2, 2020.
- 35 Nudiclo TabPak [package insert]. NuCare; October 2019. Available at: <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=c1f0ff98-c000-40a9-b770-da5813647dd8>. Accessed June 2, 2020.
- 36 Xenafamm [package insert]. Panorama City, CA; Shoreline; May 2016.
- 37 NuDroxiPak DSDR-75 [package insert]. NuCare Pharmaceuticals. February 2018. Available at: <https://dailymed.nlm.nih.gov/dailymed/search.cfm?labeltype=all&query=nudroxipak&pagesize=200&page=1&audience=consumer>. Accessed June 22, 2020.
- 38 Inflattherm (diclofenac/menthol/camphor) [package insert]. Portland, OR. V2 Pharma. July 2019.
- 39 Arthrotec [package insert]. New York, NY; Pfizer; May 2016.
- 39 NuDroxiPak Pain Relief Pak [package insert]. NuCare Pharmaceuticals. February 2018. Available at: <https://dailymed.nlm.nih.gov/dailymed/search.cfm?labeltype=all&query=nudroxipak&pagesize=200&page=1&audience=consumer>. Accessed June 22, 2020.
- 41 Vimovo [package insert]. Deerfield, IL; Horizon; June 2018.

- 42 Duexis [package insert]. Deerfield, IL; Horizon; July 2019.
- 43 Comfort Pac with ibuprofen [package insert]. Oklahoma City, OK. PD-RX. October 2017.
- 44 NuDroxiPak DSDR-75 [package insert]. NuCare Pharmaceuticals. February 2018. Available at: <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=678aad78-a2b7-f7c9-e053-2a91aa0a4f26&audience=consumer>. Accessed June 2, 2020.
- 45 Comfort Pac with meloxicam [package insert]. Oklahoma City, OK. PD-RX. October 2017.
- 46 Comfort Pac with naproxen [package insert]. Oklahoma City, OK. PD-RX. October 2017. Available at: <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=91802fa6-ddf3-4356-85f0-2032cf31478b&audience=consumer>. Accessed June 22, 2020.
- 47 NuDroxiPak N-500 [package insert]. NuCare Pharmaceuticals. March 2018. Available at: <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=67903a9b-b94e-1f65-e053-2991aa0a0e43>. Accessed June 2, 2020.
- 48 Sprix [package insert]. Wayne, PA; Egalet; January 2018.
- 49 Flector [package insert]. New York, NY; Pfizer; March 2019.
- 50 Licart [package insert]. Parsippany, NJ; IBSA; May 2020.
- 51 Lexixryl [package insert]. Panorama City, CA; Shoreline; February 2017.
- 52 Diclofenac sodium solution 1.5% [package insert]. Parsippany, NJ; Actavis; May 2016.
- 53 Pennsaid 2% [package insert]. Lake Forest, IL; Horizon; May 2016.
- 54 Xrylix [package insert]. San Fernando, CA; PureTek; March 2016.
- 55 Voltaren Gel [package insert]. Malvern, PA; Endo; February 2018.
- 56 Diclozor [package insert]. Ripley, MS; Sterling Knight; October 2016.
- 57 Diclovix [package insert]. Ocean Springs, MS; Primary; March 2019. Available at: <https://www.dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=84b45252-fdf6-0ab0-e053-2991aa0ac428>. Accessed June 3, 2020.
- 58 Diclotrex [package insert]. San Fernando, CA; Purtek; October 2018.
- 59 Inflamm-K Kit [package insert]. Peoria, AZ; Solutech; August 2016.
- 60 CapsFenac Pak [package insert]. Los Angeles, CA; SA3; January 2020. Available at: <https://www.dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=88728f7a-19e5-4f31-911d-fc7d8a64ba0d>. Accessed June 3, 2020.
- 61 Capsinac [package insert]. Las Vegas, NV; Bodysphere; July 2015.
- 62 DermacinRx Lexitral PharmaPak [package insert]. San Fernando, CA; PureTek; March 2019.
- 63 Diclofex DC [package insert]. Madison, MS; Sircle; November 2019. Available at: <https://www.dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=d8ba9e7c-3e5f-4c41-add9-3e1a142c1fff>. Accessed June 3, 2020.
- 64 Diclopak [package insert]. Ripley, MS; Sterling-Knight; January 2018.
- 65 Diclotral Pak [package insert]. Everett, WA; Patchwerx; March 2016. Available at: <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=2d8f4048-14cb-7484-e054-00144ff88e88&audience=consumer>. Accessed June 3, 2020.
- 66 Nudiclo SoluPak [package insert]. Orange, CA; Nucare; December 2017. Available at: <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=d8105efe-53e9-4392-ac13-7d65fa31211e>. Accessed June 3, 2020.
- 67 Diclosiacin [package insert]. Madison, MS; Chadwick; January 2020.
- 68 Sure Result DSS Premium Pak [package insert]. Doral, FL; International Brand Management; March 2016.
- 69 Xelitral [package insert]. Panorama City, CA; Shoreline; February 2017.
- 70 Trixylylral. San Fernando, CA; Shoreline. Available at: <https://www.drugs.com/pro/trixylitral.html>. Accessed June 3, 2020.
- 71 Dithol [package insert]. Fortus. December 2017.
- 72 Dimentho [package insert]. Biloxi, MS; Levins; October 2019.
- 73 Diclovix M [package insert]. Ocean Springs, MS; Primary; January 2020. Available at: <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=9d0012ed-a50a-a936-e053-2995a90a6100>. Accessed June 8, 2020.
- 74 DicloPR [package insert]. Tampa, FL; 19&Pacific; November 2017.
- 75 Varophen [package insert]. Portland, OR. V2 Pharma. June 2019.
- 76 Nonsteroidal Anti-inflammatory Drug (NSAID) Toxicity. Medscape. Available at: <https://emedicine.medscape.com/article/816117-overview>. Accessed June 16, 2020.
- 77 El-Serag HB, Graham DY, Richardson P, et al. Prevention of complicated ulcer disease among chronic users of nonsteroidal anti-inflammatory drugs: the use of a nomogram in cost-effectiveness analysis. Arch Intern Med. 2002; 162:2105-2110.
- 78 Ruffalo RL, Jackson RL, Ofman JJ. The impact of NSAID selection on gastrointestinal injury and risk for cardiovascular events: identifying and treating patients at risk. P & T. 2002; 27:570-576.
- 79 Available at: <https://www.fda.gov/downloads/Drugs/DrugSafety/UCM453941.pdf>. Accessed June 3, 2020.
- 80 Singh G. Recent considerations in nonsteroidal anti-inflammatory drug gastropathy. Am J Med. 1998; 105(Suppl 1B):315-385.
- 81 Cordero JA, Alarcon L, Escnbano E, et al. A comparative study of the transdermal penetration of a series of nonsteroidal anti-inflammatory drugs. J Pharm Sciences. 2002; 86(24): 503-8.
- 82 Kolasinski SL, Neogi T, Hochberg MC, et al. 2019 American College of Rheumatology/Arthritis Foundation guideline for the management of osteoarthritis of the hand, hip, and knee. Arthrit Car Resear. 2020; 72(2): 149-162. DOI: 10.1002/acr.24131 Available at: <https://www.rheumatology.org/Practice-Quality/Clinical-Support/Clinical-Practice-Guidelines/Osteoarthritis>. Accessed June 3, 2020.
- 83 American Academy of Orthopedic Surgeons. Treatment of Osteoarthritis of the Knee, Evidence-Based Guideline. Second Edition. 2013. Available at: <https://www.aaos.org/quality/quality-programs/lower-extremity-programs/osteoarthritis-of-the-knee/>. Accessed June 3, 2020.
- 84 American Academy of Orthopaedic Surgeons. Management of Osteoarthritis of the Hip Evidence-Based Clinical Practice Guideline. March 2017. Available at: <https://www.aaos.org/quality/quality-programs/lower-extremity-programs/osteoarthritis-of-the-hip/?ssopc=1>. Accessed June 3, 2020.
- 85 Qaseem A, Wilt TJ, McLean RM, et al. Noninvasive treatments for acute, subacute, and chronic low back pain: A clinical practice guideline from the American College of Physicians. 2017. Ann Intern Med. 2017; 166(7): 514-530. DOI: 10.7326/M16-2367. Available at: <https://www.acponline.org/clinical-information/guidelines>. Accessed June 3, 2020.
- 86 Qaseem A, Harris RP, Forciea MA, et al. Management of Acute and Recurrent Gout: A Clinical Practice Guideline from the American College of Physicians. 2016 Ann Intern Med. 2017;166:58-68. DOI: 10.7326/M16-0570. Available at: <https://www.acponline.org/clinical-information/guidelines>. Accessed June 3, 2020.

-
- 87 FitzGerald JD, Dalbeth N, Mikuls T, et al. 2020 American College of Rheumatology Guideline for the Management of Gout. *Arthritis Care Res.* 2020; 72(6): 744-760. DOI: 10.1002/acr.24180. Epub 2020 May 11. Available at: <https://www.rheumatology.org/Practice-Quality/Clinical-Support/Clinical-Practice-Guidelines/Gout>. Accessed June 22, 2020
- 88 Crofford LJ. COX-1 and COX-2 tissue expression: Implications and predictions. *J Rheumatol.* 1997; 24(Suppl 49):15-19.
- 89 Needleman P, Isakson PC. The discovery and function of COX-2. *J Rheumatol.* 1997; 24(Suppl 49):6-8.
- 90 Lipsky PE, Isakson PC. Outcome of specific COX-2 inhibition in rheumatoid arthritis. *J Rheumatol.* 1997; 24(Suppl 49):9-14.
- 91 Lane NE. Pain management in osteoarthritis: The role of COX-2 inhibitors. *J Rheumatol.* 1997; 24(Suppl 49):20-24.
- 92 Ziegler J. Cancer and arthritis share underlying processes. *J Natl Cancer Inst.* 1998; 90(11):802-803.
- 93 Vivlodex [package insert]. Philadelphia, PA; Iroko; October 2015.
- 94 Vimovo [package insert]. Deerfield, IL; Horizon; July 2019.
- 95 Arthrotec [package insert]. New York, NY; Pfizer; May 2016.
- 96 Duexis [package insert]. Deerfield, IL; Horizon; July 2019.
- 97 Clinical Pharmacology. Available at: <http://www.clinicalpharmacology-ip.com/default.aspx>. Accessed June 22, 2020.
- 98 Methyl salicylate and menthol cream and ointment. Available at: <https://www.drugs.com/cdi/methyl-salicylate-menthol-cream.html>. Accessed June 22, 2020.
- 99 Crofford LJ. COX-1 and COX-2 tissue expression: implications and predictions. *J Rheumatol.* 1997; 24(Suppl 49):15-19.
- 100 McAdam BF, Catella-Lawson F, Mardini IA, et al. Systemic biosynthesis of prostacyclin by cyclooxygenase (COX)-2: The human pharmacology of a selective inhibitor of COX-2. *Proc Natl Acad Sci.* 1999; 96:272-277.
- 101 Lipsky PE. Specific COX-2 inhibitors in arthritis, oncology, and beyond: Where is the science headed? *J Rheumatol.* 1999; 26(Suppl 56):25-30.
- 102 Celebrex [package insert]. New York, NY; Pfizer; May 2019.
- 103 Flurbiprofen [package insert]. Morgantown, WV; Mylan; May 2016.
- 104 Zorvolex [package insert]. Philadelphia, PA; Iroko; May 2016.
- 105 Zipsor [package insert]. Newark, CA; DepoMed; May 2016.
- 106 Diclofenac potassium [package insert]. Morgantown, WV; Mylan; August 2017.
- 107 Sulindac [package insert]. Parsippany, NJ; Actavis; May 2016.
- 108 Diflunisal [package insert]. Sellersville, PA; Teva; July 2018.
- 109 Flector [package insert]. New York, NY; Pfizer; March 2019.
- 110 Licart [package insert]. Parsippany, NJ; IBSA; May 2020.
- 111 Indocin [package insert]. Philadelphia, PA; Iroko; March 2019.
- 112 Tivorbex [package insert]. Philadelphia, PA; Iroko; March 2019.
- 113 Etodolac [package insert]. Hawthorne, NY; Taro; December 2015.
- 114 Etodolac extended release [package insert]. Hawthorne, NY; Taro; April 2016.
- 115 Mobic [package insert]. Ridgefield, CT; Boehringer-Ingelheim; June 2016.
- 116 Qmii ODT [package insert]. Lake Forest, IL; TerSera; October 2018.
- 117 Ibuprofen tablet [package insert]. Bridgewater, NJ; Amneal; May 2019.
- 118 Nalfon [package insert]. Ridgeland, MS; Xspire; May 2016.
- 119 Naprelan [package insert]. San Diego, CA; Victory; July 2019.
- 120 Mefenamic acid [package insert]. Basking Ridge, NJ; Micro; January 2020.
- 121 Anaprox/Anaprox DS [package insert]. Alpharetta, GA; Canton; July 2019.
- 122 Nabumetone [package insert]. Parsippany, NJ; Actavis; May 2016.
- 123 Ketorolac tromethamine tablet [package insert]. Morgantown, WV; Mylan; July 2015.
- 124 Tolmetin [package insert]. Morgantown, WV; Mylan; July 2015.
- 125 Daypro [package insert]. New York, NY; Pfizer; May 2019.
- 126 Feldene [package insert]. New York, NY; Pfizer; May 2019.
- 127 Diclofenac sodium tablet, delayed release [package insert]. Parsippany, NJ; Actavis; October 2017.
- 128 Voltaren Gel [package insert]. Malvern, PA; Endo; February 2018.
- 129 Diclofenac sodium tablet, film coated, extended release [package insert]. Parsippany, NJ; Actavis; June 2017.
- 130 Arthrotec [package insert]. New York, NY; Pfizer; May 2016.
- 131 Meclofenamate [package insert]. Morgantown, WV; Mylan; July 2015.
- 132 Vimovo [package insert]. Deerfield, IL; Horizon; July 2019.
- 133 Duexis [package insert]. Deerfield, IL; Horizon; May 2016.
- 134 Flector [package insert]. New York, NY; Pfizer; March 2019.
- 135 Licart [package insert]. Parsippany, NJ; IBSA; May 2020.
- 136 Voltaren Gel [package insert]. Malvern, PA; Endo; February 2018.
- 137 Pennsaid 2% [package insert]. Lake Forest, IL; Horizon; May 2016.
- 138 Diclofenac sodium solution 1.5% [package insert]. Parsippany, NJ; Actavis; May 2016.
- 139 Sprix [package insert]. Wayne, PA; Egalet; January 2018.
- 140 DermacinRx Lexitral PharmaPak [package insert]. San Fernando, CA; PureTek; July 2015.
- 141 Sure Result DSS Premium Pak [package insert]. Doral, FL; International Brand Management; March 2016.
- 142 DS Prep Pak [package insert]. Ocean Springs, MS; Alvix; November 2015.
- 143 Xrylix [package insert]. San Fernando, CA; PureTek; March 2016.
- 144 Vivlodex [package insert]. Philadelphia, PA; Iroko; October 2015.
- 145 Inflamm-K Kit [package insert]. Peoria, AZ; Solutech; August 2016.
- 146 Diclozor [package insert]. Ripley, MS; Sterling Knight; October 2016.
- 147 Relafen DS [package insert]. Hazlet, NJ; Carwin; January 2020.
- 148 Flurbiprofen [package insert]. Morgantown, WV; Mylan; May 2016.
-

-
- 149 Zorvolex [package insert]. Philadelphia, PA; Iroko; May 2016.
- 150 Diclofenac potassium [package insert]. Morgantown, WV; Mylan; August 2017.
- 151 Sulindac [package insert]. Parsippany, NJ; Actavis; May 2016.
- 152 Diflunisal [package insert]. Sellersville, PA; Teva; July 2018.
- 153 Flector [package insert]. New York, NY; Pfizer; March 2019.
- 154 Licart [package insert]. Parsippany, NJ; IBSA; May 2020.
- 155 Indocin [package insert]. Philadelphia, PA; Iroko; March 2019.
- 156 Tivorbex [package insert]. Philadelphia, PA; Iroko; March 2019.
- 157 Etodolac [package insert]. Hawthorne, NY; Taro; December 2015.
- 158 Etodolac extended release [package insert]. Hawthorne, NY; Taro; April 2016.
- 159 Mobic [package insert]. Ridgefield, CT; Boehringer-Ingelheim; June 2016.
- 160 Qmii ODT [package insert]. Lake Forest, IL; TerSera; October 2018.
- 161 Ibuprofen tablet [package insert]. Bridgewater, NJ; Amneal; May 2019.
- 162 Nalfon [package insert]. Ridgeland, MS; Xspire; May 2016.
- 163 Naprelan [package insert]. San Diego, CA; Victory; July 2019.
- 164 Mefenamic acid [package insert]. Basking Ridge, NJ; Micro; January 2020.
- 165 Anaprox/Anaprox DS [package insert]. Alpharetta, GA; Canton; July 2019.
- 166 Nabumetone [package insert]. Parsippany, NJ; Actavis; May 2016.
- 167 Ketorolac tromethamine tablet [package insert]. Morgantown, WV; Mylan; July 2015.
- 168 Tolmetin [package insert]. Morgantown, WV; Mylan; July 2015.
- 169 Daypro [package insert]. New York, NY; Pfizer; May 2019.
- 170 Feldene [package insert]. New York, NY; Pfizer; May 2019.
- 171 Diclofenac sodium tablet, delayed release [package insert]. Parsippany, NJ; Actavis; October 2017.
- 172 Voltaren Gel [package insert]. Malvern, PA; Endo; February 2018.
- 173 Diclofenac sodium tablet, film coated, extended release [package insert]. Parsippany, NJ; Actavis; June 2017.
- 174 Arthrotec [package insert]. New York, NY; Pfizer; May 2016.
- 175 Available at: <http://www.micromedexsolutions.com>. Accessed June 8, 2020.
- 176 Zipsor [package insert]. Newark, CA; DepoMed; May 2016.
- 177 Vimovo [package insert]. Deerfield, IL; Horizon; July 2019.
- 178 Sprix [package insert]. Wayne, PA; Egalet; January 2018.
- 179 Duexis [package insert]. Deerfield, IL; Horizon; July 2019. .
- 180 Flector [package insert]. New York, NY; Pfizer; March 2019.
- 181 Licart [package insert]. Parsippany, NJ; IBSA; May 2020.
- 182 Voltaren Gel [package insert]. Malvern, PA; Endo; February 2018.
- 183 Diclofenac sodium solution 1.5% [package insert]. Parsippany, NJ; Actavis; May 2016.
- 184 Pennsaid 2% [package insert]. Lake Forest, IL; Horizon Pharma; May 2016.
- 185 DermacinRx Lexitral PharmaPak [package insert]. San Fernando, CA; PureTek; July 2015.
- 186 Sure Result DSS Premium Pak [package insert]. Doral, FL; International Brand Management; March 2016.
- 187 DS Prep Pak [package insert]. Ocean Springs, MS; Alvix; November 2015.
- 188 Xrylix [package insert].]. San Fernando, CA; PureTek; March 2016.
- 189 Vivlodex [package insert]. Philadelphia, PA; Iroko; October 2015.
- 190 Inflamm-K Kit [package insert]. Peoria, AZ; Solutech; August 2016.
- 191 Diclozor [package insert]. Ripley, MS; Sterling Knight; October 2016.
- 192 Relafen DS [package insert]. Hazlet, NJ; Carwin; January 2020.
- 193 FDA Drug Safety Communication: Possible increased risk of fractures of the hip, wrist, and spine with the use of proton pump inhibitors. Available at: <https://www.fda.gov/Drugs/DrugSafety/PostmarketDrugSafetyInformationforPatientsandProviders/ucm213206.htm>. Accessed June 5, 2020.
- 194 FDA Drug Safety Communication: Low magnesium levels can be associated with long-term use of Proton Pump Inhibitor drugs (PPIs). Available at: <http://www.fda.gov/Drugs/DrugSafety/ucm245011.htm>. Accessed June 5, 2020.
- 195 FDA Drug Safety Communication: Clostridium difficile associated diarrhea can be associated with stomach acid drugs known as proton pump inhibitors (PPIs). Available at: <http://www.fda.gov/Drugs/DrugSafety/ucm290510.htm>. Accessed June 5, 2020.
- 196 COX-2 Selective (includes Bextra, Celebrex, and Vioxx) and Non-Selective Non-Steroidal Anti-Inflammatory Drugs (NSAIDs). Available at: <http://www.fda.gov/drugs/drugsafety/postmarketdrugsafetyinformationforpatientsandproviders/ucm429364.htm>. Accessed June 5, 2020.
- 197 FDA strengthens warning that non-aspirin nonsteroidal anti-inflammatory drugs (NSAIDs) can cause heart attacks or strokes. Available at: <https://www.fda.gov/downloads/Drugs/DrugSafety/UCM453941.pdf>. Accessed June 5, 2020.
- 198 Graham DJ, Campen D, Hui R, et al. Risk of acute myocardial infarction and sudden cardiac death in patients treated with cyclo-oxygenase 2 selective and non-selective non-steroidal anti-inflammatory drugs: nested case-control study. *Lancet*. 2005; 365(9458):475-81.
- 199 Mukherjee D, Nissen SE, Topol EJ. Risk of Cardiovascular events associated with selective COX-2 inhibitors. *JAMA*. 2001; 286:954-959.
- 200 Mamdani M, Juurlink DN, Lee DS, et al. Cyclo-oxygenase-2 inhibitors versus non-selective non-steroidal anti-inflammatory drugs and congestive heart failure outcomes in elderly patients: a population-based cohort study. *Lancet*. 2004; 363(9423):1751-1756.
- 201 Bresalier RS, Sandler RS, Quan H, et al. Cardiovascular events associated with rofecoxib in a colorectal adenoma chemoprevention trial. *N Engl J Med*. 2005; 352(11):1092-102.
- 202 Cardiovascular events associated with rofecoxib in a colorectal adenoma chemoprevention trial. *N Engl J Med*. 2006; 355(2):221. Available at: <http://content.nejm.org/cgi/content/short/NEJMx060029>. Accessed June 5, 2020.
- 203 Nussmeier NA, Whelton AA, Brown MT, et al. Complications of the COX-2 inhibitors parecoxib and valdecoxib after cardiac surgery. *N Engl J Med*. 2005; 352(11):1081-91.
- 204 Aw TJ, Haas SJ, Liew D, et al. Meta-analysis of cyclooxygenase-2 inhibitors and their effects on blood pressure. *Arch Intern Med*. 2005; 165(5):490-6.
-

-
- 205 COX-2 Selective (includes Bextra, Celebrex, and Vioxx) and Non-Selective Non-Steroidal Anti-Inflammatory Drugs (NSAIDs). Available at: www.fda.gov/drugs/drugsafety/postmarketdrugsafetyinformationforpatientsandproviders/ucm429364.htm. Accessed June 5, 2020.
- 206 White WB, Faich G, Borer JS, et al. Cardiovascular thrombotic events in arthritis trials of the cyclooxygenase-2 inhibitor celecoxib. *Am J Cardiol.* 2003; 92(4):411-8.
- 207 White EB, Faich G, Whelton A, et al. Comparison of thromboembolic events in patients treated with celecoxib, a cyclooxygenase-2 specific inhibitor, versus ibuprofen or diclofenac. *Am J Cardiol.* 2002; 89:425-30.
- 208 Bennett JS, Daugherty A, Herrington D, et al. The use of nonsteroidal anti-inflammatory drugs (NSAIDs): a science advisory from the American Heart Association. *Circulation.* 2005; 111:1713–1716.
- 209 Antman EM, Bennett JS, Daugherty A, et al. Use of nonsteroidal anti-inflammatory drugs: an update for clinicians: a Scientific Statement from the American Heart Association. *Circulation.* 2007; 115; 1634-1642. Available at: https://professional.heart.org/professional/GuidelinesStatements/UCM_492626_Guidelines-Statements-Search-Page.jsp. Accessed June 5, 2020.
- 210 FDA strengthens warning that non-aspirin nonsteroidal anti-inflammatory drugs (NSAIDs) can cause heart attacks or strokes. Available at: <https://www.fda.gov/downloads/Drugs/DrugSafety/UCM453941.pdf>. Accessed June 5, 2020.
- 211 Flurbiprofen [package insert]. Morgantown, WV; Mylan; May 2016.
- 212 Zorvolex [package insert]. Philadelphia, PA; Iroko; May 2016.
- 213 Diclofenac potassium [package insert]. Morgantown, WV; Mylan; August 2017.
- 214 Sulindac [package insert]. Parsippany, NJ; Actavis; May 2016.
- 215 Diflunisal [package insert]. Sellersville, PA; Teva; July 2018.
- 216 Flector [package insert]. New York, NY; Pfizer; March 2019.
- 217 Licart [package insert]. Parsippany, NJ; IBSA; May 2020.
- 218 Indocin [package insert]. Philadelphia, PA; Iroko; March 2019.
- 219 Tivorbex [package insert]. Philadelphia, PA; Iroko; March 2019.
- 220 Etodolac [package insert]. Hawthorne, NY; Taro; December 2015.
- 221 Etodolac extended release [package insert]. Hawthorne, NY; Taro; April 2016.
- 222 Mobic [package insert]. Ridgefield, CT; Boehringer-Ingelheim; June 2016.
- 223 Qmiiz ODT [package insert]. Lake Forest, IL; TerSera; October 2018.
- 224 Ibuprofen tablet [package insert]. Bridgewater, NJ; Amneal; May 2019.
- 225 Nalfon [package insert]. Ridgeland, MS; Xspire; May 2016.
- 226 Naprelan [package insert]. San Diego, CA; Victory; July 2019.
- 227 Mefenamic acid [package insert]. Basking Ridge, NJ; Micro; January 2020.
- 228 Anaprox/Anaprox DS [package insert]. Alpharetta, GA; Canton; July 2019.
- 229 Nabumetone [package insert]. Parsippany, NJ; Actavis; May 2016.
- 230 Ketorolac tromethamine tablet [package insert]. Morgantown, WV; Mylan; July 2015.
- 231 Tolmetin [package insert]. Morgantown, WV; Mylan; July 2015.
- 232 Daypro [package insert]. New York, NY; Pfizer; May 2019.
- 233 Feldene [package insert]. New York, NY; Pfizer; May 2019.
- 234 Diclofenac sodium tablet, delayed release [package insert]. Parsippany, NJ; Actavis; October 2017.
- 235 Voltaren Gel [package insert]. Malvern, PA; Endo; February 2018.
- 236 Diclofenac sodium tablet, film coated, extended release [package insert]. Parsippany, NJ; Actavis; June 2017.
- 237 Available at: <http://www.micromedexsolutions.com>. Accessed June 5, 2020.
- 238 Arthrotec [package insert]. New York, NY; Pfizer; May 2016.
- 239 Vimovo [package insert]. Deerfield, IL; Horizon; July 2019.
- 240 Sprix [package insert]. Wayne, PA; Egalet; January 2018.
- 241 Duexis [package insert]. Deerfield, IL; Horizon; July 2019.
- 242 Flector [package insert]. New York, NY; Pfizer; March 2019.
- 243 Voltaren Gel [package insert]. Malvern, PA; Endo; February 2018.
- 244 Diclofenac sodium solution 1.5% [package insert]. Parsippany, NJ; Actavis; May 2016.
- 245 Pennsaid 2% [package insert]. Lake Forest, IL; Horizon; May 2016.
- 246 DermacinRx Lexitral PharmaPak [package insert]. San Fernando, CA; PureTek; July 2015.
- 247 Sure Result DSS Premium Pak [package insert]. Doral, FL; International Brand Management; March 2016.
- 248 DS Prep Pak [package insert]. Ocean Springs, MS; Alvix; November 2015.
- 249 Xrylix [package insert]. San Fernando, CA; PureTek; March 2016.
- 250 Vopac MDS [package insert], Madison, MS; Sircle; April 2016
- 251 Vivlodex [package insert]. Philadelphia, PA; Iroko; October 2015.
- 252 Inflamma-K Kit [package insert]. Peoria, AZ; Solutech; August 2016.
- 253 Diclozor [package insert]. Ripley, MS; Sterling Knight; October 2016.
- 254 Relafen DS [package insert]. Hazlet, NJ; Carwin; January 2020.
- 255 Flurbiprofen [package insert]. Morgantown, WV; Mylan; May 2016.
- 256 Zorvolex [package insert]. Philadelphia, PA; Iroko; May 2016.
- 257 Diclofenac potassium [package insert]. Morgantown, WV; Mylan; August 2017.
- 258 Sulindac [package insert]. Parsippany, NJ; Actavis; May 2016.
- 259 Diflunisal [package insert]. Sellersville, PA; Teva; July 2018.
- 260 Flector [package insert]. New York, NY; Pfizer; March 2019.
- 261 Licart [package insert]. Parsippany, NJ; IBSA; May 2020.
- 262 Indocin [package insert]. Philadelphia, PA; Iroko; March 2019.
- 263 Tivorbex [package insert]. Philadelphia, PA; Iroko; March 2019.
-

264 Etodolac [package insert]. Hawthorne, NY; Taro; December 2015.
265 Etodolac extended release [package insert]. Hawthorne, NY; Taro; April 2016.
266 Mobic [package insert]. Ridgefield, CT; Boehringer-Ingelheim; June 2016.
267 Qmiiz ODT [package insert]. Lake Forest, IL; TerSera; October 2018.
268 Ibuprofen tablet [package insert]. Bridgewater, NJ; Amneal; May 2019.
269 Nalfon [package insert]. Ridgeland, MS; Xspire; May 2016.
270 Naprelan [package insert]. San Diego, CA; Victory; July 2019.
271 Mefenamic acid [package insert]. Basking Ridge, NJ; Micro; January 2020.
272 Anaprox/Anaprox DS [package insert]. Alpharetta, GA; Canton; July 2019.
273 Nabumetone [package insert]. Parsippany, NJ; Actavis; May 2016.
274 Ketorolac tromethamine tablet [package insert]. Morgantown, WV; Mylan; July 2015.
275 Tolmetin [package insert]. Morgantown, WV; Mylan; July 2015.
276 Daypro [package insert]. New York, NY; Pfizer; May 2019.
277 Feldene [package insert]. New York, NY; Pfizer; May 2019.
278 Diclofenac sodium tablet, delayed release [package insert]. Parsippany, NJ; Actavis; October 2017.
279 Voltaren Gel [package insert]. Malvern, PA; Endo; February 2018.
280 Diclofenac sodium tablet, film coated, extended release [package insert]. Parsippany, NJ; Actavis; June 2017.
281 Arthrotec [package insert]. New York, NY; Pfizer; May 2016.
282 Meclofenamate [package insert]. Morgantown, WV; Mylan; July 2015.
283 Celebrex [package insert]. New York, NY; Pfizer; May 2019.
284 Zipsor [package insert]. Newark, CA; DepoMed; May 2016.
285 Vimovo [package insert]. Deerfield, IL; Horizon; July 2019.
286 Duexis [package insert]. Deerfield, IL; Horizon; July 2019.
287 Sprix [package insert]. Wayne, PA; Egalet; January 2018.
288 Flector [package insert]. New York, NY; Pfizer; March 2019.
289 Diclofenac sodium solution 1.5% [package insert]. Parsippany, NJ; Actavis; May 2016.
290 Pennsaid 2% [package insert]. Lake Forest, IL; Horizon; May 2016.
291 Voltaren Gel [package insert]. Malvern, PA; Endo; February 2018.
292 DermacinRx Lexitral PharmaPak [package insert]. San Fernando, CA; PureTek; July 2015.
293 Sure Result DSS Premium Pak [package insert]. Doral, FL; International Brand Management; March 2016.
294 DS Prep Pak [package insert]. Ocean Springs, MS; Alvix; November 2015.
295 Xrylix [package insert]. San Fernando, CA; PureTek; March 2016.
296 Vopac MDS [package insert]. Madison, MS; Sircle; April 2016.
297 Vivlodex [package insert]. Philadelphia, PA; Iroko; October 2015.
298 Inflamm-K Kit [package insert]. Peoria, AZ; Solutech; August 2016.
299 Diclozor [package insert]. Ripley, MS; Sterling Knight; October 2016.
300 Relafen DS [package insert]. Hazlet, NJ; Carwin; January 2020.
301 Flurbiprofen [package insert]. Morgantown, WV; Mylan; May 2016.
302 Zorvolex [package insert]. Philadelphia, PA; Iroko; May 2016.
303 Diclofenac potassium [package insert]. Morgantown, WV; Mylan; August 2017.
304 Sulindac [package insert]. Parsippany, NJ; Actavis; May 2016.
305 Diflunisal [package insert]. Sellersville, PA; Teva; July 2018.
306 Flector [package insert]. New York, NY; Pfizer; March 2019.
307 Licart [package insert]. Parsippany, NJ; IBSA; May 2020.
308 Indocin [package insert]. Philadelphia, PA; Iroko; March 2019.
309 Tivorbex [package insert]. Philadelphia, PA; Iroko; March 2019.
310 Etodolac [package insert]. Hawthorne, NY; Taro; December 2015.
311 Etodolac extended release [package insert]. Hawthorne, NY; Taro; April 2016.
312 Mobic [package insert]. Ridgefield, CT; Boehringer-Ingelheim; June 2016.
313 Qmiiz ODT [package insert]. Lake Forest, IL; TerSera; October 2018.
314 Ibuprofen tablet [package insert]. Bridgewater, NJ; Amneal; May 2019.
315 Nalfon [package insert]. Ridgeland, MS; Xspire; May 2016.
316 Naprelan [package insert]. San Diego, CA; Victory; July 2019.
317 Mefenamic acid [package insert]. Basking Ridge, NJ; Micro; January 2020.
318 Anaprox/Anaprox DS [package insert]. Alpharetta, GA; Canton; July 2019.
319 Nabumetone [package insert]. Parsippany, NJ; Actavis; May 2016.
320 Ketorolac tromethamine tablet [package insert]. Morgantown, WV; Mylan; July 2015.
321 Tolmetin [package insert]. Morgantown, WV; Mylan; July 2015.
322 Daypro [package insert]. New York, NY; Pfizer; May 2019.
323 Feldene [package insert]. New York, NY; Pfizer; May 2019.
324 Diclofenac sodium tablet, delayed release [package insert]. Parsippany, NJ; Actavis; October 2017.
325 Voltaren Gel [package insert]. Malvern, PA; Endo; February 2018.
326 Diclofenac sodium tablet, film coated, extended release [package insert]. Parsippany, NJ; Actavis; June 2017.
327 Arthrotec [package insert]. New York, NY; Pfizer; May 2016.
328 Meclofenamate [package insert]. Morgantown, WV; Mylan; July 2015.
329 Celebrex [package insert]. New York, NY; Pfizer; May 2019.

330 Zipsor [package insert]. Newark, CA; DepoMed; May 2016.
331 Vimovo [package insert]. Deerfield, IL; Horizon; July 2019.
332 Sprix [package insert]. Wayne, PA; Egalet; January 2018.
333 Duexis [package insert]. Deerfield, IL; Horizon; July 2019.
334 Flector [package insert]. New York, NY; Pfizer; March 2019.
335 Voltaren Gel [package insert]. Malvern, PA; Endo; February 2018.
336 Diclofenac sodium solution 1.5% [package insert]. Parsippany, NJ; Actavis; May 2016.
337 Pennsaid 2% [package insert]. Lake Forest, IL; Horizon; May 2016.
338 DermacinRx Lexitral PharmaPak [package insert]. San Fernando, CA; PureTek; July 2015.
339 Sure Result DSS Premium Pak [package insert]. Doral, FL; International Brand Management; March 2016.
340 DS Prep Pak [package insert]. Ocean Springs, MS; Alvix; November 2015.
341 Xrylix [package insert].]. San Fernando, CA; PureTek; March 2016.
342 Vopac MDS [package insert]. Madison, MS; Sircl; April 2016.
343 Inflamma-K Kit [package insert]. Peoria, AZ; Solutech; August 2016.
344 Diclozor [package insert]. Ripley, MS; Sterling Knight; October 2016.
345 Relafen DS [package insert]. Hazlet, NJ; Carwin; January 2020.
346 Flurbiprofen [package insert]. Morgantown, WV; Mylan; May 2016.
347 Zorvolex [package insert]. Philadelphia, PA; Iroko; May 2016.
348 Diclofenac potassium [package insert]. Morgantown, WV; Mylan; August 2017.
349 Sulindac [package insert]. Parsippany, NJ; Actavis; May 2016.
350 Diflunisal [package insert]. Sellersville, PA; Teva; July 2018.
351 Licart [package insert]. Parsippany, NJ; IBSA; May 2020.
352 Flector [package insert]. New York, NY; Pfizer; March 2019.
353 Indocin [package insert]. Philadelphia, PA; Iroko; March 2019.
354 Etodolac [package insert]. Hawthorne, NY; Taro; December 2015.
355 Tivorbex [package insert]. Philadelphia, PA; Iroko; March 2019.
356 Etodolac extended release [package insert]. Hawthorne, NY; Taro; April 2016.
357 Mobic [package insert]. Ridgefield, CT; Boehringer-Ingelheim; June 2016.
358 Qmiiz ODT [package insert]. Lake Forest, IL; TerSera; October 2018.
359 Ibuprofen tablet [package insert]. Bridgewater, NJ; Amneal; May 2019.
360 Nalfon [package insert]. Ridgeland, MS; Xspire; May 2016.
361 Naprelan [package insert]. San Diego, CA; Victory; July 2019.
362 Mefenamic acid [package insert]. Basking Ridge, NJ; Micro; January 2020.
363 Anaprox/Anaprox DS [package insert]. Alpharetta, GA; Canton; July 2019.
364 Nabumetone [package insert]. Parsippany, NJ; Actavis; May 2016.
365 Ketorolac tromethamine tablet [package insert]. Morgantown, WV; Mylan; July 2015.
366 Tolmetin [package insert]. Morgantown, WV; Mylan; July 2015.
367 Daypro [package insert]. New York, NY; Pfizer; May 2019.
368 Feldene [package insert]. New York, NY; Pfizer; May 2019.
369 Diclofenac sodium tablet, delayed release [package insert]. Parsippany, NJ; Actavis; October 2017.
370 Voltaren Gel [package insert]. Malvern, PA; Endo; February 2018.
371 Diclofenac sodium tablet, film coated, extended release [package insert]. Parsippany, NJ; Actavis; June 2017.
372 Arthrotec [package insert]. New York, NY; Pfizer; May 2016.
373 Nudiclo TabPak [package insert]. Nucare; August 2016. Available at: <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=c1f0ff98-c000-40a9-b770-da5813647dd8>. Accessed June 5, 2020.
374 Meclofenamate [package insert]. Morgantown, WV; Mylan; July 2015.
375 Celebrex [package insert]. New York, NY; Pfizer; May 2019.
376 Zipsor [package insert]. Newark, CA; DepoMed; May 2016.
377 Vimovo [package insert]. Deerfield, IL; Horizon; July 2019.
378 Sprix [package insert]. Wayne, PA; Egalet; January 2018.
379 Duexis [package insert]. Deerfield, IL; Horizon; July 2019.
380 Flector [package insert]. New York NY; Pfizer; March 2019.
381 Diclofenac sodium solution 1.5% [package insert]. Parsippany, NJ; Actavis; May 2016.
382 Voltaren Gel [package insert]. Malvern, PA; Endo; February 2018.
383 Pennsaid 2% [package insert]. Lake Forest, IL; Horizon; May 2016.
384 DermacinRx Lexitral PharmaPak [package insert]. San Fernando, CA; PureTek; July 2015.
385 Nudiclo Solupak [package insert]. Nucare; August 2016. Available at: <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=d8105efe-53e9-4392-ac13-7d65fa31211e>. Accessed June 5, 2020.
386 Sure Result DSS Premium Pak [package insert]. Doral, FL; International Brand Management; March 2016.
387 DS Prep Pak [package insert]. Ocean Springs, MS; Alvix; November 2015.
388 Lexixryl [package insert]. Panorama City, CA; Shoreline; February 2017.
389 Xrylix [package insert].]. San Fernando, CA; PureTek; March 2016.
390 Flexipak [package insert]. Ripley, MS; Sterling-Knight; January 2018.
391 Vopac MDS [package insert]. Madison, MS; Sircl; April 2016.
392 Trans-D patch [package insert]. Keego Harbor, MI; Advance Medical Sales; March 2016.
393 Inflamma-K Kit [package insert]. Peoria, AZ; Solutech; August 2016.

- 394 Diclozor [package insert]. Ripley, MS; Sterling Knight; October 2016.
- 395 Comfort Pac with ibuprofen [package insert]. Oklahoma City, OK. PD-RX. October 2017.
- 396 Comfort Pac with meloxicam [package insert]. Oklahoma City, OK. PD-RX. October 2017.
- 397 Comfort Pac with naproxen [package insert]. Oklahoma City, OK. PD-RX. October 2017. Available at: <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=91802fa6-ddf3-4356-85f0-2032cf31478b&audience=consumer>. Accessed June 5, 2020.
- 398 Inflammacin [package insert]. San Fernando, CA; PureTek; April 2019. Available at: <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=91d25301-fac9-41f0-8a4e-e22d44926347&audience=consumer>. Accessed June 5, 2020.
- 399 Diclopak [package insert]. Ripley, MS; Sterling-Knight; January 2018.
- 400 Diclotral Pak [package insert]. Everett, WA; Patchwerx; March 2016. Available at: <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=2d8f4048-14cb-7484-e054-00144ff88e88&audience=consumer>. Accessed June 5, 2020.
- 401 Xelital [package insert] Panorama City, CA; Shoreline; February 2017.
- 402 Xenaflam [package insert]. Panorama City, CA; Shoreline.
- 403 Dithol [package insert]. Ocean Springs, MS; Fortus. December 2017.
- 404 Trixyltral. Available at: <https://www.drugs.com/pro/trixyltral.html>. Accessed June 5, 2020.
- 405 DicloPR [package insert]. Tampa, FL; 19&Pacific; November 2017.
- 406 NuDroxiPak Pain Relief Pak [package insert]. NuCare Pharmaceuticals. February 2018. Available at: <https://dailymed.nlm.nih.gov/dailymed/search.cfm?labeltype=all&query=nudroxipak&pagesize=200&page=1&audience=consumer>. Accessed June 5, 2020.
- 407 NuDroxiPak DSDR-75 [package insert]. NuCare Pharmaceuticals. February 2018. Available at: <https://dailymed.nlm.nih.gov/dailymed/search.cfm?labeltype=all&query=nudroxipak&pagesize=200&page=1&audience=consumer>. Accessed June 5, 2020.
- 408 CapsFenac Pak [package insert]. Los Angeles, CA; SA3; December 2018. Available at: <https://www.dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=88728f7a-19e5-4f31-911d-fc7d8a64ba0d>. Accessed June 5, 2020.
- 409 Diclofex DC [package insert]. Madison, MS; Sirlec; January 2019. Available at: <https://www.dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=d8ba9e7c-3e5f-4c41-add9-3e1a142c1fff>. Accessed June 5, 2020.
- 410 Diclovix [package insert]. Ocean Springs, MS; Primary; March 2019. Available at: <https://www.dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=84b45252-fdf6-0ab0-e053-2991aa0ac428>. Accessed June 5, 2020.
- 411 Inavix [package insert]. Ocean Springs, MS; Primary; July 2020.
- 412 Capsinac [package insert]. Las Vegas, NV; Bodysphere; July 2015.
- 413 Diclotrex [package insert]. San Fernando, CA; Purtek; October 2018.
- 414 Diclovix M [package insert]. Ocean Springs, MS; Primary; January 2020. Available at: <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=9d0012ed-a50a-a936-e053-2995a90a6100>. Accessed June 8, 2020.
- 415 Relafen DS [package insert]. Hazlet, NJ; Carwin; January 2020.
- 416 Inflattherm (diclofenac/menthol/camphor) [package insert]. Portland, OR. V2 Pharma. July 2019.
- 417 Inflattherm (diclofenac/trolamine salicylate) [package insert]. Portland, OR. V2 Pharma. July 2019.
- 418 Diclosiacin [package insert]. Madison, MS; Chadwick; July 2018.
- 419 Dimenth [package insert]. Biloxi, MS; Levins; October 2019.
- 420 Varophen [package insert]. Portland, OR. V2 Pharma. June 2019.
- 421 Available at: <https://www.drugs.com/ppa/ketoprofen.html>. Accessed June 5, 2020.
- 422 Valat JP, Accardo S, Reginster JY, et al. A comparison of the efficacy and tolerability of meloxicam and diclofenac in the treatment of patients with osteoarthritis of the lumbar spine. *Inflamm Res*. 2001; 50(Suppl 1):S30–4.
- 423 Linden B, Distel M, Bluhmki E. A double-blind study to compare the efficacy and safety of meloxicam 15 mg with piroxicam 20 mg in patients with osteoarthritis of the hip. *British Journal of Rheumatology*. 1996; 35(Suppl 1):35–8.
- 424 Wojtulewski JA, Schattenkirchner M, Barcelo P, et al. A six-month double-blind trial to compare the efficacy and safety of meloxicam 7.5 mg daily and naproxen 750 mg daily in patients with rheumatoid arthritis. *British Journal of Rheumatology*. 1996; 35(Suppl 1):22–8.
- 425 Rogind H, Bliddal H, Klokke D, et al. Comparison of etodolac and piroxicam in patients with osteoarthritis of the hip or knee: A prospective, randomised, double-blind, controlled multicentre study. *Clinical Drug Investigation*. 1997; 13(2):66–75.
- 426 Liang TH, Hsu PN. Double-blind, randomised, comparative trial of etodolac SR versus diclofenac in the treatment of osteoarthritis of the knee. *Curr Med Res Opin*. 2003; 19(4):336–41.
- 427 Chan FK, Hung LC, Suen BY, et al. Celecoxib versus diclofenac and omeprazole in reducing the risk of recurrent ulcer bleeding in patients with arthritis. *N Engl J Med*. 2002; 347(26):2104–2110.
- 428 Chan FK, Hung LC, Suen BY, et al. Celecoxib versus diclofenac plus omeprazole in high-risk arthritis patients: results of a randomized double-blind trial. *Gastroenterology*. 2004; 127(4):1038–1043.
- 429 Silverstein FE, Faich G, Goldstein JL, et al. Gastrointestinal toxicity with celecoxib vs. nonsteroidal anti-inflammatory drugs for osteoarthritis and rheumatoid arthritis. The CLASS Study: a randomized controlled trial. *JAMA*. 2000; 284:1247–1255.
- 430 Dougados M, Behier JM, Jolchine I, et al. Efficacy of celecoxib, a cyclooxygenase 2-specific inhibitor, in the treatment of ankylosing spondylitis: a six-week controlled study with comparison against placebo and against a conventional nonsteroidal anti-inflammatory drug. *Arthritis Rheum*. 2001; 44(1):180–185.
- 431 Kivitz AJ, Moskowitz RW, Woods E, et al. Comparative efficacy and safety of celecoxib and naproxen in the treatment of osteoarthritis of the hip. *J Int Med Res*. 2001; 29(6):467–479.
- 432 Bensen WG, Fiechtner JJ, McMillen JJ, et al. Treatment of osteoarthritis with celecoxib, a cyclooxygenase-2 inhibitor: a randomized controlled trial. *Mayo Clin Proc*. 1999; 74(11):1095–1105.
- 433 Goldstein JL, Correa P, Zhao WW, et al. Reduced incidence of gastroduodenal ulcers with celecoxib, a novel cyclooxygenase-2 inhibitor, compared to naproxen in patients with arthritis. *Am J Gastroenterol*. 2001; 96(4):1019–1027.
- 434 Singh G, Fort JG, Goldstein JL, et al. Celecoxib versus naproxen and diclofenac in osteoarthritis patients: SUCCESS-I Study. *Am J Med*. 2006; 119(3):255–266.

- 435 Gibofsky A, Silberstein S, Argoff C, et al. Lower-dose diclofenac submicron particle capsules provide early and sustained acute pain patient relief in a phase 3 study. *Postgrad Med*. 2013; 125(5):130-8. DOI: 10.3810/pgm.2013.09.2693.
- 436 Petri M, Hufman SL, Waser G, et al. Celecoxib effectively treats patients with acute shoulder tendinitis/bursitis. *J Rheumatol*. 2004; 31(8):1614-1620.
- 437 Petrella R, Ekman EF, Schuller R, et al. Efficacy of celecoxib, a COX-2-specific inhibitor, and naproxen in the management of acute ankle sprain: results of a double-blind, randomized controlled trial. *Clin J Sport Med*. 2004; 14(4):225-231.
- 438 Salo DF, Lavery R, Varma V, et al. A randomized, clinical trial comparing oral celecoxib 200 mg, celecoxib 400 mg, and ibuprofen 600 mg for acute pain. *Acad Emerg Med*. 2003; 10(1):22-30.
- 439 Ekman EF, Fiechtner JJ, Levy S, et al. Efficacy of celecoxib versus ibuprofen in the treatment of acute pain: a multicenter, double-blind, randomized controlled trial in acute ankle sprain. *Am J Orthop*. 2002; 31(8):445-451.
- 440 Nikanne E, Kokki H, Salo J, et al. Celecoxib and ketoprofen for pain management during tonsillectomy: a placebo-controlled clinical trial. *Otolaryngol Head Neck Surg*. 2005; 132(2):287-294.
- 441 Nissen Se, Yeomans ND, Solomon DH, et al. Cardiovascular safety of celecoxib, naproxen, or ibuprofen for arthritis. *N Engl J Med*. 2016; 375 (26): 2519-2529. DOI: 10.1056/NEJMoa1611593.
- 442 Bicameral TS, Weaver AL, Tindal EA, et al. Diclofenac/misoprostol compared with diclofenac in the treatment of osteoarthritis of the knee or hip: a randomized, placebo controlled trial. *Arthrosc Osteoarthritis Study Group. J Rheumatol*. 1998; 25(8):1602-11.
- 443 Bolten W, Gomes JA, Stead H, et al. The gastro duodenal safety and efficacy of the fixed combination of diclofenac and misoprostol in the treatment of osteoarthritis. *Br J Rheumatol*. 1992; 31(11):753-758.
- 444 de Queiroz MV, Beaulieu A, Kruger K, et al. Double-blind comparison of the efficacy of diclofenac/misoprostol and diclofenac in the treatment of rheumatoid arthritis. *Euro J Rheumatol Inflamm*. 1994; 14(2):5-13.
- 445 Agawam NM, Caldwell J, Kibitz AJ, et al. Comparison of the upper gastrointestinal safety of Arthrotec 75 and nabumetone in osteoarthritis patients at high risk for developing nonsteroidal anti-inflammatory drug-induced gastrointestinal ulcers. *Clin Ther*. 1999; 21(4):659-74.
- 446 Gibofsky A, Hochberg MC, Jaros MJ, et al. Efficacy and safety of low-dose submicron diclofenac for the treatment of osteoarthritis pain: a 12 week, phase 3 study. *Curr Med Res Opin*. 2014; 30(9):1883-1893. DOI: 10.1185/03007995.2014.946123.
- 447 Duexis [package insert]. Deerfield, IL; Horizon; July 2019.
- 448 Tivorbex [package insert]. Philadelphia, PA; Iroko; March 2019.
- 449 Vivlodex [package insert]. Philadelphia, PA; Iroko; October 2015.
- 450 Altman R, Hochberg M, Gibofsky A, et al. Efficacy and safety of low-dose SoluMatrix meloxicam in the treatment of osteoarthritis pain: a 12-week, phase 3 study. *Curr Med Res Opin*. 2015; 31(12): 2331-2343. DOI: 10.1185/03007995.2015.1112772.
- 451 Vimovo [package insert]. Deerfield, IL; Horizon Pharma USA; July 2017.
- 452 Goldstein J, Hochberg MC, Fort JG, et al. Clinical trial: the incidence of NSAID-associated endoscopic gastric ulcers in patients treated with PN 400 (naproxen plus esomeprazole magnesium) vs. enteric-coated naproxen alone. *Aliment Pharmacol Ther*. 2010; 32:401-413.
- 453 Vimovo [package insert]. Deerfield, IL; Horizon Pharma USA; July 2017.
- 454 Graham DY, Agrawal NM, Campbell DR, et al. Ulcer prevention in long-term users of nonsteroidal anti-inflammatory drugs: results of a double-blind, randomized, multicenter, active- and placebo-controlled study of misoprostol vs. lansoprazole. *Arch Intern Med*. 2002; 162(2):169-175.
- 455 Cullen D, Bardhan KD, Eisner M, et al. Primary gastroduodenal prophylaxis with omeprazole for non-steroidal anti-inflammatory drug users. *Aliment Pharmacol Ther*. 1998; 12(2):135-140.
- 456 Hawkey CJ, Karrasch JA, Szczepanski L, et al. Omeprazole compared with misoprostol for ulcers associated with nonsteroidal antiinflammatory drugs. Omeprazole versus Misoprostol for NSAID-induced Ulcer Management (OMNIUM) Study Group. *N Engl J Med*. 1998; 338(11):727-734.
- 457 Sprix [package insert]. Wayne, PA; Egalet; January 2018.
- 458 Sprix [package insert]. Wayne, PA; Egalet; January 2018.
- 459 Predel H, Koll R, Pabst H, et al. Diclofenac patch for topical treatment of acute impact injuries: a randomized, double-blind, placebo-controlled, multicenter study. *Br J Sports Med*. 2004; 38:318-323.
- 460 Galer BS, Rowbatham M, Perander J, et al. Topical diclofenac patch relieves minor sports injury pain: results of a multicenter controlled clinical trial. *J Pain Symptom Manage*. 2000; 19(4):287-294.
- 461 Bookman AA, Williams KS, Shainhouse JZ. Effect of a topical diclofenac solution for relieving symptoms of primary osteoarthritis of the knee: a randomized controlled trial. *CMAJ*. 2004; 171(4):333-338.
- 462 Baer PA, Thomas LM, Shainhouse Z. Treatment of osteoarthritis of the knee with a topical diclofenac solution: a randomised controlled, 6-week trial. *BMC Musculoskelet Disord*. 2005; 6:44.
- 463 Roth SH, Shainhouse JZ. Efficacy and safety of a topical diclofenac solution (Pennsaid) in the treatment of primary osteoarthritis of the knee: a randomized, double-blind, vehicle-controlled clinical trial. *Arch Intern Med*. 2004; 164(18):2017-2023.
- 464 Simon LS, Grierson LM, Naseer Z, et al. Efficacy and safety of topical diclofenac containing dimethyl sulfoxide (DMSO) compared with those of topical placebo, DMSO vehicle and oral diclofenac for knee osteoarthritis. *Pain*. 2009; 143(3):238-245.
- 465 Altman RD, Dreiser RL, Fisher CL, et al. Diclofenac sodium gel in patients with primary hand osteoarthritis: a randomized, double-blind, placebo-controlled trial. *J Rheumatol*. 2009; 36(9):1991-1999.
- 466 Barthel HR, Haselwood D, Longley S 3rd, et al. Randomized controlled trial of diclofenac sodium gel in knee osteoarthritis. *Semin Arthritis Rheum*. 2009; 39(3):203-212.