

Technical Notes Hospital Quality-Based Program: Potentially Preventable Readmissions

Patient Population: Texas Medicaid and CHIP Reporting Period: State Fiscal Year 2020 Midyear

The Institute for Child Health Policy University of Florida

The External Quality Review Organization for Texas Medicaid Managed Care and CHIP

Issue Date: July 22, 2020

Table of Contents

Section 1. Introduction1
Section 2. Data
Section 3. PPR Logic and Calculation for Facilities2
Section 4. How Hospitals Should Use This Hospital Level Report4
Section 5. Guide to the PPR Hospital Level Report4
Hospital4
National Provider Identifier (NPI)4
Texas Provider Identifier (TPI)4
Table 1: Hospital PPR Rates 5
Total Admissions at Risk for PPR5
Actual Number of PPR Chains5
Actual PPR Weights5
Expected PPR Weights
Actual to Expected Ratio for PPR Weights5
Total Reimbursement Reduction5
Table 2: Hospital PPR Expenditures 6
Members with PPRs6
Number of PPR Events
Actual PPR Expenditures
Table 3: State-Wide Hospital PPR Rate (information can be found in the State Norm file on the HHSC PPE webpage, see References)
Percentiles6
Table 4: State-Wide Hospital Distributions (information can be found in the State Norm file on the HHSC PPE webpage, see References)
Percentiles6
Table 5: Hospital PPR Results by PPR Reason6
Table 6: Hospital PPR Results by APR-DRG 6
Section 6. Underlying Admissions Data8
References

Section 1. Introduction

Readmissions have potential value as an indicator of quality of care because they may reflect poor clinical care and poor coordination of services either during hospitalization or in the immediate post discharge period. A potentially preventable readmission (PPR) is a readmission (return hospitalization within the specified readmission time interval) that is clinically related to the initial hospital admission. "Clinically related" is defined as a requirement that the underlying reason for readmission be plausibly related to the care rendered during or immediately following a prior hospital admission. A readmission within a specified time interval, called the readmission time interval. The readmission time interval is the maximum number of days allowed between the discharge date of a prior admission and the admitting date of a subsequent admission. If a subsequent admission occurs within the readmission time interval and is clinically related to a prior admission, it is considered a PPR. The hospitalization triggering a PPR is called an Initial Admission. Subsequent PPRs relate back to the care rendered during or following the Initial Admission.

Using the 3M approach, PPRs are also measured in other quality initiatives in Texas, namely the Delivery System Reform Incentive Payment (DSRIP program, website: <u>https://hhs.texas.gov/laws-</u> <u>regulations/policies-rules/waivers/medicaid-1115-waiver</u>) and in the future, the Department of State Health Services (DSHS) hospital quality reporting process (website: <u>https://www.dshs.state.tx.us/thcic/default.shtm</u>).

For this report a 15 day readmission window was applied.

Section 2. Data

Inpatient facility admissions for all Medicaid and CHIP population (MCO encounters and FFS claims) with an admission date for state fiscal year 2020 midyear (September 1, 2019 through February 29, 2020) are included with three exceptions:

- **Undocumented Alien Status** If the patient had undocumented alien status and the client was discharged and readmitted, the readmission may not have been captured in the Medicaid database.
- *Medicaid / Medicare Dual Eligibility* Patients who are dually eligible for both Medicaid and Medicare during the measurement year were excluded.
- Global PPR exclusions Admissions with certain major or metastatic malignancy conditions; a
 situation where it is very likely that a readmission was either planned, unpreventable, or beyond a
 hospital's influence; and encounters with a discharge status of "left against medical advice". Refer to
 the 3M specifications manual for the complete list of exclusions.

Section 3. PPR Logic and Calculation for Facilities

The 3M[™] PPR methodology is a computerized algorithm to identify readmissions with a plausible clinical relationship to the care rendered during or immediately following a prior hospital admission. Every stay is assigned to an All Patient Refined (APR)-Diagnosis Related Group (DRG). There are more than three hundred base APR-DRGs and each base APR-DRG has four levels of severity. Severity of illness (SOI) is defined as the extent of organ system derangement or physiologic decompensation for a patient. It gives a medical classification into 1=minor, 2=moderate, 3=major, and 4=extreme.

When a patient was admitted to acute inpatient hospital more than once during the measurement period, whether a subsequent admission would be classified as a potentially preventable admission (PPR) was determined by readmission interval and clinical relationship. Only when the subsequent admission occurred within the pre-defined readmission interval (15 days) after discharge from the initial admission would the readmission be further considered for preventability. If the qualified readmission (admitted within 15 days after previous discharge) was clinically related to the initial admission as determined by their APR-DRGs, the readmission would be considered a PPR. The initial admission and the PPR formed a PPR chain. A PPR chain could contain more than one PPR. In the case when two or more readmissions followed the initial admission, and the intervals between each other were within 15 days, then the clinical relationship of each readmission would be assessed based on the relationship between each adjacent readmission pair.

A 3M panel of clinicians determined whether each possible admission/readmission pair represented a PPR. For some pairs, additional factors were considered, including patient age or particular diagnoses and procedures within an APR-DRG. A readmission is considered clinically-related to the Initial Admission if the reason for the readmission falls into one of three categories for medical readmissions, one of two categories for surgical readmissions and one category for mental health or substance abuse conditions. Readmissions for medical reasons are much more common than readmissions for surgical procedures, regardless of the reason for the Initial Admission. The three categories of clinically-related medical readmissions are as follows:

- A continuation or recurrence of the reason for the Initial admission, or for a condition closely related to the reason for the Initial Admission (e.g. a readmission for diabetes following an Initial Admission for diabetes).
- An ambulatory care sensitive condition (as designated by the Agency for Healthcare Research and Quality, AHRQ), or a chronic problem that may be related to care either during or after the initial admission.
- An acute medical condition or complication that may be related to or have resulted from care during or after the initial admission.

Surgical readmissions were generally considered not preventable unless they met one of the two criteria for a clinical relationship to the Initial Admission:

- A readmission for a surgical procedure that addressed a continuation or a recurrence of the problem causing the Initial Admission (a patient readmitted for an appendectomy following an Initial Admission for abdominal pain and fever).
- A readmission for a surgical procedure that addressed a complication resulting from care during the Initial Admission (a readmission for drainage of a post-operative wound abscess following an Initial Admission for a bowel resection).

Readmissions for mental health or substance abuse reasons will be considered clinically related regardless of the diagnoses for the initial admission.

- Readmission for mental health reasons following an initial admission for a non-mental health, nonsubstance abuse reason.
- Readmission for a substance abuse diagnosis reason following an initial admission for a non-mental health, non-substance abuse reason.
- Mental health or substance abuse readmission following an initial admission for a substance abuse or mental health diagnosis.

A readmission that did not fit one of these categories (e.g., a readmission for trauma) is classified as a clinically-unrelated readmission and, therefore, not potentially preventable (i.e. not a PPR). A readmission that is not clinically-related to the Initial Admission in a readmission chain terminates the readmission chain. A readmission that has a discharge status of transferred to an acute care hospital, left against medical advice or died terminates a readmission chain.

Relative weights are assigned to each admission by APR-DRG. These weights (Texas relative weights for APR-DRG Grouper 37, effective 10/01/2019) were determined based on resource utilization from Texas Medicaid data, and can be downloaded from

<u>http://www.tmhp.com/Pages/Medicaid/Medicaid_hosp_reimb.aspx</u>. Weighting the PPRs is important because not all PPRs have the same resource costs. High resource PPR should weigh more when comparing the impact of PPRs.

In the PPR reports prior to SFY2014, all PPR chains were event-based. No relative weights were assigned to the PPR chains. There are potential drawbacks with the unweighted approach, because 1) not all PPRs have the same resource costs, 2) more than one PPRs can happen in a single PPR chain. Starting SFY2014, PPR chains will be weighted and the Actual to Expected (A/E) ratio will also be based on the PPR weights. In SFY2020 midyear, the hospital-level reports were revised to also include information on event-based/unweighted PPRs to provide hospitals with a better view of their performance.

Using all the qualifying encounters data from the state, state norm PPR weight is calculated for each APR-DRG and SOI (severity of illness) level of the initial admissions. The initial admissions are first grouped by their APR-DRGs and SOI level, then APR-DRG weights of the PPRs following the initial admissions were summed up respectively for different APR-DRG and SOI level of the initial admission. For each APR-DRG and SOI level within each hospital, the expected PPR weight was calculated by multiplying the state norm PPR weight for each APR-DRG and SOI level by the number of candidate

admissions in the hospital in the corresponding APR-DRG and SOI level. The overall expected PPR weights for the hospital is the sum of the expected PPR weight for all APR-DRGs and SOI levels.

Section 4. How Hospitals Should Use This Hospital Level Report

The "Hospital PPR Results by APR-DRG" table in the report lists the top 40 APR-DRGs of the initial admissions, ranked by the total counts of the PPR events following the initial admission. Associated PPR weights following the same initial admission will also be provided in the same table. Hospitals may focus on the top APR-DRGs to target opportunities for improvement. The underlying detail data will be provided to hospitals upon request. The underlying data is a good resource for hospitals to identity which claims/encounters are potentially preventable. Based on the information provided, hospitals can design their own intervention strategies.

The underlying data used to generate the hospital level report can be requested via an email to <u>MCD_PPR_PPC@hhsc.state.tx.us</u> (please provide full name, email, phone number, NPI, TPI and hospital name).

Section 5. Guide to the PPR Hospital Level Report

Using the 3M[™] PPR Grouping software and methodology (3M Core Grouping Software Version 2020.1.4, PPR Version 37.0), encounter and eligibility data for Texas Medicaid for SFY2020 midyear was used to calculate facility rates for PPRs.

Low volume hospitals can affect the reliability and interpretability of hospital-based summary statistics such as statewide percentile rankings. Hospitals meeting any of the following criteria below were considered low volume. These hospitals will receive a report, but will be excluded from reimbursement reductions.

- Less than 40 total admissions at risk for PPR or,
- Less than 5 actual PPR chains or,
- Less than 5 expected PPR chains

Hospital

The hospital name associated with the NPI.

National Provider Identifier (NPI)

The NPI associated with the hospital, and identified as the billing hospital in the encounters attributed to the hospital and included in the hospital results.

Texas Provider Identifier (TPI)

The TPIs corresponding to the hospital NPI based on the crosswalk provided by Texas Medicaid Healthcare Partnership (TMHP).

Table 1: Hospital PPR Rates

Total Admissions at Risk for PPR

All encounters with bill type = '11x', '12x', '41x' (hospital inpatient encounters) minus the global exclusions and non-events per the software. For the fiscal year 2020 midyear reporting period, admissions between September 1, 2019 and February 14, 2020 were evaluated. The last 15 days of the reporting period are excluded to allow for the 15 days readmission window. The total admissions at risk for PPR can be found in the underlying data by filtering the 'Candidate Admissions' column to only show = 1 (yes).

Actual Number of PPR Chains

A readmission chain is a sequence of PPRs that are all clinically related to the Initial Admission. A readmission chain may contain an Initial Admission and only one PPR, which is the most common situation, or may contain multiple PPRs following the Initial Admission. The actual number of PPR chains can be found in the underlying data by filtering the 'Initial Admissions' column to only show = 1 (yes).

Actual PPR Weights

The sum of APR-DRG weights (Texas relative weights for APR-DRG Grouper 37, effective 10/01/2019) for all PPRs. The hospital level report only lists the top 40 initial admissions by APR-DRG. To calculate the actual PPR weights you will need the underlying data that lists all APR-DRGs. For calculations, see the section on "Hospital PPR Results by APR-DRG." APR-DRG weights can be found on the TMHP website at: http://www.tmhp.com/Pages/Medicaid/Medicaid_hosp_reimb.aspx

Expected PPR Weights

Expected PPR weights reflects the sum of APR-DRG weights that would be expected based on your hospital's case mix, calculated based on APR-DRG (including severity of illness), the presence or absence of a major mental health or substance abuse comorbidity, and patient age.

Actual to Expected Ratio for PPR Weights

The ratio of the actual PPR weights to the expected PPR weights indicates the performance of the hospital relative to the overall performance for all hospitals in Texas. This is calculated by dividing the Actual PPR Weights by the Expected PPR Weights.

Total Reimbursement Reduction

For the midyear report there is no reimbursement reduction.

Actual PPR Rate

Actual number of PPR chains divided by total admissions at risk for PPR.

Expected PPR Rate

Expected PPR rate is obtained by dividing the expected number of PPR chains by the total admissions at risk of PPR.

Table 2: Hospital PPR Expenditures

Members with PPRs

Number of unique clients with PPRs. The unique members with PPRs can be found in the underlying data by filtering for the 'Initial Admission' column = 1 (yes). Then de-duplicate the 'Medicaid Client ID' to find the unique members with PPRs.

Number of PPR Events

Count of all Readmissions. The number of PPR events can be found in the underlying data by filtering 'Potentially Preventable Readmission (PPR)' column to only show = 1 (yes).

Actual PPR Expenditures

Sum of payment amounts associated with all PPRs.

Table 3: State-Wide Hospital PPR Rate (information can be found in the State Norm file on the HHSC PPE webpage, see References)

Percentiles

Calculated from 'Actual PPR Weights' for all hospitals, excluding those identified as low volume hospitals.

Table 4: State-Wide Hospital Distributions (information can be found in the StateNorm file on the HHSC PPE webpage, see References)

Percentiles

Calculated from 'Total Admissions at Risk for PPR', 'Actual Number of PPR Chains, 'Members with PPRs', and 'Number of PPR Events' for all hospitals, excluding those identified as low volume hospitals.

Table 5: Hospital PPR Results by PPR Reason

This table provides a distribution of Number of PPR Events, PPR weights and expenditures for each PPR Reason for the hospital based on the clinical-related reasons defined by 3M. PPR Reasons are identified by APR-DRG.

Table 6: Hospital PPR Results by APR-DRG

This table lists the top 40 APR-DRGs of the initial admission, not the PPR admission. The table is shown this way because the initial admission is crucial in the PPR analysis. The table is ranked by total number of PPR events following these initial admissions. The table provides hospitals an entry point to explore the possibility to maximize the inpatient care quality by studying the subset of diagnosis groups that had

PPR Hospital Reporting Technical Notes

cause the most resource utilization due to PPR. PPRs often result from defective care in initial admissions. Hospitals are recommended to consult clinicians in order to improve the performance.

This table consists of seven columns:

- 1) Initial Admission APR DRG The APR DRG in this table is listed as the overall APR DRG. APR DRGs are further broken down by severity of illness. This is indicated as 1 through 4 (from low to high severity of illness) after the overall APR DRG. For example, APR DRG 254 | Other Digestive System Diagnosis has four levels of severity indicated as 254-1, 254-2, 254-3, and 254-4 (this is indicated in the underlying data in the 'APR DRG' column). It is important to note that the higher the severity of illness, the higher the PPR weight.
- 2) PPR Count This is a multi-step process in calculating the PPR Count of each overall APR DRG by using the underlying data. Begin by creating a new table in a separate spreadsheet or paper with three columns that you will need to fill out for Unique PPR Chain, APR DRG for PPRs, and APR DRG Weights for PPRs. Follow steps a. through d.
 - a. In the underlying data, filter the records which are not considered initial admission or a PPR by filtering 'Unique Chain ID' column to show all chains except F0.
 - b. Then filter the 'Initial Admission' column to only show = 1 (yes) and filter the 'APR DRG' column to only show the corresponding APR DRG (include all severities of illness). Write down the 'Unique Chain IDs' in the new table mentioned in 2) under Unique PPR Chain.
 - c. Un-filter all columns and identify chains by filtering the 'Unique Chain ID' column to show only the unique chain IDs that you wrote down in the new table for b. Then identify the PPRs by filtering the 'Potentially Preventable Readmission (PPR)' column to only show = 1 (yes). Write down the PPR weight for each unique chain in the new table mentioned in 2) under the corresponding APR DRG for all PPRs.
 - d. Sum up all PPRs to arrive as the PPR count for that particular APR DRG.
- 3) Fraction of Total PPR Events -- This is calculated by dividing the 'Number of PPR Events' for each overall APR DRG by the 'Number of PPR Events' Column in the second table of the hospital report, and multiplying by 100.
- 4) PPR Weights Use the table generated above, and sum up all PPR weights for all PPRs to arrive as the PPR weights for that particular APR DRG.
- 5) Fraction of Total PPR Weights This is calculated by dividing the 'PPR Weights' for each overall APR DRG by the 'Actual PPR Weights' (found in the first table of the hospital report), and multiplying by 100.

- 6) PPR Expenditures Medicaid payment amount associated with each encounter is considered confidential, and is not shown in the underlying data. This number can be found by matching the 'Medicaid Claim ID' column in the underlying data to the hospital's claims data.
- 7) Fraction of PPR Expenditures This is calculated by dividing the 'PPR Expenditures' for each overall APR DRG by the 'Actual PPR Expenditures' (found in the second table of the hospital report), and multiplying by 100.

Section 6. Underlying Admissions Data

The underlying inpatient admissions data are supplemented with the report cards for the providers to reproduce the PPR results. The data sets contain patients' Medicaid client IDs, claim IDs, initial admission and readmission hospital identifiers, discharge APR-DRG and SOI, discharge status, mental health comorbidity flag, unique PPR chain ID, clinical relationship reason code, primary and secondary diagnosis codes, and indicators for candidate admission, initial admission, and potentially preventable readmission.

The Medicaid client ID and claim ID allow providers to match the admission data with their own database to obtain additional information. The indicators for candidate admission, initial admission, and potentially preventable readmission allow the providers to reproduce the PPR results. A total PPR count is the sum of the "1"s in the column "Potential Preventable Readmission". PPR counts were multiplied by the corresponding APR-DRG weights to calculate the total PPR weights.

References

3M[™] Health Information Systems. Potentially Preventable Readmissions Classification System: Methodology Overview. GRP-118, 2020.

3M[™] Potentially Preventable Readmissions (PPRs) v37.0: Definitions Manual.

3M[™] Definitions Manuals via <u>www.aprdrgassign.com</u> (Username: TXHosp; Password: aprdrg004)

HHS PPE Webpage: <u>http://hhs.texas.gov/about-hhs/process-improvement/medicaid-chip-quality-</u> efficiency-improvement/potentially-preventable-events

Contact email for questions: MCD_PPR_PPC@hhsc.state.tx.us