



Colony Stimulating Factors Therapeutic Class Review (TCR)

February 6, 2023

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, digital scanning, or via any information storage or retrieval system without the express written consent of Magellan Rx Management.

All requests for permission should be mailed to:

Magellan Rx Management
Attention: Legal Department
6950 Columbia Gateway Drive
Columbia, Maryland 21046

The materials contained herein represent the opinions of the collective authors and editors and should not be construed to be the official representation of any professional organization or group, any state Pharmacy and Therapeutics committee, any state Medicaid Agency, or any other clinical committee. This material is not intended to be relied upon as medical advice for specific medical cases and nothing contained herein should be relied upon by any patient, medical professional or layperson seeking information about a specific course of treatment for a specific medical condition. All readers of this material are responsible for independently obtaining medical advice and guidance from their own physician and/or other medical professional in regard to the best course of treatment for their specific medical condition. This publication, inclusive of all forms contained herein, is intended to be educational in nature and is intended to be used for informational purposes only. Send comments and suggestions to PSTCEditor@magellanhealth.com.

FDA-APPROVED INDICATIONS

Drug	Manufacturer	Cancer patients receiving myelosuppressive chemotherapy (To reduce incidence of infection [febrile neutropenia])	Acute Myeloid Leukemia (AML) patients receiving chemotherapy (Following induction or consolidation chemotherapy to reduce time to neutrophil recovery and the duration of fever in adults)	Bone Marrow Transplant (BMT)	Peripheral Blood Progenitor Cell Collection and Therapy	Severe Chronic Neutropenia (To reduce the incidence and duration of neutropenia in symptomatic patients with congenital neutropenia, cyclic neutropenia, or idiopathic neutropenia)	Hematopoietic Syndrome of Acute Radiation Syndrome (To Increase survival in patients acutely exposed to myelosuppressive doses of radiation)
eflapegrastim-xnst (Rolvedon™) ¹	Spectrum	X	--	--	--	--	--
filgrastim (Neupogen®) ²	Amgen	X	X	X ^a	X	X	X
filgrastim-aafi (Nivestym®) ^{*3}	Pfizer	X	X	X ^a	X	X	--
filgrastim-ayow (Releuko®) ^{*4}	Amneal	X	X	X ^a	--	X	--
filgrastim-sndz (Zarxio®) ^{*5}	Sandoz	X	X	X ^a	X	X	--
pegfilgrastim (Neulasta®) ⁶	Amgen	X	--	--	--	--	X
pegfilgrastim-apgf (Nyvepria™) ^{†7}	Pfizer	X	--	--	--	--	--
pegfilgrastim-cbqv (Udenyca®) ^{†8}	Coherus	X	--	--	--	--	X
pegfilgrastim-jmdb (Fulphila®) ^{†9}	Mylan	X	--	--	--	--	--
pegfilgrastim-bmez (Ziextenzo®) ^{†10}	Sandoz	X	--	--	--	--	--
pegfilgrastim-fpgk (Stimufend®) ^{†11}	Fresenius Kabi	X	--	--	--	--	--

* Filgrastim-aafi (Nivestym), filgrastim-ayow (Releuko), and filgrastim-sndz (Zarxio) are biosimilars to filgrastim (Neupogen) and were approved via the FDA 351(k) biosimilar pathway.

† Pegfilgrastim-apgf (Nyvepria), pegfilgrastim-cbqv (Udenyca), pegfilgrastim-jmdb (Fulphila), pegfilgrastim-bmez (Ziextenzo), pegfilgrastim-fpgk (Stimufend), and pegfilgrastim-pbbk (Fylnetra) are biosimilars to the originator pegfilgrastim (Neulasta), approved through the FDA 351(k) biosimilar pathway.

^a In cancer patients receiving BMT to reduce duration of neutropenia and febrile neutropenia

FDA Approved Indication (continued)

Drug	Manufacturer	Cancer patients receiving myelosuppressive chemotherapy (To reduce incidence of infection [febrile neutropenia])	Acute Myeloid Leukemia (AML) patients receiving chemotherapy (Following induction or consolidation chemotherapy to reduce time to neutrophil recovery and the duration of fever in adults)	Bone Marrow Transplant (BMT)	Peripheral Blood Progenitor Cell Collection and Therapy	Severe Chronic Neutropenia (To reduce the incidence and duration of neutropenia in symptomatic patients with congenital neutropenia, cyclic neutropenia, or idiopathic neutropenia)	Hematopoietic Syndrome of Acute Radiation Syndrome (To Increase survival in patients acutely exposed to myelosuppressive doses of radiation)
pegfilgrastim-pbbk (Fylnetra®) ⁺¹²	Amneal	X	--	--	--	--	--
sargramostim (Leukine®) ¹³	Partner	--	X	X ^b	X	--	X
tbo-filgrastim (Granix®) ¹⁴	Cephalon/Teva	X	--	--	--	--	--

† Pegfilgrastim-apgf (Nyvepria), pegfilgrastim-cbqv (Udenyca), pegfilgrastim-jmdb (Fulphila), pegfilgrastim-bmez (Ziextenzo), pegfilgrastim-fpgk (Stimufend), and pegfilgrastim-pbbk (Fylnetra) are biosimilars to the originator pegfilgrastim (Neulasta), approved through the FDA 351(k) biosimilar pathway.

^b For acceleration of myeloid reconstitution after PBPC/BMT; treatment of delayed neutrophil recovery or graft failure after BMT

OVERVIEW

Myelosuppressive chemotherapy can induce neutropenia (< 500 neutrophils/ μL or $< 1,000$ neutrophils/ μL and a predicted decline to $\leq 500/\mu\text{L}$ during the 48 hours after the dose) and febrile neutropenia ($\geq 38.3^\circ\text{C}$ orally or $\geq 38^\circ\text{C}$ sustained over 1 hour) which is a dose-limiting toxicity of chemotherapy.¹⁵ Febrile neutropenia can cause increased diagnostic and treatment costs, prolonged hospitalizations, and broad-spectrum antibiotic use which may necessitate chemotherapy dose reductions and treatment delays, and which may ultimately compromise treatment outcomes. The risk of febrile neutropenia is dependent on treatment and dose intensity, which are often underreported.

Colony stimulating factors (CSF) are hematopoietic growth factors that have been shown to decrease the likelihood of neutropenic complications resulting from chemotherapy and improve relative chemotherapy dose intensity. Prophylactic CSF use can reduce the severity, risk, and duration of febrile neutropenia and decrease rates of infection and hospitalizations. Eflapegrastim-xnst (Rolvedon), filgrastim (Neupogen), filgrastim-aafi (Nivestym), filgrastim-ayow (Releuko), filgrastim-sndz (Zarxio), pegfilgrastim (Neulasta), pegfilgrastim-apgf (Nyvepria), pegfilgrastim-cbqv (Udenyca), pegfilgrastim-jmdb (Fulphila), pegfilgrastim-bmez (Ziextenzo), pegfilgrastim-fpgk (Stimufend), pegfilgrastim-pbbk (Fylnetra), and tbo-filgrastim (Granix) are granulocyte colony-stimulating factors (G-CSF). Sargramostim (Leukine) is a granulocyte-macrophage colony stimulating factor (GM-CSF). CSFs act on hematopoietic cells and stimulate proliferation, differentiation commitment, and some end-cell functional activation.

The National Comprehensive Cancer Network (NCCN) v1.2023 updated practice guidelines for Hematopoietic Growth Factors base recommendations on evidence derived mainly from G-CSF studies in patients with solid tumors and lymphoid blood cancers.¹⁶ Safety data appear similar between filgrastim (Neupogen), pegfilgrastim (Neulasta), and their biosimilars, and the subcutaneous (SC) route is preferred for all agents. To date, there are insufficient head-to-head comparative studies on the clinical benefits of G-CSFs and GM-CSFs. Filgrastim, tbo-filgrastim, and pegfilgrastim administered SC have a category 1 recommendation stating there is high-level evidence from randomized, controlled clinical trials, and there is uniform NCCN consensus that they prophylactically reduce the risk of febrile neutropenia. However, the guidelines advise caution should be used with prophylactic use of G-CSFs administered with chemotherapy and radiation concurrently.¹⁷ The guidelines advise that FDA-approved biosimilars are a reasonable substitute for filgrastim and pegfilgrastim. Due to their recent approvals, pegfilgrastim-apgf (Nyvepria), filgrastim-ayow (Releuko), pegfilgrastim-fpgk (Stimufend), and pegfilgrastim-pbbk (Fylnetra) were not addressed by NCCN in their v1.2023 guidance. Eflapegrastim-xnst is included in the guideline and can be administered 24 hours after cytotoxic chemotherapy, but not within 14 days prior to 24 hours after chemotherapy. Filgrastim, filgrastim biosimilars, and tbo-filgrastim can be administered the day after chemotherapy, up to 3 to 4 days after chemotherapy, and through post-nadir recovery (category 1). Based on data from clinical trials, pegfilgrastim and its biosimilars should be administered the day after chemotherapy (category 1); however, administration up to 3 to 4 days after chemotherapy is also reasonable according to the NCCN guidelines. For patients unable to return to the clinic the next day for medication administration, a delivery device, Neulasta Onpro®, is available that allows for application to the patient the same day as chemotherapy administration, but the device does not release the medication until approximately 27 hours after application. There is evidence to support the use of chemotherapy regimens every 3 weeks with pegfilgrastim or its biosimilars (category 1). Efficacy data exist for pegfilgrastim products in chemotherapy regimens given every 2 weeks (category 2A). There are insufficient data to support dose/schedule of weekly

chemotherapy regimens; therefore, pegfilgrastim products should not be used. Sargramostim is no longer recommended for prophylactic use in patients with solid tumors receiving myelosuppressive chemotherapy. Since the approval of chimeric antigen receptor-modified T cell (CAR-T) therapies in recent years, NCCN advises against use of G-CSFs within 14 days after receipt of CAR-T therapy due to concern for exacerbation of cytokine release syndrome (CRS). Use after this time period is considered for treatment of neutropenia. Trilaciclib (Cosela™) was FDA approved in February 2021 to decrease the incidence of chemotherapy-induced myelosuppression in adults when administered prior to a platinum/etoposide-containing regimen or topotecan-containing regimen for extensive-stage small cell lung cancer.¹⁸ The NCCN recommends its use according to the approved indication (recommendation 2A); prophylactic G-CSF may be given after cycle 1.¹⁹ In clinical trials, the use of G-CSF was decreased in patients who received trilaciclib.²⁰ Trilaciclib is an intravenously-administered kinase inhibitor and will not be discussed in this Therapeutic Class Review.

Compared to prophylactic use, there is less evidence to support the CSFs for therapeutic use for febrile neutropenia and the guidelines note that it remains unknown whether CSF use translates into survival advantage.²¹ The NCCN guidelines recommend therapeutic treatment based on the patient's prophylactic therapy use. Patients presenting with febrile neutropenia who are on prophylactic filgrastim, filgrastim biosimilars, or tbo-filgrastim should continue the G-CSF. However, since pegfilgrastim, its biosimilars, and eflapegrastim-xnst (Rolvedon) are long acting, patients who received prophylactic pegfilgrastim products or eflapegrastim-xnst should not receive additional CSF. For patients who have not received prophylactic CSF, the guidelines recommend an evaluation of risk factors related to infection complications or poor clinical outcomes; if risk factors are present, then CSF should be considered. Filgrastim, filgrastim biosimilars, tbo-filgrastim, and sargramostim have a 2A recommendation (based on lower-level evidence, there is uniform NCCN consensus that the intervention is appropriate) for therapeutic use and can be used until post-nadir absolute neutrophil count (ANC) recovery to normal or near-normal levels. Pegfilgrastim, its biosimilars, and eflapegrastim-xnst have only been studied for prophylactic use.

The NCCN practice guidelines stratify patients into 3 risk groups based on the chemotherapy regimen and patient-related risk factors: high risk (> 20% risk of developing febrile neutropenia), intermediate risk (10% to 20% risk of developing febrile neutropenia), and low risk (< 10% risk of developing febrile neutropenia).²² It is recommended that high-risk patients receive prophylactic CSF regardless of the intent of treatment (category 1). If the patient falls into the intermediate risk group, NCCN recommends individualized consideration of CSF based on the likelihood of developing febrile neutropenia, consequences of developing febrile neutropenia, and the implications of interfering with chemotherapy treatments. Patients with ≥ 1 risk factor (e.g., age ≥ 65 years old, prior exposure to chemotherapy/radiation, poor performance status, recent surgery and/or open wounds, persistent neutropenia) should be considered for prophylactic therapy, whereas patients with no risk factors should be observed. Lastly, NCCN does not recommend the routine use of CSF in patients with low risk of developing febrile neutropenia due to lack of cost effectiveness and availability of alternative treatments but advises that CSF use can be considered for low-risk patients with risk factors. Choosing to administer a CSF may be considered if the treatment is curative or adjuvant and the patient is at serious medical consequences of febrile neutropenia.

The v1.2023 NCCN Hematopoietic Growth Factors guidelines also address the role of biosimilars.²³ The guidelines note that a biosimilar is a biological product that is highly similar to the FDA-approved originator product with very small, clinically inactive differences but no differences in efficacy, safety, or

purity. The first biosimilar, filgrastim-sndz, was approved in March 2015 with the second and third filgrastim biosimilars, filgrastim-aafi and filgrastim-ayow, approved in 2018 and 2022, respectively. However, tbo-filgrastim was approved in 2012 as a biologic rather than a biosimilar in the United States; in Europe, tbo-filgrastim is available as a biosimilar to filgrastim. The first pegfilgrastim biosimilars, pegfilgrastim-jmdb and pegfilgrastim-cbqv, were approved in 2018, followed by pegfilgrastim-bmez in 2019, pegfilgrastim-apgf in 2020, and both pegfilgrastim-pbbk and pegfilgrastim-fpgk in 2022. Studies have shown these products have similar safety and efficacy profiles as the originator product. The guidelines state if overall safety and efficacy are equivalent, biosimilars may be substituted for the originator product. While current biosimilars are not approved as interchangeable by the FDA, NCCN states limited data suggest that patients can alternate between the originator product and the biosimilar without any clinically meaningful differences regarding efficacy or safety. The guidelines also note that the use of biosimilars may provide opportunities for cost containment. The panel endorses the use of biosimilars of filgrastim and pegfilgrastim for myelosuppressive doses of radiation. For mobilization of hematopoietic progenitor cells in the autologous setting, the use of concurrent filgrastim (or its biosimilars) or tbo-filgrastim with (category 2B) or without (category 2A) sargramostim is recommended. Filgrastim, tbo-filgrastim, or filgrastim biosimilars are all NCCN category 1 G-CCF options for prophylaxis of febrile neutropenia and maintenance of scheduled dose delivery. Based on data from the World Marrow Donor Association (WMDA), NCCN recommends filgrastim or filgrastim biosimilars (category 2A) or tbo-filgrastim (category 2B) for the mobilization of peripheral blood progenitor cells in healthy donors in the allogeneic setting. The NCCN Panel does not recommend pegfilgrastim or its biosimilars for mobilization at this time.

The American Society of Clinical Oncology (ASCO) published an updated set of guidelines in 2015 for the use of white blood cell growth factors.²⁴ They note the ability of these agents to reduce the duration and severity of neutropenia and febrile neutropenia and to allow more intensive or dose-dense chemotherapy. The guidelines make no recommendation regarding the equivalency of the 2 CSFs, G-CSFs and GM-CSFs. Pegfilgrastim, filgrastim, filgrastim-sndz, and tbo-filgrastim can be used for the prevention of treatment-related febrile neutropenia; other biosimilars for filgrastim and the biosimilars for pegfilgrastim were not available at the time this guideline was published, and therefore, are not specifically addressed. ASCO advises that the choice of agent should be based on the clinical situation, convenience, and cost (Type: evidence-based, benefit outweigh harms; Evidence quality: high; Strength of recommendation: strong). The recommendations for the use of CSF for primary prophylaxis include the prevention of febrile neutropenia in patients who are at high risk based on age, medical history, disease characteristics, and myelotoxicity of the chemotherapy regimen. Clinical trial data support the use of CSF when the risk of febrile neutropenia is $\geq 20\%$ starting with the first cycle and continuing through subsequent cycles of chemotherapy. The decision on whether to use prophylactic CSF should take into consideration concerns such as the optimal chemotherapy regimen, individual patient risk factors, and the intention of treatment, that is, curative, prolongation of life, or symptom control and palliation. Alternative, but equally effective and safe, chemotherapy options not requiring CSFs should be considered (Type: evidence-based, benefit outweigh harms; Evidence quality: high; Strength of recommendation: strong). ASCO recommends secondary prophylaxis with CSF for patients who experienced a neutropenic complication from a prior cycle of chemotherapy (for which primary prophylaxis was not received), in which a reduced dose may compromise disease-free, overall survival, or treatment outcome. In many clinical situations, dose reduction or delay may be a reasonable alternative (Type: evidence-based, benefits outweigh harms; Evidence quality: high; Strength of recommendation: strong). CSF should not be routinely used for patients with neutropenia who are

afebrile (Type: evidence-based, benefits outweigh harms; Evidence quality: high; Strength of recommendation: strong). CSF should not be routinely used as adjunctive treatment with antibiotic therapy for patients with fever and neutropenia. However, CSF should be considered in patients with fever and neutropenia who are at high-risk for infection-related complications or who have prognostic factors that are predictive of poor clinical outcomes (Type: evidence-based, benefits outweigh harms; Evidence quality: high; Strength of recommendation: strong). High-risk features include expected prolonged (> 10 days) and profound (< 0.1 x 10⁹/L) neutropenia, age > 65 years, uncontrolled primary disease, pneumonia, hypotension and multiorgan dysfunction (sepsis syndrome), invasive fungal infection, or being hospitalized at the time of the development of fever.

Per the 2015 ASCO guidelines, CSFs can be used during or after chemotherapy or with plerixafor (Mozobil®) to mobilize peripheral-blood progenitor cells (PBPC) depending on the type of cancer and transplantation (Type: evidence-based, benefit outweigh harms; Evidence quality: strong; Strength of recommendation: high). CSF should be administered after autologous stem-cell transplantation (Type: evidence-based, benefits outweigh harms; Evidence quality: high; Strength of recommendation: strong) and may be administered after allogenic stem-cell transplantation (Type: evidence-based; Evidence quality: low; Strength of recommendation: weak). They recommend filgrastim-sndz; future biosimilars may be used for the prevention of neutropenia. ASCO recommends that the choice of agent depends on convenience, cost, and clinical situation. ASCO's dosing and administration recommendations for filgrastim-sndz are identical to those for filgrastim.

PHARMACOLOGY^{25,26,27,28,29,30,31,32,33,34,35,36,37,38}

Sargramostim (Leukine) is a recombinant human GM-CSF produced by recombinant DNA technology in yeast. GM-CSF is a hematopoietic growth factor, which triggers proliferation and differentiation of hematopoietic progenitor cells. GM-CSF is included in a group of growth factors termed CSF that promote survival, clonal expansion, and differentiation of hematopoietic progenitor cells. GM-CSF induces partially-committed progenitor cells to divide and differentiate in the granulocyte-macrophage pathways which include neutrophils, monocytes/macrophages, and myeloid derived dendritic cells. GM-CSF also has the capability of activating mature granulocytes and macrophages. GM-CSF is a multilineage factor and, in addition to dose-dependent effects on the myelomonocytic lineage, can encourage the proliferation of megakaryocytic and erythroid progenitors.

Eflapegrastim-xnst (Rolvedon), filgrastim (Neupogen), filgrastim-aafi (Nivestym), filgrastim-ayow (Releuko), filgrastim-sndz (Zarxio), pegfilgrastim (Neulasta), pegfilgrastim-apgf (Nyvepria), pegfilgrastim-cbqv (Udenyca), pegfilgrastim-jmdb (Fulphila), pegfilgrastim-bmez (Ziextenzo), pegfilgrastim-fpgk (Stimufend), pegfilgrastim-pbbk (Fylnetra), and tbo-filgrastim (Granix) are G-CSFs that are produced by recombinant technology using *Escherichia coli*. G-CSF acts on hematopoietic cells by binding to specific cell surface receptors, thereby stimulating differentiation, commitment, proliferation, and end cell functional activation. Filgrastim, filgrastim-aafi, filgrastim-ayow, filgrastim-sndz, and tbo-filgrastim stimulate the growth and development of neutrophils within the bone marrow. Labeling for filgrastim, filgrastim-aafi, filgrastim-ayow, and filgrastim-sndz report a shift towards earlier granulocyte progenitor cells (left shift), including the appearance of promyelocytes and myeloblasts, usually during neutrophil recovery following the chemotherapy-induced nadir; this shift is not reported in the labeling for tbo-filgrastim. Studies on cellular proliferation, receptor binding, and neutrophil function demonstrate that filgrastim and pegfilgrastim have the same mechanism of action. Pegfilgrastim, pegfilgrastim-apgf, pegfilgrastim-cbqv, pegfilgrastim-jmdb, pegfilgrastim-fpgk, and pegfilgrastim-pbbk are conjugates of

filgrastim and monomethoxypolyethylene glycol. Neutrophil receptor binding is important for the clearance of pegfilgrastim and its biosimilars; serum clearance is directly related to the level of neutrophils. Filgrastim-aafi (Nivestym), filgrastim-ayow (Releuko), and filgrastim-sndz (Zarxio) are biosimilars for filgrastim (Neupogen); pegfilgrastim-apgf (Nyvepria), pegfilgrastim-cbqv (Udenyca), pegfilgrastim-jmdb (Fulphila), pegfilgrastim-bmez (Ziextenzo), **pegfilgrastim-fpgk (Stimufend)**, and **pegfilgrastim-pbbk (Fylnetra)** are biosimilars to pegfilgrastim (Neulasta).

PHARMACOKINETICS^{39,40,41,42,43,44,45,46,47,48,49,50,51,52}

Drug	Half-Life Intravenous Administration	Half-Life Subcutaneous Administration
eflapegrastim-xnst (Rolvedon)	--	36.4 hours
filgrastim (Neupogen)	231 minutes	210 minutes
filgrastim-aafi (Nivestym)		
filgrastim-ayow (Releuko)		
filgrastim-sndz (Zarxio)		
pegfilgrastim (Neulasta)	--	15–80 hours
pegfilgrastim-apgf (Nyvepria)		
pegfilgrastim-cbqv (Udenyca)		
pegfilgrastim-jmdb (Fulphila)		
pegfilgrastim-bmez (Ziextenzo)		
pegfilgrastim-fpgk (Stimufend)		
pegfilgrastim-pbbk (Fylnetra)	3.84 hours	1.4 hours
sargramostim (Leukine)		
tbo-filgrastim (Granix)	--	3–3.5 hours

CONTRAINDICATIONS/WARNINGS^{53,54,55,56,57,58,59,60,61,62,63,64,65,66}

eflapegrastim-xnst (Rolvedon), filgrastim (Neupogen), filgrastim-aafi (Nivestym), filgrastim-ayow (Releuko), filgrastim-sndz (Zarxio), pegfilgrastim (Neulasta), pegfilgrastim-apgf (Nyvepria), pegfilgrastim-cbqv (Udenyca), pegfilgrastim-jmdb (Fulphila), pegfilgrastim-bmez (Ziextenzo), **pegfilgrastim-fpgk (Stimufend)**, **pegfilgrastim-pbbk (Fylnetra)**, and tbo-filgrastim (Granix)

Eflapegrastim-xnst, filgrastim, filgrastim-aafi, filgrastim-ayow, filgrastim-sndz, pegfilgrastim, pegfilgrastim-apgf, pegfilgrastim-cbqv, pegfilgrastim-jmdb, pegfilgrastim-bmez, **pegfilgrastim-fpgk**, **pegfilgrastim-pbbk**, and tbo-filgrastim are contraindicated in patients with known hypersensitivity to **eflapegrastim products**, filgrastim products, pegfilgrastim products, or any other component of medication.

Uses of **eflapegrastim-xnst**, filgrastim, filgrastim biosimilars, pegfilgrastim, pegfilgrastim biosimilars, and tbo-filgrastim have reported allergic-type reactions during initial or subsequent treatments, including anaphylactic reactions. However, a majority of allergic type reactions occurred with initial exposure. These have generally been characterized by systemic symptoms involving at least 2 body systems, most often skin (e.g., rash, urticaria, facial edema), cardiovascular (e.g., hypotension, tachycardia), and

respiratory (e.g., wheezing, dyspnea). In rare cases following administration of pegfilgrastim, filgrastim, or their biosimilars, allergic reactions, including anaphylaxis, recurred within days after the initial anti-allergic treatment was discontinued; recurrence is not discussed in the labeling for eflapegrastim-xnst or tbo-filgrastim. If a serious allergic reaction occurs, appropriate therapy should be administered, with close patient follow-up over several days. Administration of steroids, bronchodilators, antihistamines, and/or epinephrine may reduce the severity of symptoms. If rechallenged, symptoms recurred in more than half the patients. Filgrastim, filgrastim biosimilars, pegfilgrastim, pegfilgrastim biosimilars, and tbo-filgrastim should be permanently discontinued in patients with serious allergic reactions. The needle cap for the single-dose prefilled syringe contains latex; therefore, persons with latex allergies should not administer pegfilgrastim, pegfilgrastim-bmez, filgrastim, or filgrastim-sndz.

Splenic rupture, including fatal cases, has been reported following the administration of eflapegrastim-xnst, filgrastim, filgrastim biosimilars, pegfilgrastim, pegfilgrastim biosimilars, and tbo-filgrastim. Patients receiving any agent who report left upper abdominal and/or shoulder pain should be evaluated for an enlarged spleen or splenic rupture.

The on-body injector (Neulasta Onpro) for pegfilgrastim (Neulasta) delivery kit uses acrylic adhesive; therefore, its use may result in serious reactions in patients who have allergies to acrylic adhesives. Patients should avoid activities such as traveling, driving, or operating heavy machinery 26 to 29 hours following application of the on-body injector for pegfilgrastim. This includes the 45-minute delivery period, plus 1 hour post-delivery. A caregiver should be close by for the first use. Use of the on-body injector is not recommended for patients with hematopoietic acute radiation syndrome (H-ARS), and it has not been studied in pediatric patients. Missed or partial doses of pegfilgrastim have been reported with the on-body injector. Patients should be aware to notify their healthcare professional (HCP) immediately if they suspect a missed or partial dose has been delivered in order to get replacement medication. If a dose is missed, a new dose should be administered manually by a single prefilled syringe as soon as possible. The pegfilgrastim on-body injector is intended to be used in an electromagnetic environment; however, it should not be exposed to medical imaging studies (e.g., X-ray scan, magnetic resonance imaging [MRI], computerized tomography [CT] scan, ultrasound), oxygen-rich environments (e.g., hyperbaric chambers), or radiation treatment, as it may cause injector damage or patient injury. The injector must be at least 4 inches away from electrical equipment, including cell phones, cordless telephones, microwaves, and other common appliances, as these may interfere with operation and can lead to a missed or incomplete dose. The pegfilgrastim injector should not be used in hot tubs, saunas, and whirlpools. Direct sunlight should be avoided and the injector should be worn under clothing. Patients should not sleep on the injector or apply pressure during wear. Lotions, creams, oils, and cleaning agents can loosen the adhesive and should be avoided near the injector. Patients should avoid activities and places that may interfere with monitoring during the dosing (e.g., traveling, driving, or operating heavy machinery during hours 26 to 29 following application of the on-body injector).

Acute respiratory distress syndrome (ARDS) has been reported in patients receiving eflapegrastim-xnst, filgrastim, filgrastim biosimilars, pegfilgrastim, pegfilgrastim biosimilars, and tbo-filgrastim, and is suggested to be secondary to an invasion of neutrophils to sites of inflammation in the lungs. Patients receiving eflapegrastim-xnst, filgrastim, filgrastim biosimilars, pegfilgrastim, pegfilgrastim biosimilars, or tbo-filgrastim who develop fever, lung infiltrates, or respiratory distress should be evaluated for the possibility of ARDS. If ARDS occurs, eflapegrastim-xnst, filgrastim, filgrastim biosimilars, pegfilgrastim, pegfilgrastim biosimilars, and tbo-filgrastim should be discontinued and/or withheld until resolution of ARDS, and patients should receive appropriate medical management.

Healthy donors undergoing PBPC mobilization have reported alveolar hemorrhage manifesting as pulmonary infiltrates and hemoptysis and have required hospitalization. Hemoptysis resolved with discontinuation of filgrastim products. The use of filgrastim or filgrastim biosimilars for PBPC mobilization in healthy donors is not an approved indication. In cancer patients using filgrastim or its biosimilars, the medication should be discontinued if the leukocyte count is $> 100,000/\text{mm}^3$. Alveolar hemorrhage has also been reported in the post-marketing setting for pegfilgrastim and its biosimilars. **Eflapegrastim-xnst**, pegfilgrastim products and tbo-filgrastim are not indicated for PBPC mobilization.

Severe sickle cell crisis, some cases resulting in death, have been associated with the use of **eflapegrastim-xnst**, filgrastim, filgrastim biosimilars, pegfilgrastim, pegfilgrastim biosimilars, and tbo-filgrastim in patients with sickle cell disorders. After careful consideration of the potential risks and benefits, only physicians qualified by specialized training or experience in the treatment of patients with sickle cell disorders should prescribe filgrastim, filgrastim biosimilars, pegfilgrastim, pegfilgrastim biosimilars, or tbo-filgrastim for such patients. Patients experiencing a sickle cell crisis should discontinue therapy.

The G-CSF receptor that **eflapegrastim-xnst**, filgrastim, pegfilgrastim, and tbo-filgrastim act upon has been found on tumor cell lines. It is possible that **eflapegrastim-xnst**, filgrastim, pegfilgrastim, and tbo-filgrastim act as growth factors for any tumor type; however, more data are needed since the limited data that are available are inconclusive. This also applies to biosimilar products of filgrastim and pegfilgrastim. The safety of filgrastim and its biosimilars in treating chronic myeloid leukemia (CML) and myelodysplasia has not been established.

The safety and efficacy of filgrastim products used concurrently with chemotherapy have not been established. However, due to the sensitivity of rapidly dividing myeloid cells during chemotherapy, the use of filgrastim products should be avoided for 24 hours before and through 24 hours after chemotherapy. Concurrent therapy with radiation should also be avoided.

For breast and lung cancer patients, **eflapegrastim-xnst**, filgrastim, pegfilgrastim, and their biosimilars are associated with an increased risk of myelodysplastic syndrome (MDS) and acute myeloid leukemia (AML) when used in conjunction with chemotherapy and/or radiation. Patients should be monitored for signs and symptoms of MDS and AML.

The diagnosis of severe chronic neutropenia (SCN) should be confirmed prior to initiating therapy with filgrastim or its biosimilars since cytogenetic abnormalities and transformation to MDS and AML have been observed in patients treated with filgrastim products for SCN. The risk of developing MDS and AML appears to be confined to the subset of patients with congenital neutropenia. Eventual development of myeloid leukemia has been associated with abnormal cytogenetics and MDS. The effect of filgrastim or its biosimilars on the development of abnormal cytogenetics and the effect of their continued administration in patients with abnormal cytogenetics or MDS are unknown. The risks and benefits of continuing filgrastim and its biosimilars should be carefully considered if a patient with SCN develops abnormal cytogenetics or myelodysplasia.

Cases of moderate to severe cutaneous vasculitis have been reported in patients taking filgrastim products. These cases most often occurred in patients with SCN receiving long-term filgrastim products. Therapy with filgrastim, including its biosimilars, should be held if cutaneous vasculitis occurs and may be restarted at a reduced dose when symptoms resolve and the ANC has decreased.

Thrombocytopenia has been reported in patients taking **eflapegrastim-xnst**, filgrastim, pegfilgrastim, their biosimilars, and tbo-filgrastim; therefore, platelet counts should be monitored.

Capillary leak syndrome (CLS) can occur in patients using human G-CSFs. Symptoms include hypotension, hypoalbuminemia, edema, and hemoconcentration and can be life-threatening, especially if treatment is delayed. Patients who develop symptoms of CLS should be monitored closely and receive symptomatic treatment.

Glomerulonephritis has occurred in patients taking **eflapegrastim-xnst**, filgrastim, or pegfilgrastim products, including their biosimilars, based on findings of azotemia, hematuria, proteinuria, and renal biopsy. Reducing the dose or discontinuing filgrastim or pegfilgrastim products should result in resolution of glomerulonephritis.

White blood cell (WBC) counts $\geq 100,000/\text{mm}^3$ were observed in approximately 2% of patients taking filgrastim at doses $> 5 \text{ mcg/kg/day}$ when receiving myelosuppressive chemotherapy and have been reported in patients receiving **eflapegrastim-xnst** and pegfilgrastim products (WBC $\geq 100 \times 10^9/\text{L}$) and tbo-filgrastim (WBC $> 100,000 \times 10^6/\text{L}$). In order to avoid potential risks of excessive leukocytosis, it is recommended that **eflapegrastim-xnst**, filgrastim, filgrastim biosimilars, and tbo-filgrastim be stopped if the ANC surpasses $10,000/\text{mm}^3$ after the chemotherapy-induced ANC nadir occurs. Dosages of filgrastim, filgrastim biosimilars, or tbo-filgrastim that increase ANC $> 10,000/\text{mm}^3$ may not result in additional clinical benefit. Patients using filgrastim, filgrastim biosimilars, and tbo-filgrastim should have their complete blood count (CBC) monitored at least twice weekly during therapy. Monitoring of CBC is also recommended for **eflapegrastim-xnst** and for pegfilgrastim products, although a monitoring schedule is not specified. Filgrastim and its biosimilars should be discontinued if the leukocyte count becomes $> 100,000/\text{mm}^3$ during the period of administration for PBPC mobilization.

When interpreting bone-imaging results, it should be noted that increased hematopoietic activity of bone marrow in response to growth factor has been associated with transient positive bone-imaging changes.

Aortitis has been reported in patients taking **eflapegrastim-xnst**, filgrastim, and pegfilgrastim products as early as the first week of therapy. Signs and symptoms of aortitis include fever, abdominal pain, malaise, back pain, and increased inflammatory markers. If suspected, the product should be discontinued.

sargramostim (Leukine)

Sargramostim (Leukine) is contraindicated in patients with a history of serious allergic reactions to human granulocyte-macrophage colony stimulating factor, yeast-derived products, or any component of the product; anaphylactic reactions have been reported.

Sargramostim should not be administered simultaneously with cytotoxic chemotherapy or radiotherapy or within 24 hours preceding or following chemotherapy or radiotherapy due to the potential sensitivity of rapidly dividing hematopoietic progenitor cells.

Sargramostim contains benzyl alcohol in the liquid formulations and in the bacteriostatic water for injection diluent. Benzyl alcohol has been reported to be associated with a fatal “gaspings syndrome” in premature and low birth weight infants. Formulations containing benzyl alcohol should not be administered to neonates. Instead, administer lyophilized sargramostim reconstituted with sterile water for injection.

Reports of fluid retention (11%), edema, capillary leak syndrome (< 1%), and pleural (1%) and/or pericardial effusion (4%) have been reported in patients after sargramostim administration. In patients with pre-existing pleural and pericardial effusions, the use of sargramostim may worsen fluid retention; however, fluid retention linked with or worsened by sargramostim has been reversible after interruption or dose reduction of sargramostim, with or without diuretic therapy. Patients with pre-existing fluid retention, pulmonary infiltrates, or congestive heart failure should use sargramostim with caution.

Transient supraventricular arrhythmia has been reported in uncontrolled studies during sargramostim administration especially in patients with a history of cardiac arrhythmia. These arrhythmias have been reversible after discontinuation of sargramostim. Sargramostim should be used with caution in patients with pre-existing cardiac disease.

Respiratory distress, hypoxia, flushing, hypotension, syncope, and/or tachycardia have been reported following the first administration of sargramostim in a particular treatment cycle. These signs have resolved with symptomatic treatment and usually do not reappear with successive doses in the same cycle of treatment. If a patient displays symptoms, the rate of infusion should be reduced by 50% and therapy should be discontinued if symptoms persist.

Stimulation of marrow precursors may cause an increase in WBC count. If the ANC exceeds 20,000 cells/mm³ sargramostim therapy should be stopped or the dose reduced in half. Within 3 to 7 days following therapy termination, excessive blood counts should return to normal or baseline levels. Twice weekly CBC with differential levels should be performed thereafter.

Sargramostim primarily stimulates normal myeloid precursors but may act as a growth factor for any tumor type including myeloid malignancies; therefore, caution should be used when prescribing sargramostim in patients with any malignancy with myeloid characteristics. If disease progression occurs during sargramostim therapy, the medication should be discontinued.

DRUG INTERACTIONS^{67,68,69,70,71,72,73,74,75,76,77,78,79,80}

Interactions between agents in this class and other drugs have not been fully evaluated.

Drugs that may potentiate the myeloproliferative effects of these agents, such as lithium and corticosteroids, should be used with caution. Patients receiving lithium and these agents should have more frequent monitoring of neutrophil counts.

ADVERSE EFFECTS^{81,82,83,84,85,86,87,88,89,90,91,92,93,94}

Drug	Bone (Skeletal) Pain	Pyrexia	Skin Rash	Edema
eflapegrastim-xnst (Rolvedon)	38	28	25	18
filgrastim (Neupogen), filgrastim-aafi (Nivestym), filgrastim-ayow (Releuko), filgrastim-sndz (Zarxio) cancer patients receiving chemotherapy	11 (6)	48 (29)	14 (5)	reported
pegfilgrastim (Neulasta), pegfilgrastim-apgf (Nyvepria), pegfilgrastim-cbqv (Udenyca), pegfilgrastim-jmdb (Fulphila), pegfilgrastim-bmez (Ziextenzo), pegfilgrastim-fpgk (Stimufend), pegfilgrastim-pbbk (Fylnetra)	31	reported	reported	reported
sargramostim (Leukine) patients with AML	21 (5)	81–95 (74–96)	44–70 (38–73)	13–34 (11–35)
tbo-filgrastim (Granix)	3.4 (1.4)	8 (nr)	reported	reported

Adverse effects are reported as a percentage. Adverse effects data are obtained from package inserts and are not meant to be comparative or all inclusive. nr = not reported.

Lactate dehydrogenase and alkaline phosphatase occurred in 6% of 294 patients receiving filgrastim following chemotherapy treatment.

Adverse reactions with $\geq 10\%$ incidence in eflapegrastim-xnst through week 14 in patients with early-stage breast cancer include fatigue, nausea, diarrhea, headache, anemia, myalgia, arthralgia, back pain, decreased appetite, abdominal pain, dizziness, dyspnea, cough, thrombocytopenia, pain, pain in extremity, local administration reactions, and flushing.

There is a potential for immunogenicity when using therapeutic proteins; however, the incidence of antibody development in patients taking filgrastim, filgrastim-aafi, filgrastim-ayow, filgrastim-sndz, or tbo-filgrastim has not been adequately determined. In clinical studies, the incidence of antibodies binding to filgrastim was 3% but no evidence of a neutralizing response was observed. In clinical trials the incidence of binding antibodies to tbo-filgrastim was 1.4% in patients who received prior chemotherapy; no cross-reactive antibodies were detected, and all antibody responses were transient and of low titers. In studies, evidence of binding antibodies was detected ($< 1\%$) with pegfilgrastim use but there was no evidence of neutralizing antibodies. In clinical trials, antibodies to eflapegrastim-xnst were detected in 7.1% of patients during a 12-week treatment period, with neutralizing antibodies detected in 0.3% of patients post-treatment. Cytopenias resulting from an antibody response to exogenous growth factor have been reported rarely. Antibody studies involving patients with Crohn's disease, melanoma in complete remission (unapproved use), and a variety of other diseases revealed that 1.3%, 82.9%, and 2.3%, respectively, had neutralizing anti-sargramostim antibodies detected. The incidence of anti-sargramostim neutralizing antibodies may be related to duration of exposure to the product.

Adverse reactions with $\geq 2\%$ higher incidence in filgrastim and its biosimilars compared to placebo-treated patients for AML include epistaxis, back pain, extremity pain, erythema, maculopapular rash, diarrhea, constipation, and transfusion reaction. Adverse reactions with $\geq 5\%$ higher incidence in filgrastim and its biosimilars patients compared to placebo-treated patients for BMT include rash, hypersensitivity, thrombocytopenia, anemia, hypertension, sepsis, bronchitis, and insomnia. Adverse reactions for the treatment of PBPC when using filgrastim or its biosimilars include bone pain (30%), pyrexia (16%), alkaline phosphatase increases (11%), and headache (10%). Adverse reactions with $\geq 5\%$ higher incidence in patients treated with filgrastim, or its biosimilars, compared to placebo when treating SCN include arthralgia, bone pain, back pain, muscle spasms, musculoskeletal pain, extremity pain, splenomegaly, anemia, upper respiratory tract infection, urinary tract infection, epistaxis, chest pain, diarrhea, hypoesthesia, and alopecia.

Monitoring

For cancer patients receiving myelosuppressive chemotherapy, a CBC and platelet count should be obtained prior to chemotherapy and at regular intervals (twice weekly) during treatment with filgrastim or pegfilgrastim products; if ANC increases beyond $10,000/\text{mm}^3$ therapy termination should be considered. For cancer patients receiving BMT, frequent CBC and platelet counts are recommended during filgrastim therapy, including biosimilars.

For patients with severe chronic neutropenia, serial CBC with differential, platelet counts, and an evaluation of bone marrow morphology and karyotype should be performed prior to starting therapy. Twice weekly CBC with differential and platelet counts should be obtained during the initial 4 weeks of therapy with filgrastim or its biosimilars and during the 2 weeks following any dose adjustment. Once a patient is clinically stable, a monthly CBC with differential and platelet count should be performed during the first year of therapy with filgrastim or its biosimilars. Thereafter, quarterly CBC with differential and platelet counts is recommended.

Patients using filgrastim, filgrastim-aafi, or filgrastim-sndz for PBPC mobilization should have their neutrophil counts monitored after 4 days of therapy and discontinued therapy if the WBC count increases $> 100,000/\text{mm}^3$.

For hematopoietic syndrome of acute radiation syndrome, a baseline CBC and serial CBC should be taken approximately every third day until the ANC remains $> 1,000/\text{mm}^3$ for 3 consecutive CBCs when using filgrastim or sargramostim. Filgrastim and sargramostim therapy should not be delayed if a CBC is not readily available. Filgrastim and sargramostim therapy should continue until the ANC remains $> 1,000/\text{mm}^3$ for 3 consecutive CBCs or exceeds $10,000/\text{mm}^3$ after radiation-induced nadir. When using pegfilgrastim, a baseline CBC should be obtained. However, therapy should not be delayed if a CBC is not readily available. The amount of radiation absorbed should be estimated based on information from public authorities, biodosimetry, or clinical findings, such as time to onset of vomiting or lymphocyte depletion kinetics.

With sargramostim, a CBC is recommended twice per week to avoid excessive leukocytosis (WBC $> 50,000$ cells/ mm^3 ; ANC $> 20,000$ cells/ mm^3). Patients with pre-existing renal or hepatic dysfunction should have their renal and hepatic function monitored at least biweekly during sargramostim therapy. Patient hydration and body weight should be monitored carefully during therapy.

SPECIAL POPULATIONS^{95,96,97,98,99,100,101,102,103,104,105,106,107,108}

Pediatrics

The safety and efficacy of eflapegrastim-xnst (Rolvedon) have not been established in pediatric patients.

The safety and efficacy of filgrastim (Neupogen), filgrastim-aafi (Nivestym), filgrastim-ayow (Releuko), filgrastim-sndz (Zarxio), pegfilgrastim (Neulasta), pegfilgrastim-apgf (Nyvepria), pegfilgrastim-cbqv (Udenyca), pegfilgrastim-jmdb (Fulphila), pegfilgrastim-bmez (Ziextenzo), pegfilgrastim-fpgk (Stimufend), pegfilgrastim-pbbk (Fylnetra), tbo-filgrastim (Granix), and sargramostim have been established in the treatment of pediatric patients. The pharmacokinetics of filgrastim in pediatric patients after chemotherapy are similar to those in adults receiving the same weight-normalized doses, suggesting no age-related differences in the pharmacokinetics of this product. Similarly, there are no overall differences in safety between adults and pediatrics based on postmarketing surveillance and review of scientific literature when using pegfilgrastim. Pharmacokinetics and safety of tbo-filgrastim in pediatrics is similar to what is seen in adults. Safety and efficacy of sargramostim have been established in pediatrics ≥ 2 years based on clinical studies in adults for autologous peripheral blood progenitor cells and bone marrow transplantation, allogenic bone marrow transplantation, and treatment of delayed neutrophil recovery or graft failure. The adverse reactions were consistent with those reported in adults. Approval of sargramostim for pediatric patients from birth to 17 years of age acutely exposed to myelosuppressive doses of radiation was based on animal studies since human studies with H-ARS could not be conducted due to ethical and feasibility reasons.

The pegfilgrastim on-body injector (Neulasta Onpro) has not been studied in pediatrics.

Pregnancy

For all agents in this class review, data on use during pregnancy are insufficient to inform prescribers of the drug-associated risk; the benefits of the drug should outweigh the risks when prescribing to pregnant women. The drug labels for eflapegrastim-xnst, filgrastim, filgrastim-aafi, filgrastim-ayow, filgrastim-sndz, tbo-filgrastim, pegfilgrastim, pegfilgrastim-apgf, pegfilgrastim-cbqv, pegfilgrastim-jmdb, pegfilgrastim-bmez, pegfilgrastim-fpgk, pegfilgrastim-pbbk, and sargramostim comply with the Pregnancy and Lactation Labeling Rule (PLLR).

Geriatric

In clinical studies, there were no overall differences in safety or efficacy between patients ≥ 65 years of age and younger patients treated with eflapegrastim-xnst, filgrastim, tbo-filgrastim, or pegfilgrastim.

Clinical trials with sargramostim did not include a sufficient number of patients ≥ 65 years of age to compare response to younger individuals.

DOSAGES^{109,110,111,112,113,114,115,116,117,118,119,120,121,122}

Drug	Cancer patients receiving myelosuppressive chemotherapy	Acute Myeloid Leukemia (AML) patients receiving chemotherapy	Bone Marrow Transplant (BMT)	Peripheral Blood Progenitor Cell Collection and Therapy	Severe Chronic Neutropenia	Hematopoietic Syndrome of Acute Radiation Syndrome	Availability
eflapegrastim-xnst (Rolvedon)	13.2 mg SC once per chemotherapy cycle	--	--	--	--	--	Single-dose prefilled syringe: 13.2 mg/0.6 mL
filgrastim (Neupogen)	5 mcg/kg/day, administered as a single daily injection by SC bolus, by short IV infusion (15 to 30 minutes), or continuous IV infusion; doses may be increased by increments of 5 mcg/kg for each chemotherapy cycle according to the duration and severity of the ANC nadir	5 mcg/kg/day as single daily injection by SC injection, short IV infusion (15 to 30 minutes), or continuous IV infusion; doses may be increased by increments of 5 mcg/kg for each chemotherapy cycle according to the duration and severity of the ANC nadir	10 mcg/kg/day given as an IV infusion no longer than 24 hours; during periods of neutrophil recovery, the daily dose should be titrated against the neutrophil response dosing schedule	10 mcg/kg/day SC; give for at least 4 days before the first leukapheresis procedure and continued until the last leukapheresis	Starting Dose: Congenital Neutropenia: 6 mcg/kg SC twice daily Idiopathic or Cyclic Neutropenia: 5 mcg/kg SC daily	10 mcg/kg as single daily SC injection; administer as soon as possible after suspected/confirmed exposure to radiation doses > 2 gray (Gy)	Single-dose vials: 300 mcg/1 mL, 480 mcg/1.6 mL Prefilled single-use syringes (SingleJect®): 300 mcg/0.5 mL, 480 mcg/0.8 mL
filgrastim-aafi (Nivestym)				--		Single-dose vials: 300 mcg/1 mL, 480 mcg/1.6 mL Prefilled single-dose syringes: 300 mcg/0.5 mL, 480 mcg/0.8 mL	
filgrastim-ayow (Releuko)				--		Single-dose vials: 300 mcg/1 mL, 480 mcg/1.6 mL Prefilled single-dose syringes: 300 mcg/0.5 mL, 480 mcg/0.8 mL	
filgrastim-sndz (Zarxio)				--		Prefilled single-dose syringes: 300 mcg/0.5 mL, 480 mcg/0.8 mL	

Dosages (continued)

Drug	Cancer patients receiving myelosuppressive chemotherapy	Acute Myeloid Leukemia (AML) patients receiving chemotherapy	Bone Marrow Transplant (BMT)	Peripheral Blood Progenitor Cell Collection and Therapy	Severe Chronic Neutropenia	Hematopoietic Syndrome of Acute Radiation Syndrome	Availability
pegfilgrastim (Neulasta)	6 mg SC once per chemotherapy cycle; pediatric (weight < 45 kg) dosing is weight-based per dosing schedule in drug PI	--	--	--	--	2 doses, 6 mg each, given SC 1 week apart; pediatric (weight < 45 kg) dosing is weight based per dosing schedule in drug PI; first dose given as soon as possible after suspected / confirmed radiation exposure of > 2 gray; second dose given 1 week after	Neulasta: single-use prefilled syringe for manual use: 6 mg/0.6 mL Neulasta Onpro® Delivery Kit: single-use delivery kit: 1 prefilled syringe (6 mg/0.6 mL) copackaged with 1 on-body injector for HCP administration
pegfilgrastim-apgf (Nyvepria)						--	Single-dose prefilled syringe: 6 mg/0.6 mL
pegfilgrastim-cbqv (Udenyca)						2 doses, 6 mg each, given SC 1 week apart; pediatric (weight < 45 kg) dosing is weight based per dosing schedule in drug PI; first dose given as soon as possible after suspected / confirmed radiation exposure of > 2 gray; second dose given 1 week after	Single-dose prefilled syringe: 6 mg/0.6 mL
pegfilgrastim-jmdb (Fulphila)						--	Single-dose prefilled syringe: 6 mg/0.6 mL
pegfilgrastim-bmez (Ziextenzo)						--	Single-dose prefilled syringe: 6 mg/0.6 mL

PI=Prescribing Information

Dosages (continued)

Drug	Cancer patients receiving myelosuppressive chemotherapy	Acute Myeloid Leukemia (AML) patients receiving chemotherapy	Bone Marrow Transplant (BMT)	Peripheral Blood Progenitor Cell Collection and Therapy	Severe Chronic Neutropenia	Hematopoietic Syndrome of Acute Radiation Syndrome	Availability
pegfilgrastim-fpgk (Stimufend)	6 mg SC once per chemotherapy cycle; pediatric (weight < 45 kg) dosing is weight-based per dosing schedule in drug PI	--	--	--	--	--	Single-dose prefilled syringe: 6 mg/0.6 mL
pegfilgrastim-pbbk (Fylnetra)		--	--	--	--	--	Single-dose prefilled syringe: 6 mg/0.6 mL
sargramostim (Leukine)	--	250 mcg/m ² /day given IV over 4 hours starting approximately day 11 or 4 days following completion of induction chemotherapy, if the day 10 bone marrow is hypoplastic with < 5% blasts	250mcg/m ² /day given IV over 2 hours beginning 2 to 4 hours after bone marrow infusion, and not < 24 hours after the last dose of chemotherapy or radiotherapy; therapy should not begin until post-marrow infusion ANC is < 500 cells/mm ³ and continued until ANC is > 1,500 cells/mm ³ for 3 consecutive days; treatment of BMT failure or engraftment delay: 250 mcg/m ² /day given IV over 2 hours for 14 days	Mobilization/collection: 250 mcg/m ² /day IV over 24 hours or SC once daily; continue at same dose through PBPC collection, usually after 5 days; if WBC > 50,000 cells/mm ³ reduce dose by 50% Transplantation: 250 mcg/m ² /day IV over 24 hours or SC once daily until ANC > 1,500 cells/mm ³ for 3 consecutive days	--	Administer weight-based dose as SC injection once daily after suspected or confirmed exposure to radiation doses > 2 gray; continue until ANC > 1,000/mm ³ for 3 consecutive CBCs or exceeds 10,000/mm ³ after radiation induced nadir Adults/pediatrics > 40 kg: 7 mcg/kg; pediatrics 15 kg to 40 kg: 10 mcg/kg; and pediatrics < 15kg: 12 mcg/kg	Single-dose vial: 250 mcg

Dosages (continued)

Drug	Cancer patients receiving myelosuppressive chemotherapy	Acute Myeloid Leukemia (AML) patients receiving chemotherapy	Bone Marrow Transplant (BMT)	Peripheral Blood Progenitor Cell Collection and Therapy	Hematopoietic Syndrome of Acute Radiation Syndrome	Severe Chronic Neutropenia	Availability
tbo-filgrastim (Granix)	5 mcg/kg/day as a SC injection until expected neutrophil nadir is passed and neutrophil count is in normal range; administer no earlier than 24 hours following myelosuppressive chemotherapy	--	--	--	--	--	Single-use, preservative-free, prefilled syringes: 300 mcg/0.5 mL, 480 mcg/0.8 mL with needle guard (HCP-administered); without needle guard (self- or caregiver-administered) Single use vials: 300 mcg/1 mL, 480 mcg/1.6 mL

Due to the potential sensitivity of rapidly dividing myeloid cells to chemotherapy, **eflapegrastim-xnst**, filgrastim, filgrastim biosimilars, and tbo-filgrastim should not be administered within 24 hours before administration of chemotherapy. These products should be administered no earlier than 24 hours after the administration of cytotoxic chemotherapy. Filgrastim or its biosimilars should be administered daily for up to 2 weeks or until the ANC has reached 10,000/mm³ following the expected chemotherapy-induced neutrophil nadir. Daily dosing of tbo-filgrastim should continue until the expected neutrophil nadir is passed and the ANC has recovered to the normal range; in clinical trials tbo-filgrastim was given for up to 14 days or until and ANC > 10,000 x 10⁶/L after the nadir. The duration of filgrastim and filgrastim biosimilar therapy needed to attenuate chemotherapy-induced neutropenia may be dependent on the myelosuppressive capabilities of the chemotherapy regimen being used. Filgrastim or its biosimilars should be discontinued if the ANC exceeds 10,000/mm³ after the expected chemotherapy-induced neutrophil nadir.

The first dose of filgrastim or its biosimilars for patients receiving BMT should be administered at least 24 hours after chemotherapy treatment and at least 24 hours after bone marrow infusion. **Eflapegrastim-xnst**, pegfilgrastim, and pegfilgrastim biosimilars should not be administered in the period between 14 days before and 24 hours after administration of cytotoxic chemotherapy.

The filgrastim-aafi, filgrastim-ayow, and filgrastim-sndz single-dose prefilled syringe are not designed to allow for direct administration of doses < 0.3 mL. The spring-mechanism interferes with the visibility of the graduation markings for doses < 0.3 mL. As a result, patient administration of doses < 0.3 mL is not recommended.

Eflapegrastim-xnst, pegfilgrastim, pegfilgrastim-apgf, pegfilgrastim-cbqv, pegfilgrastim-jmdb, pegfilgrastim-bmez, **pegfilgrastim-fpgk**, and **pegfilgrastim-pbbk** prefilled syringes are not designed to allow for direct administration of doses < 0.6 mL as they do not bear the necessary graduation marks. Direct administration to patients needing doses < 0.6 mL is not recommended due to potential dosing errors. Pediatric dosage of 0.1 mg/kg should be used for those weighing < 10 kg.

The prefilled syringe co-packaged with pegfilgrastim (Neulasta Onpro) kit should only be used with the on-body injector as it contains additional solution to compensate for liquid loss during delivery through the on-body injector. If the prefilled syringe co-packaged in the Neulasta Onpro kit is used for manual subcutaneous injection, overdose will occur. Likewise, if the manual single-use prefilled syringe is used with the on-body injector, the patient will be under-dosed. The on-body injector should not be used with any other drugs, and it is not recommended for patients with H-ARS.

An HCP must fill the on-body injector with pegfilgrastim using the prefilled syringe and apply the injector to the patient's intact, non-irritated skin (abdomen or back of arm). The back of the arm may only be used if there is a caregiver available to monitor the status of the on-body injector. An HCP may initiate administration with the on-body injector on the same day as the administration of chemotherapy, as long as the injector delivers pegfilgrastim no less than 24 hours after administration of chemotherapy. Approximately 27 hours after the on-body injector is applied to the patient's skin, pegfilgrastim will be delivered over approximately 45 minutes.

Like the single-use manual syringes, the single-use pegfilgrastim (Neulasta) delivery kits should be refrigerated until ready for use. Because the on-body injector for pegfilgrastim is at room temperature during use, the kit should not be held at room temperature for > 12 hours prior to use. The kit should be discarded if stored at room temperature > 12 hours. The prefilled syringe should be discarded if stored at room temperature > 48 hours. The injector should be kept dry for the last 3 hours prior to the

beginning of dose delivery in order to identify potential leaks. It is waterproof for up to 8 feet for 1 hour. It should not be reapplied if the on-body injector falls off.

For neutrophil recovery after chemotherapy in AML, if a second cycle of induction is needed, sargramostim should be administered approximately 4 days after the completion of chemotherapy if bone marrow is hypoplastic with < 5% blasts. Sargramostim should be continued until the ANC is > 1,500 cells/mm³ for 3 consecutive days; not to exceed 42 days. Sargramostim should not be administered within 24 hours preceding or following chemotherapy or radiation. For bone marrow transplantation failure or engraftment delay, another dose can be administered after 7 days off therapy if engraftment has not occurred; if engraftment still does not occur, a third course of 500 mcg/m²/day for 14 days can be used after another 7 days off therapy; additional dose escalation is unlikely to be beneficial.

Sargramostim treatment should be interrupted, or the dose reduced by half, if the ANC exceeds 20,000 cells/mm³ in order to avoid complications of excessive leukocytosis or if WBC is > 50,000 cells/mm³. If a severe reaction occurs, the sargramostim dose can be decreased by 50% or temporarily discontinued until the reaction goes away. If leukemic regrowth, disease progression, or blast cells appear, sargramostim should be discontinued.

CLINICAL TRIALS

Search Strategy

Articles were identified through searches performed on PubMed and review of information sent by manufacturers. Search strategy included the use of all drugs in this class for FDA-approved indications. Randomized, controlled, double-blind, comparative trials are considered the most relevant in this category. Studies included for analysis in the review were published in English, performed with human participants and randomly allocated participants to comparison groups. In addition, studies must contain clearly stated, predetermined outcome measure(s) of known or probable clinical importance, use data analysis techniques consistent with the study question and include follow-up (endpoint assessment) of at least 80% of participants entering the investigation. Despite some inherent bias found in all studies, including those sponsored and/or funded by pharmaceutical manufacturers, the studies in this therapeutic class review were determined to have results or conclusions that do not suggest systematic error in their experimental study design. While the potential influence of manufacturer sponsorship and/or funding must be considered, the studies in this review have also been evaluated for validity and importance.

Biosimilars must demonstrate that there are no clinically meaningful differences in safety or effectiveness from the reference product. Often biosimilars are compared to products that are not available in the US. In this therapeutic class review, only biosimilar clinical studies that meet the above search criteria and compare the biosimilar product to an FDA-approved reference product will be included.

Pegfilgrastim-apgf (Nyvepria) was compared to pegfilgrastim (Neulasta) for immunogenicity, safety, pharmacokinetics, and pharmacodynamics; no significant differences were found.¹²³

Pegfilgrastim-cbqv (Udenyca) was compared to pegfilgrastim (Neulasta) for immunogenicity, adverse reactions, safety, pharmacokinetics, and pharmacodynamics; no significant differences were found.¹²⁴

Pegfilgrastim-jmdb (Fulphila) was compared to European-sourced pegfilgrastim in a phase 3 trial, but not an FDA-approved product.¹²⁵ The FDA states approval is based on review of evidence that included

extensive structural and functional characterization, animal study data, human pharmacokinetic and pharmacodynamic data, clinical immunogenicity data, and other clinical safety and effectiveness data that demonstrates pegfilgrastim-jmdb (Fulphila) is biosimilar to pegfilgrastim (Neulasta).^{126,127}

Pegfilgrastim-bmez (Ziextenzo) was compared to pegfilgrastim (Neulasta) for pharmacokinetics, pharmacodynamics, safety, local tolerability, and immunogenicity; pegfilgrastim-bmez demonstrated similarity to Neulasta for these endpoints.¹²⁸

Pegfilgrastim-fpgk (Stimufend) was compared to pegfilgrastim (Neulasta) for pharmacokinetics, pharmacodynamics, safety, potency, purity, and immunogenicity; pegfilgrastim-fpgk was found to be similar to Neulasta for these endpoints with no clinically meaningful differences.¹²⁹

Pegfilgrastim-pbbk (Fylnetra) was compared to pegfilgrastim (Neulasta) for pharmacokinetics, pharmacodynamics, safety, local tolerability, and immunogenicity; pegfilgrastim-pbbk was found to be similar to Neulasta for these endpoints with no clinically meaningful differences.¹³⁰

Three open-label studies compared filgrastim-aafi (Nivestym) and US-sourced filgrastim (Neupogen). No significant difference was found between the products regarding pharmacokinetics, pharmacodynamics, safety, and immunogenicity.^{131,132,133}

eflapegrastim-xnst (Rovedon) and pegfilgrastim (Neulasta)

Two randomized, open-label, multicenter, active-controlled, non-inferiority studies of similar design compared the efficacy and safety of fixed doses of eflapegrastim to pegfilgrastim in 643 adult patients with early-stage breast cancer receiving up to 4 cycles of standard docetaxel plus cyclophosphamide (TC) chemotherapy.^{134,135} The primary endpoint was duration of severe neutropenia (DSN) ($ANC < 0.5 \times 10^9/L$) during cycle 1 of TC. Patients received a fixed dose of eflapegrastim 13.2 mg/0.6 mL (3.6 mg G-CSF) or pegfilgrastim 6 mg/0.6 mL (6 mg G-CSF). Study 1 enrolled 406 patients (196 received eflapegrastim, 210 received pegfilgrastim), and at cycle 1, 16% ($n = 31$) of patients receiving eflapegrastim and 24% ($n = 51$) of patients receiving pegfilgrastim had severe neutropenia, showing a relative risk decrease of 35% ($p=0.034$) with eflapegrastim and a difference in mean DSN of -0.148 day, thereby meeting the primary endpoint ($p<0.0001$). Noninferiority was observed at each of the 4 treatment cycles ($p<0.0001$, all cycles), and the study met all secondary endpoints. Study 2 included 237 patients (118 received eflapegrastim, 119 received pegfilgrastim). At cycle 1, 20.3% ($n = 24$) of patients receiving eflapegrastim and 23.5% ($n = 28$) of patients receiving pegfilgrastim had severe neutropenia ($ANC < 0.5 \times 10^9/L$). The primary endpoint, duration of severe neutropenia in patients receiving eflapegrastim, at a difference in mean DSN of -0.074 days, was noninferior to those receiving pegfilgrastim ($p<0.0001$). Noninferiority was observed at each of the 4 treatment cycles ($p<0.0001$, all cycles). Secondary endpoints included time-to-ANC recovery, occurrence of febrile neutropenia, occurrence of neutropenic complications, relative dose intensity, and safety; the study met all secondary endpoints and demonstrated noninferior efficacy of eflapegrastim in comparison to pegfilgrastim. Reported adverse events matched those reported in patients receiving TC and both eflapegrastim and pegfilgrastim were shown to have comparable safety profiles.

filgrastim (Neupogen) and pegfilgrastim (Neulasta)

Pegfilgrastim 6 mg SC once per chemotherapy cycle (every 21 days) and filgrastim 5 mcg/kg SC daily were compared in 157 patients with breast cancer (stage II–IV).¹³⁶ The placebo and filgrastim injections were given daily until the ANC was $> 10 \times 10^9/L$ or a total of 14 days. This randomized, double-blind,

multicenter study evaluated patients for duration of grade 4 neutropenia, depth of neutrophil nadir, incidence of febrile neutropenia, time to neutrophil recovery, and safety data. Patients received doxorubicin 60 mg/m² and docetaxel 75 mg/m² chemotherapy. A total of 152 patients were evaluated for efficacy. Pegfilgrastim and filgrastim had similar efficacy for all efficacy measures for all cycles. The incidence of severe neutropenia was 83% with filgrastim therapy and 84% with pegfilgrastim during cycle 1. The duration of severe neutropenia during cycle 1 was 1.6 days with filgrastim and 1.8 days with pegfilgrastim. The incidence of febrile neutropenia was 20% with filgrastim and 13% with pegfilgrastim. The median time of recovery to an ANC of > 2 x 10⁹ mm³ in all cycles for both treatment groups was 9 days. The most frequently reported adverse effect was bone pain (37% pegfilgrastim; 42% filgrastim).

A randomized, double-blind, non-inferiority study compared the safety and efficacy of filgrastim and pegfilgrastim in 310 patients with advanced breast cancer (stage II-IV) receiving chemotherapy.¹³⁷ Patients were randomized to filgrastim 5 mcg/kg body weight once daily or pegfilgrastim 100 mcg/kg body weight once every 21-day chemotherapy cycle plus placebo injection once daily. G-CSF therapy began 24 hours post-chemotherapy. Patients received doxorubicin 60 mg/m² and docetaxel 75 mg/m² for up to 4 cycles. Daily injections of filgrastim or placebo were given until the ANC was 10 x 10⁹/L or a total of 14 days. The primary outcome was duration of grade 4 neutropenia (ANC < 500 in cycle 1); the percentage of patients experiencing grade 4 neutropenia in cycle 1 was 79% and 77% for filgrastim and pegfilgrastim, respectively (p>0.5). The mean duration of grade 4 neutropenia for cycle 1 was 1.8 and 1.7 days for filgrastim and pegfilgrastim, respectively (p>0.5). For subsequent cycles (2 through 4), the mean duration of severe neutropenia for cycles 2, 3, and 4 were 0.7, 0.6, and 0.9 days for pegfilgrastim and 1.1, 1.2, and 1.3 days for filgrastim (p≤0.001 cycles 2 and 3, p=0.019 cycle 4). The incidence of febrile neutropenia (defined as fever > 38.2° C with ANC < 0.5 x 10⁹/L) was 12% with filgrastim and 7% with pegfilgrastim during the first cycle. Incidence of febrile neutropenia over the entire study period was 18% with filgrastim and 9% with pegfilgrastim (p=0.029). Time to ANC recovery was 9.7 days with filgrastim and 9.3 days with pegfilgrastim (95% confidence interval [CI], -0.88 to 0.08). Adverse event profiles were similar in both groups. The pegfilgrastim dose is different from the current FDA-approved labeling dose of 6 mg as a single dose.

filgrastim (Neupogen) and sargramostim (Leukine)

A randomized, double-blind, multicenter study compared sargramostim and filgrastim in the treatment of chemotherapy-induced myelosuppression in 181 afebrile cancer patients with ANC levels < 500/μL.¹³⁸ Patients received daily SC injections of either agent until ANC levels reached at least 1,500/μL. There was no statistical difference between treatment groups in the mean number of days to reach an ANC of 500/μL, but the mean number of days to reach ANC levels of 1,000/μL and 1,500/μL was approximately 1 day less in patients receiving filgrastim. Fewer patients in the sargramostim arm were hospitalized, and they had a shorter mean length of hospitalization, mean duration of fever, and mean duration of intravenous antibiotic therapy compared with patients who received filgrastim. Both growth factors were well tolerated. Sargramostim and filgrastim have comparable efficacy and tolerability in the treatment of standard-dose chemotherapy-induced myelosuppression in community practice.

A randomized, double-blind, multicenter study in 137 cancer patients receiving myelosuppressive chemotherapy compared the tolerability of sargramostim and filgrastim in the prevention or treatment of chemotherapy-induced neutropenia.¹³⁹ Patients received sargramostim SC 300 mcg daily or filgrastim SC 480 mcg daily starting 1 to 2 days after completion of chemotherapy. The drugs were given

prophylactically to 82% of patients within 48 hours; the other patients (18%) received the drugs when the ANC decreased to < 500/mL. No statistically significant differences in the incidence or severity of adverse events were detected with the exception of a slightly higher incidence of grade 1 fever (< 38.1 degrees C) with sargramostim. The study was not designed to evaluate efficacy, but it was noted that there were no statistically significant differences between treatment groups in total days of growth factor therapy, days of hospitalization, or days of intravenous antibiotics during the treatment period.

filgrastim-sndz (Zarxio) and filgrastim (Neupogen)

The efficacy and safety of filgrastim-sndz were evaluated in a multicenter, phase 3, randomized, double-blind, active-control, noninferiority study in 218 adult, female breast cancer patients receiving adjuvant myelosuppressive chemotherapy (TAC: docetaxel, doxorubicin, cyclophosphamide).¹⁴⁰ A notable inclusion criterion was adequate bone marrow function, while exclusion criteria included: history of myelogenous leukemia, myelodysplastic syndrome, or concomitant sickle cell disease; concurrent or prior radiotherapy within 4 weeks of randomization; and use of prophylactic antibiotics, prior chemotherapy, or anticancer treatment of breast cancer, or previous G-CSF therapy. Patients were randomized 1:1:1:1 to 4 treatment arms prior to 6 cycles of chemotherapy: (1) filgrastim-sndz, (2) reference filgrastim, (3) alternating filgrastim-sndz and reference filgrastim each cycle, or (4) alternating reference filgrastim and filgrastim-sndz each cycle. Chemotherapy was administered on day 1 of each cycle and given every 3 weeks for 6 cycles. The assigned filgrastim agent (5 mcg/kg/day) was administered daily as a SC injection beginning on day 2 of each cycle until the ANC recovered to 10,000/mm³ following the nadir or for a maximum of 14 days. The primary endpoint was duration of severe neutropenia, defined as a difference in mean DSN (number of consecutive days from ANC < 500/mm³ to ≥ 500/mm³) between the 2 filgrastim agents during cycle 1. Filgrastim-sndz was defined as noninferior if the lower 97.5% CI for the DSN mean difference in the per-protocol population was larger than -1 day. The mean DSN for the per-protocol population in cycle 1 was 1.17 days (95% CI, 0.95 to 1.39) in the biosimilar group (n=101) and 1.2 days (95% CI, 1 to 1.4) in the reference arm (n=103). The DSN mean difference was 0.04 days, and the lower 97.5% CI was -0.26 days. Thus, filgrastim-sndz met the noninferiority requirement. Results in the full analysis set for cycle 1 were similar to the per-protocol analysis. The authors found no clinically meaningful differences between the 2 agents across all cycles in secondary efficacy measures, including incidence of febrile neutropenia (FN), hospitalization secondary to FN, infection incidence, and extent and timing of nadir.

META-ANALYSIS

A meta-analysis evaluated a total of 5 head-to-head studies comparing pegfilgrastim and filgrastim for reducing chemotherapy-induced neutropenia among a total of 617 patients with solid tumors and malignant lymphomas.¹⁴¹ Studies included used the approved doses as indicated in the package insert. Although only 1 study had a statistically significant difference in febrile neutropenia reductions favoring pegfilgrastim over filgrastim (relative risk [RR] reduction of 50%; p=0.027), the pooled RR showed a statistically significant favorable result for pegfilgrastim (RR, 0.64; 95% CI, 0.43 to 0.97). Grade 4 neutropenia rates (for cycle 1: RR, 0.99; 95% CI, 0.91 to 1.08; cycle 2: RR, 0.88; 95% CI, 0.7 to 1.11; cycle 3: RR, 0.80; 95% CI, 0.47 to 1.36; cycle 4: RR, 0.9; 95% CI, 0.71 to 1.13), time to ANC (95% CI, -0.34 to 0.56), and incidence of bone pain (RR, 0.95; 95% CI, 0.76 to 1.19) were similar between the 2 G-CSFs. A single dose of pegfilgrastim performed better than a median of 10 to 14 days of filgrastim in reducing febrile neutropenia rates for patients undergoing myelosuppressive chemotherapy.

SUMMARY

Myelosuppressive chemotherapy can induce neutropenia and febrile neutropenia, which can lead to increased healthcare costs and poor patient health outcomes. Clinical trials have shown that prophylactic use of colony stimulating factors (CSFs) promotes the recovery of neutrophils following myelosuppressive chemotherapy and decreases the likelihood of neutropenic complications. Limited comparative data suggest that filgrastim (Neupogen) and pegfilgrastim (Neulasta) have similar efficacy, tolerability, and adverse drug reaction profiles. However, though data are limited, pegfilgrastim may have a slightly higher rate of reducing febrile neutropenia compared to filgrastim. In comparison to other products, the administration frequency of **eflapegrastim-xnst (Rolvedon)**, pegfilgrastim, and pegfilgrastim biosimilars may be viewed as more favorable since these products only require a single subcutaneous injection per chemotherapy cycle, whereas filgrastim, filgrastim-aafi (Nivestym), filgrastim-ayow (Releuko), filgrastim-sndz (Zarxio), tbo-filgrastim (Granix), and sargramostim (Leukine) administration requires daily subcutaneous injection. Additionally, the pegfilgrastim delivery kit (Neulasta Onpro) offers an alternative delivery option with the on-body injector. Several biosimilars to the originator products, filgrastim (filgrastim-aafi, filgrastim-ayow, and filgrastim-sndz) and pegfilgrastim (pegfilgrastim-apgf, pegfilgrastim-cbqv, pegfilgrastim-bmez, pegfilgrastim-jmdb, **pegfilgrastim-fpgk**, and **pegfilgrastim-pbbk**), are now available; tbo-filgrastim was not approved as a biosimilar. Further clinical trials comparing clinical activity, toxicity, cost-effectiveness, and long-term outcomes of these products are warranted since head-to-head trials are extremely limited.

The **v1.2023** NCCN practice guidelines for hematologic growth factors indicate filgrastim, tbo-filgrastim, and pegfilgrastim have a category 1 recommendation for prophylactic use against febrile neutropenia. The guidelines advise biosimilars are a reasonable substitute for filgrastim and pegfilgrastim. Due to the recent approvals, pegfilgrastim-apgf (Nyvepria), filgrastim-ayow (Releuko), **pegfilgrastim-fpgk (Stimufend)**, and **pegfilgrastim-pbbk (Fynetra)** are not currently addressed by NCCN in this guidance. Sargramostim is used for treatment but no longer recommended for prophylaxis. The guidelines also acknowledge that there are insufficient head-to-head comparative studies on the clinical benefits of G-CSFs and GM-CSFs. A delivery device for pegfilgrastim is also available that has impacted NCCN guidelines for pegfilgrastim use, eliminating same day pegfilgrastim/chemotherapy recommendations.

The NCCN guidelines also address the use of biosimilars. The guidelines note that biosimilars are biological products that are highly similar to their FDA-approved originator product with very small, clinically inactive differences but have no difference in efficacy, safety, or purity. The guidelines state that if overall safety and efficacy are equivalent, biosimilars may be substituted for the originator product and may provide opportunities for cost containment. While the FDA currently does not consider any biosimilar for filgrastim or pegfilgrastim to be interchangeable (substitutable without physician involvement) with its originator product, NCCN states limited data suggest that patients can alternate between the originator product and the biosimilar without any clinically meaningful differences regarding efficacy or safety.

To conclude, the NCCN guidelines recommend clinicians assess the patient and treatment-related risk factors for the development of neutropenic complications, and clinicians should use independent judgment when considering appropriate therapy.

REFERENCES

- 1 Rolvedon [package insert]. Irvine, CA; Spectrum; September 2022.
- 2 Neupogen [package insert]. Thousand Oaks, CA; Amgen; February 2021.
- 3 Nivestym [package insert]. New York, NY; Pfizer; April 2021.
- 4 Releuko [package insert]. Bridgewater, NJ; Amneal; April 2022.
- 5 Zarxio [package insert]. Princeton, NJ; Sandoz; August 2022.
- 6 Neulasta [package insert]. Thousand Oaks, CA; Amgen; February 2021.
- 7 Nyvepria [package insert]. New York, NY; Pfizer; October 2021.
- 8 Udenyca [package insert]. Redwood City, CA; Coherus; November 2022.
- 9 Fulphila [package insert]. Morgantown, WV; Mylan; October 2021.
- 10 Ziextenzo [package insert]. Princeton, NJ; Sandoz; March 2021.
- 11 Stimufend [package insert]. Lake Zurich, IL; Fresenius Kabi; September 2022.
- 12 Fylnetra [package insert]. Bridgewater, NJ; Amneal; May 2022.
- 13 Leukine [package insert]. Bridgewater, NJ; Sanofi-Aventis; May 2022.
- 14 Granix [package insert]. North Wales, PA; Teva; November 2019.
- 15 The National Comprehensive Cancer Network practice guidelines for Hematopoietic Growth Factors v1.2023. Available at: https://www.nccn.org/professionals/physician_gls/default.aspx. Accessed February 8, 2023.
- 16 The National Comprehensive Cancer Network practice guidelines for Hematopoietic Growth Factors v1.2023. Available at: https://www.nccn.org/professionals/physician_gls/default.aspx. Accessed February 8, 2023.
- 17 The National Comprehensive Cancer Network practice guidelines for Hematopoietic Growth Factors v1.2023. Available at: https://www.nccn.org/professionals/physician_gls/default.aspx. Accessed February 8, 2023.
- 18 Cosela [package insert]. Durham, NC; G1 Therapeutics; February 2021.
- 19 The National Comprehensive Cancer Network practice guidelines for Hematopoietic Growth Factors v1.2023. Available at: https://www.nccn.org/professionals/physician_gls/default.aspx. Accessed February 8, 2023.
- 20 Cosela [package insert]. Durham, NC; G1 Therapeutics; February 2021.
- 21 The National Comprehensive Cancer Network practice guidelines for Hematopoietic Growth Factors v1.2023. Available at: https://www.nccn.org/professionals/physician_gls/default.aspx. Accessed February 8, 2023.
- 22 The National Comprehensive Cancer Network practice guidelines for Hematopoietic Growth Factors v1.2023. Available at: https://www.nccn.org/professionals/physician_gls/default.aspx. Accessed February 8, 2023.
- 23 The National Comprehensive Cancer Network practice guidelines for Hematopoietic Growth Factors v1.2023. Available at: https://www.nccn.org/professionals/physician_gls/default.aspx. Accessed February 8, 2023.
- 24 Smith TJ, Bohlke K, Lyman GH, et. al. Recommendations for the use of WBC growth factors: American Society of Clinical Oncology Clinical Practice Guideline Update. 2015. Available at: <https://www.asco.org/research-guidelines/quality-guidelines/guidelines>. Accessed February 8, 2023.
- 25 Rolvedon [package insert]. Irvine, CA; Spectrum; September 2022.
- 26 Neupogen [package insert]. Thousand Oaks, CA; Amgen; February 2021.
- 27 Neulasta [package insert]. Thousand Oaks, CA; Amgen; February 2021.
- 28 Releuko [package insert]. Bridgewater, NJ; Amneal; April 2022.
- 29 Leukine [package insert]. Bridgewater, NJ; Sanofi-Aventis; May 2022.
- 30 Granix [package insert]. North Wales, PA; Teva; November 2019.
- 31 Zarxio [package insert]. Princeton, NJ; Sandoz; August 2022.
- 32 Fulphila [package insert]. Morgantown, WV; Mylan; October 2021.
- 33 Nivestym [package insert]. New York, NY; Pfizer; April 2021.
- 34 Nyvepria [package insert]. New York, NY; Pfizer; October 2021.
- 35 Udenyca [package insert]. Redwood City, CA; Coherus; November 2022.
- 36 Stimufend [package insert]. Lake Zurich, IL; Fresenius Kabi; September 2022.
- 37 Ziextenzo [package insert]. Princeton, NJ; Sandoz; March 2021.
- 38 Fylnetra [package insert]. Bridgewater, NJ; Amneal; May 2022.
- 39 Rolvedon [package insert]. Irvine, CA; Spectrum; September 2022.
- 40 Neupogen [package insert]. Thousand Oaks, CA; Amgen; February 2021.
- 41 Zarxio [package insert]. Princeton, NJ; Sandoz; August 2022.
- 42 Releuko [package insert]. Bridgewater, NJ; Amneal; April 2022.
- 43 Neulasta [package insert]. Thousand Oaks, CA; Amgen; February 2021.
- 44 Leukine [package insert]. Bridgewater, NJ; Sanofi-Aventis; May 2022.
- 45 Granix [package insert]. North Wales, PA; Teva; November 2019.
- 46 Fulphila [package insert]. Morgantown, WV; Mylan; October 2021.
- 47 Nivestym [package insert]. New York, NY; Pfizer; April 2021.
- 48 Nyvepria [package insert]. New York, NY; Pfizer; October 2021.
- 49 Udenyca [package insert]. Redwood City, CA; Coherus; November 2022.
- 50 Stimufend [package insert]. Lake Zurich, IL; Fresenius Kabi; September 2022.
- 51 Ziextenzo [package insert]. Princeton, NJ; Sandoz; March 2021.
- 52 Fylnetra [package insert]. Bridgewater, NJ; Amneal; May 2022.
- 53 Rolvedon [package insert]. Irvine, CA; Spectrum; September 2022.

54 Neupogen [package insert]. Thousand Oaks, CA; Amgen; February 2021.
55 Neulasta [package insert]. Thousand Oaks, CA; Amgen; February 2021.
56 Releuko [package insert]. Bridgewater, NJ; Amneal; April 2022.
57 Leukine [package insert]. Bridgewater, NJ; Sanofi-Aventis; May 2022.
58 Granix [package insert]. North Wales, PA; Teva; November 2019.
59 Zarxio [package insert]. Princeton, NJ; Sandoz; August 2022.
60 Fulphila [package insert]. Morgantown, WV; Mylan; October 2021.
61 Nivestym [package insert]. New York, NY; Pfizer; April 2021.
62 Nyvepria [package insert]. New York, NY; Pfizer; October 2021.
63 Udenyca [package insert]. Redwood City, CA; Coherus; November 2022.
64 Stimufend [package insert]. Lake Zurich, IL; Fresenius Kabi; September 2022.
65 Ziextenzo [package insert]. Princeton, NJ; Sandoz; March 2021.
66 Fylnetra [package insert]. Bridgewater, NJ; Amneal; May 2022.
67 Rolvedon [package insert]. Irvine, CA; Spectrum; September 2022.
68 Neupogen [package insert]. Thousand Oaks, CA; Amgen; February 2021.
69 Neulasta [package insert]. Thousand Oaks, CA; Amgen; February 2021.
70 Releuko [package insert]. Bridgewater, NJ; Amneal; April 2022.
71 Leukine [package insert]. Bridgewater, NJ; Sanofi-Aventis; May 2022.
72 Granix [package insert]. North Wales, PA; Teva; November 2019.
73 Zarxio [package insert]. Princeton, NJ; Sandoz; August 2022.
74 Fulphila [package insert]. Morgantown, WV; Mylan; October 2021.
75 Nivestym [package insert]. New York, NY; Pfizer; April 2021.
76 Nyvepria [package insert]. New York, NY; Pfizer; October 2021.
77 Udenyca [package insert]. Redwood City, CA; Coherus; November 2022.
78 Stimufend [package insert]. Lake Zurich, IL; Fresenius Kabi; September 2022.
79 Ziextenzo [package insert]. Princeton, NJ; Sandoz; March 2021.
80 Fylnetra [package insert]. Bridgewater, NJ; Amneal; May 2022.
81 Rolvedon [package insert]. Irvine, CA; Spectrum; September 2022.
82 Neupogen [package insert]. Thousand Oaks, CA; Amgen; February 2021.
83 Zarxio [package insert]. Princeton, NJ; Sandoz; August 2022.
84 Releuko [package insert]. Bridgewater, NJ; Amneal; April 2022.
85 Neulasta [package insert]. Thousand Oaks, CA; Amgen; February 2021.
86 Leukine [package insert]. Bridgewater, NJ; Sanofi-Aventis; May 2022.
87 Granix [package insert]. North Wales, PA; Teva; November 2019.
88 Fulphila [package insert]. Morgantown, WV; Mylan; October 2021.
89 Nivestym [package insert]. New York, NY; Pfizer; April 2021.
90 Nyvepria [package insert]. New York, NY; Pfizer; October 2021.
91 Udenyca [package insert]. Redwood City, CA; Coherus; November 2022.
92 Stimufend [package insert]. Lake Zurich, IL; Fresenius Kabi; September 2022.
93 Ziextenzo [package insert]. Princeton, NJ; Sandoz; March 2021.
94 Fylnetra [package insert]. Bridgewater, NJ; Amneal; May 2022.
95 Rolvedon [package insert]. Irvine, CA; Spectrum; September 2022.
96 Neupogen [package insert]. Thousand Oaks, CA; Amgen; February 2021.
97 Neulasta [package insert]. Thousand Oaks, CA; Amgen; February 2021.
98 Releuko [package insert]. Bridgewater, NJ; Amneal; April 2022.
99 Leukine [package insert]. Bridgewater, NJ; Sanofi-Aventis; May 2022.
100 Granix [package insert]. North Wales, PA; Teva; November 2019.
101 Zarxio [package insert]. Princeton, NJ; Sandoz; August 2022.
102 Fulphila [package insert]. Morgantown, WV; Mylan; October 2021.
103 Nivestym [package insert]. New York, NY; Pfizer; April 2021.
104 Nyvepria [package insert]. New York, NY; Pfizer; October 2021.
105 Udenyca [package insert]. Redwood City, CA; Coherus; November 2022.
106 Stimufend [package insert]. Lake Zurich, IL; Fresenius Kabi; September 2022.
107 Ziextenzo [package insert]. Princeton, NJ; Sandoz; March 2021.
108 Fylnetra [package insert]. Bridgewater, NJ; Amneal; May 2022.
109 Rolvedon [package insert]. Irvine, CA; Spectrum; September 2022.
110 Neupogen [package insert]. Thousand Oaks, CA; Amgen; February 2021.
111 Zarxio [package insert]. Princeton, NJ; Sandoz; August 2022.
112 Releuko [package insert]. Bridgewater, NJ; Amneal; April 2022.
113 Neulasta [package insert]. Thousand Oaks, CA; Amgen; February 2021.
114 Leukine [package insert]. Bridgewater, NJ; Sanofi-Aventis; May 2022.
115 Granix [package insert]. North Wales, PA; Teva; November 2019.
116 Fulphila [package insert]. Morgantown, WV; Mylan; October 2021.
117 Nivestym [package insert]. New York, NY; Pfizer; April 2021.
118 Nyvepria [package insert]. New York, NY; Pfizer; October 2021.

-
- 119 Udenyca [package insert]. Redwood City, CA; Coherus; November 2022.
- 120 Stimufend [package insert]. Lake Zurich, IL; Fresenius Kabi; September 2022.
- 121 Ziextenzo [package insert]. Princeton, NJ; Sandoz; March 2021.
- 122 Fynetra [package insert]. Bridgewater, NJ; Amneal; May 2022.
- 123 Center for Drug Evaluation and Research. Nyvepria. Multi-Discipline Review. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/nda/2020/761111Orig1s000MultidisciplineR.pdf. Accessed February 8, 2023.
- 124 Center for Drug Evaluation and Research. Udenyca Summary Review. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/nda/2018/761039Orig1s000MedR.pdf. Accessed February 8, 2023.
- 125 Waller CF, Blakeley C, Pennella E, et al. Phase 3 efficacy and safety trial of proposed pegfilgrastim biosimilar MYL-1401H vs EU-Neulasta in the prophylaxis of chemotherapy-induced neutropenia. *Annals of Oncology*. 2016; 27(suppl 6):vi497-vi521.
- 126 FDA approves first biosimilar to Neulasta to help reduce the risk of infection during cancer treatment. Available at: <https://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm609805.htm>. Accessed February 8, 2023.
- 127 Waller CF, Ranganna GM, Pennella EJ, et al. Randomized phase 3 efficacy and safety trial of proposed pegfilgrastim biosimilar MYL-1401H in the prophylactic treatment of chemotherapy-induced neutropenia. *Ann Hemtol*. 2019; 98(5):1217-1224. DOI: 10.1007/s00277-019-03639-5.
- 128 Nakov R, Schussler S, Schier-Mumzhiu S, et al. A large multi-center, randomized, double-blind, crossover study in healthy volunteers to compare pharmacokinetics and pharmacodynamics of a proposed biosimilar pegfilgrastim with EU and US reference pegfilgrastim: Methodological approach [published online ahead of print May 26, 2019]. *Annals of Oncology*. 2019; 37(suppl 15). DOI: 10.1200/JCO.2019.37.15_suppl.e23118.
- 129 Center for Drug Evaluation and Research. Stimufend. Multi-Discipline Review. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/nda/2022/761173Orig1s000MultidisciplineR.pdf. Accessed February 9, 2023.
- 130 Center for Drug Evaluation and Research. Fynetra. Multi-Discipline Review. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/nda/2022/761084Orig1s000MultidisciplineR.pdf. Accessed February 8, 2023.
- 131 Nivestym. Office of Clinical Pharmacology 251(k) biosimilar review. FDA Available at: <https://www.fda.gov/media/115386/download>. Accessed February 13, 2023.
- 132 FDA Clinical review. Filgrastim-aafi. Available at: <https://www.fda.gov/media/115382/download>. Accessed February 13, 2023.
- 133 Yao HM, Ottery FD, Borema T, et al. PF-06881893 (Nivestym™), a filgrastim biosimilar, versus US-Licensed filgrastim reference product (US-Neupogen®): pharmacokinetics, pharmacodynamics, immunogenicity, and safety of single or multiple subcutaneous doses in healthy volunteers. *BioDrugs*. 2019; 33(2): 207-220. DOI: 10.1007/s40259-019-00343-8.
- 134 Cobb PW, Moon YW, Mezei K, et al. A comparison of eflapegrastim to pegfilgrastim in the management of chemotherapy-induced neutropenia in patients with early-stage breast cancer undergoing cytotoxic chemotherapy (RECOVER): a phase 3 study. *Cancer Med*. 2020;9(17):6234-6243. doi:10.1002/cam4.3227.
- 135 Schwartzberg LS, Bhat G, Peguero J, et al. Eflapegrastim, a long-acting granulocyte-colony stimulating factor for the management of chemotherapy-induced neutropenia: results of a phase III trial. *Oncologist*. 2020;25(8):e1233-e1241. doi:10.1634/theoncologist.2020-0105.
- 136 Green MD, Koelbl H, Baselga J, et al for the International Pegfilgrastim 749 Study Group. A randomized double-blind multicenter phase III study of fixed-dose single-administration pegfilgrastim versus daily filgrastim in patients receiving myelosuppressive chemotherapy. *Ann Oncol*. 2003; 14(1):29-35.
- 137 Holmes FA, O'Shaughnessy JA, Vukelia S, et al. Blinded, randomized, multicenter study to evaluate single administration pegfilgrastim once per cycle versus daily filgrastim as an adjunct to chemotherapy in patients with high-risk stage II or stage III/IV breast cancer. *J Clin Oncol*. 2002; 20(3):727-731.
- 138 Beveridge RA, Miller JA, Kales AN, et al. A comparison of efficacy of sargramostim (yeast-derived RhuGM-CSF) and filgrastim (bacteria-derived RhuG-CSF) in the therapeutic setting of chemotherapy-induced myelosuppression. *Cancer Invest*. 1998; 16(6):366-373.
- 139 Beveridge RA, Miller JA, Kales AN, et al. Randomized trial comparing the tolerability of sargramostim (yeast-derived RhuGM-CSF) and filgrastim (bacteria-derived RhuG-CSF) in cancer patients receiving myelosuppressive chemotherapy. *Support Care Cancer*. 1997; 5(4):289-298.
- 140 Blackwell K, Semiglazoc C, Krasnozhan D, et al. Comparison of EP2006, a filgrastim biosimilar, to the reference: a phase III, randomized, double-blind clinical study in the prevention of severe neutropenia in patients with breast cancer receiving myelosuppressive chemotherapy. *Ann Oncol*. 2015; 26(9):1948-1953.
- 141 Pinto L, Liu Z, Doan Q, et al. Comparison of pegfilgrastim with filgrastim on febrile neutropenia, grade IV neutropenia and bone pain: a meta-analysis of randomized controlled trials. *Curr Med Res Opin*. 2007; 23(9):2283-2295.
-