

**Medication Audit Checklist** 

## Bupropion (Wellbutrin®, Budeprion®, Zyban®)

January 2023

## Audit Information

Reviewer:		
Drug:		
Audit #:		
Audit Date:		
Dose:		
Does this audit require a physician review? Y or N		

## **Patient Information**

Patient #:
Age:
Ordering Provider:
Admit Date:
Gender:
Attending Provider:

Indication		
This document lists only FDA- approved indications from the product labeling. The PEFC acknowledges that there are off- label indications for use that have supporting evidence for efficacy. If a medication is prescribed for an off-label indication, documentation in the patient chart is recommended.	Comments	Does this indication require a physician review?
<ul> <li>Seasonal Affective Disorder (24 hour extended release)</li> <li>Major Depressive Disorder</li> <li>Smoking cessation (12 hour extended release)</li> </ul>		

Contraindications	Comments	Does this contraindication require a physician review?
<ul> <li>Absolute</li> <li>Abrupt discontinuation of alcohol, benzodiazepines, barbiturates, antiepileptics</li> <li>Co-administration with an MAOI, including linezolid or IV methylene blue, or use within 14 days of discontinuing an MAOI</li> <li>Current or prior diagnosis of bulimia or anorexia nervosa</li> <li>Diagnosis of a seizure disorder or history of seizures</li> <li>History of anaphylactic reaction or similarly severe significant hypersensitivity to the medication prescribed</li> </ul>		
Relative <ul> <li>Pregnancy/nursing mothers</li> </ul>		

Patient Monitoring	Comments	Does this require a physician review?
<ul> <li>Baseline Tests:</li> <li>Blood pressure and heart rate</li> <li>CBC</li> <li>Hepatic and renal function panels</li> <li>Pregnancy test (females)</li> <li>Weight</li> </ul>		
<ul> <li>Ongoing Tests:</li> <li>Blood pressure and heart rate <ul> <li>as clinically indicated</li> <li>CBC - as clinically indicated</li> <li>ECG - as clinically indicated</li> <li>Hepatic function - as clinically indicated</li> <li>Emergence of suicidal ideation or behavior - as clinically indicated, particularly during the first two months of therapy/after dosage adjustments</li> <li>Neuropsychiatric reactions if using for smoking cessation</li> <li>Pregnancy test - as clinically indicated</li> <li>Renal function - as clinically indicated</li> </ul> </li> </ul>		