

Test	Ages	Risk Factors
Dyslipidemia	2 years and older	<p>Dyslipidemia testing is recommended for anyone at increased risk for dyslipidemia, including:</p> <p>Positive family history:</p> <ul style="list-style-type: none"> • Premature coronary artery disease • Parent with known dyslipidemia or TC >240 mg/dL (6.2 mmol/L) <p>Specific conditions:</p> <ul style="list-style-type: none"> • Diabetes mellitus and familial hypercholesterolemia associated with high or moderate risk for CVD • Significant tobacco smoke exposure • Hypertension <p>Overweight:</p> <ul style="list-style-type: none"> • Elevated body mass index (BMI) • Children between 2 and 8 years of age with a BMI ≥95th percentile • Older children with a BMI ≥85th percentile
Type 2 diabetes	10 years, or onset of puberty if earlier	<p>Type 2 diabetes testing is recommended for anyone at increased risk for type 2 diabetes, including: Overweight:</p> <ul style="list-style-type: none"> • BMI >85th percentile for age/gender • Weight for height >85th percentile • Weight >120% of ideal for height <p>Plus any two of the following: Positive family history:</p> <ul style="list-style-type: none"> • Type 2 diabetes in first- or second-degree relative <p>Race/ethnicity:</p> <ul style="list-style-type: none"> • American Indian • African American • Latino • Asian American • Pacific Islander <p>Signs of insulin resistance or:</p> <ul style="list-style-type: none"> • Acanthosis nigricans • Hypertension • Dyslipidemia • PCOS • Small for gestational age birth weight Maternal history of diabetes or GDM during the child's gestation



Hepatitis C	Any	<p>HCV testing is recommended for anyone at increased risk for HCV infection, including: Injected illegal drugs:</p> <ul style="list-style-type: none"> • Persons who have ever injected illegal drugs, including those who injected only once many years ago <p>Selected medical conditions:</p> <ul style="list-style-type: none"> • Patients who have ever received long-term hemodialysis treatment • All persons with HIV infection • Patients with signs or symptoms of liver disease (e.g., abnormal liver enzyme tests) <p>Children born to HCV-positive mothers (to avoid detecting maternal antibody, these children should not be tested before age 18 months)</p>
HIV	Any	<p>HIV testing is recommended for anyone at increased risk for HIV infection, including: Exposure or potential exposure:</p> <ul style="list-style-type: none"> • Use of intravenous drugs • Sexual activity or abuse <p>History of an STI</p> <p>Maternal-child transmission</p>

REFERENCES

Dyslipidemia

http://www.nhlbi.nih.gov/guidelines/cvd_ped/peds_guidelines_sum.pdf

<http://www.uptodate.com/contents/definition-and-screening-for-dyslipidemia-in-children>

Type 2 Diabetes

Standards of Medical Care in Diabetes-2012

http://care.diabetesjournals.org/content/35/Supplement_1/S11/T5.expansion.html

Hepatitis C

Hepatitis C FAQs for Health Professionals

<http://www.cdc.gov/hepatitis/hcv/hcvfaq.htm>

HIV

Adolescents and HIV Infection: The Pediatrician's Role in Promoting Routine Testing Committee on Pediatric AIDS Pediatrics 2011;128;1023; originally published online October 31, 2011; DOI: 10.1542/peds.2011-1761

<http://pediatrics.aappublications.org/content/128/5/1023.full.pdf>

