



## Medication Audit Checklist

### Trazodone (Desyrel®), Nefazodone (Serzone®)

January 2023

#### Audit Information

Reviewer:
Drug:
Audit #:
Audit Date:
Dose:
Does this audit require a physician review? Y or N

#### Patient Information

Patient #:
Age:
Ordering Provider:
Admit Date:
Gender:
Attending Provider:

<b>Indication</b>	<b>Comments</b>	<b>Does this indication require a physician review?</b>
<p>If a medication is prescribed for an off-label indication, documentation in the patient chart is recommended.</p>		
<p>Label:</p> <ul style="list-style-type: none"> <li>● Major Depressive Disorder</li> </ul> <p>Off-label:</p> <ul style="list-style-type: none"> <li>● Insomnia</li> </ul>		

<b>Contraindications</b>	<b>Comments</b>	<b>Does this contraindication require a physician review?</b>
<ul style="list-style-type: none"> <li>● Co-administration with an MAOI, including linezolid or IV methylene blue, or use within 14 days of discontinuing an MAOI</li> <li>● History of anaphylactic reaction or similarly severe significant hypersensitivity to trazodone or nefazodone</li> <li>● Trazodone: recovery phase of myocardial infarction</li> <li>● Nefazodone: history of liver injury due to previous nefazodone treatment</li> <li>● Nefazodone: concurrent therapy with carbamazepine, cisapride, terfenadine, astemizole, pimozone, or full doses of triazolam</li> </ul>		

<b>Patient Monitoring</b>	<b>Comments</b>	<b>Does this require a physician review?</b>
<p>Baseline Tests:</p> <ul style="list-style-type: none"> <li>● Pregnancy test (females)</li> <li>● CBC</li> <li>● ECG as clinically indicated</li> <li>● Liver function</li> </ul> <p>Ongoing Tests:</p> <ul style="list-style-type: none"> <li>● Pregnancy test (females) as clinically indicated</li> <li>● Trazodone: Liver function tests as clinically indicated</li> <li>● Nefazodone: Liver function tests at 1, 2, 4, 6 and 12 months, then annually and as clinically indicated. Discontinue if AST or ALT levels are 3 times (or greater) the upper limit of normal.</li> <li>● CBC as clinically indicated</li> <li>● ECG as clinically indicated</li> <li>● Monitor for emergence of suicidal ideation or behavior or worsening depression</li> </ul>		